

1982

An Evaluation of Three Feeding Schemes to Winter Replacement Heifers

L. B. Bruce

South Dakota State University

H. L. Miller


A. Dittman

Follow this and additional works at: http://openprairie.sdstate.edu/sd_cow-calf_1982

Recommended Citation

Bruce, L. B.; Miller, H. L.; and Dittman, A., "An Evaluation of Three Feeding Schemes to Winter Replacement Heifers" (1982). *South Dakota Cow-Calf Field Day Proceedings, 1982*. Paper 11.
http://openprairie.sdstate.edu/sd_cow-calf_1982/11

This Report is brought to you for free and open access by the Animal Science Reports at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in South Dakota Cow-Calf Field Day Proceedings, 1982 by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.



AN EVALUATION OF THREE FEEDING SCHEMES TO WINTER REPLACEMENT HEIFERS

L. B. Bruce, H. L. Miller and A. Dittman

Department of Animal and Range Sciences

COW-CALF 82-5

Summary

Three feeding schemes for wintering replacement heifers were evaluated in terms of cost and animal performance. Ninety-three Simmental-Angus crossbred heifers were divided into three groups of 31 head and fed the following diets: (1) .28 Mcal of net energy for gain per pound of feed, free choice, (2) .28 Mcal of net energy for gain per pound of feed, limit fed to 13 lb dry matter per head per day and (3) .36 Mcal of net energy for gain per pound of feed, limit fed to 13 lb of dry feed per head per day. The third diet (higher energy) was the best scheme, resulting in the best gains and the lowest feed costs per pound of gain.

Introduction

Many different feeding practices have been used to grow out replacement heifers. The goal for replacement heifers is to over winter at gains sufficiently high enough that they will be at 60% of mature body weight at breeding. This should be done as efficiently as possible and to meet all nutrient requirements.

This study was designed to evaluate three methods of growing replacements as to dollar efficiency and animal performance.

Methods

Ninety-three Simmental-Angus crossbred heifers were purchased at weaning and placed on trial at the James Valley Research and Extension Center at Redfield, South Dakota. The heifers were divided into two groups, light and heavy (avg wt of 510 lb and 600 lb, respectively). Each group was divided randomly into three groups, each receiving one of the following diets: (1) .28 Mcal of net energy for gain per pound of dry feed, free choice (low energy diet, free choice), (2) the same diet as in 1 but with intake limited to 13 lb of dry feed per head per day or (3) .36 Mcal of net energy for gain per pound of dry feed (high energy diet), intake limited to 13 lb of dry matter per head per day. Composition of the diets is shown in table 1. The cattle were weighed and treatments initiated February 4, 1982. The heifers were artificially inseminated in early June and dietary treatments were terminated at that time, resulting in a 144-day feeding period. Cattle were weighed every 28 days and feed intake was measured on a pen basis each day. Subsequently, cattle were placed on pasture at Cottonwood, South Dakota.

Analyses of data included calculation of weight gain, average daily gain, feed consumption and feed cost per treatment group. The reproductive performance of the heifers will be monitored in future years.

Table 1. Diet Composition^a, Energy Level and Daily Intake Level for Each of the Diets

	Low energy Free choice	High energy Limit fed	High energy Limit fed
Composition ^b			
Corn, shelled, %	65	77	65
Prairie hay, %	34	22	34
Protein supplement, %	1	1	1
Energy, NEg Mcal/lb	.28	.36	.28
Intake per head per day, lb	12.5	12.7	14.5

^a All diets included free access to mineralized salt.

^b All numbers are on a dry matter basis.

Results

Data representing total weight gain, average daily gain, total feed consumption, pounds of feed per pound of gain and feed cost are presented in table 2. The average daily gains and feed efficiencies were lower than expected in all groups because of a month of severe weather compounded by recurring water problems. The most satisfactory overall performance was observed for the high energy limit-fed group. They gained the most with the lowest feed cost per pound of gain. The animals in the low energy free choice group consumed the largest amount of feed and had the poorest feed efficiency, resulting in the highest total feed cost as well as feed cost per pound of gain. The low energy limit-fed group was more efficient than the free-choice group but had the lowest rate of gain. The limit-fed high energy group had the best total weight gain, average daily gain and feed conversion. They also were the most economical in feed cost per pound of gain. Limit feeding seems to decrease feed costs, but the energy level in the diet must be relatively high to sustain adequate gains.

Table 2. Feedlot Performance^a as Measured by Weight Gain, Average Daily Gain, Feed Consumption, Feed Efficiency and Feed Cost of Simmental-Angus Crossbred Replacement Heifers Fed 144 Days

	Low energy Free choice	Low energy Limit fed	High energy Limit fed
Total wt gain, lb	189	171	195
Avg daily gain, lb	1.31 ± .12	1.22 ± .04	1.39 ± .03
Total feed consumed, lb ^b	2099	1810	1839
Feed efficiency, lb feed per lb gain, dry matter basis	11.24 ± 1.22	10.64 ± .77	9.43 ± .07
Feed cost, \$			
Total ^c	73	63	65
Per lb of gain	.39	.37	.33

^a Values are pen averages for one individual.

^b Dry basis.

^c Based on corn at \$2.40 per bushel and prairie hay at \$68 per ton.