

1972

Dietary Protein, Fat and Copper Levels and Its Effect on Fat Composition of Turkeys

R. A. Nelson

South Dakota State University

E. Guenther

South Dakota State University

C. W. Carlson

South Dakota State University

Follow this and additional works at: http://openprairie.sdstate.edu/sd_poultry_1972

Recommended Citation

Nelson, R. A.; Guenther, E.; and Carlson, C. W., "Dietary Protein, Fat and Copper Levels and Its Effect on Fat Composition of Turkeys" (1972). *South Dakota Poultry Field Day Proceedings and Research Reports, 1972*. Paper 11.
http://openprairie.sdstate.edu/sd_poultry_1972/11

This Report is brought to you for free and open access by the Animal Science Reports at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in South Dakota Poultry Field Day Proceedings and Research Reports, 1972 by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

Dietary Protein, Fat and Copper Levels and Its Effect
on Fat Composition of Turkeys

R. A. Nelson¹, E. Guenther² and C. W. Carlson³

Reports suggesting a relationship between diet and blood cholesterol levels have caused people to become more concerned about the type of fat in their diets. However, there is little conclusive evidence that high blood cholesterol is indeed a major cause of heart attacks in this country. Previous work at this station has indicated that the diet has a great effect on the type of body fat in birds and that what already is highly unsaturated can be readily made still further unsaturated.

In this experiment, the effects of copper, fat and protein level on carcass composition were considered. Carcasses from the growth study reported here (A.S. Series 72-13) were made available for this work.

Adipose tissue collections were taken at slaughter from a random selection of ten birds from each treatment (five hens and five toms). Two methyl esters were made of each fat sample and these in turn were analyzed in replicate for fatty acid content by gas-liquid chromatography. The results obtained are shown in table 1. Copper supplementation in this study did not increase the extent of total fat unsaturation as had been reported for swine. There also seemed to be no consistent alterations in the individual fatty acid contents due to copper supplementation.

The 5% corn-oil additions caused a large increase in unsaturation, while the 5% yellow grease addition caused a slight decrease in body fat unsaturation. The large unsaturation increase in the corn-oil diets was due mainly to an approximate doubling of linoleic acid at the expense of the other fatty acids. Diets with the yellow grease supplement caused an increase in myristic, palmitoleic, stearic and oleic acid contents, along with a decrease in linoleic acid. Diets had no consistent effect on total fat content of muscle tissue.

¹ Graduate student.

² Instructor.

³ Professor and Leader, Poultry Research and Extension.

Table 1. Fat Composition as Influenced by Diet and Type of Turkey

Fatty acids	High protein + yellow grease	High protein + corn oil	Low protein	Low protein + corn oil	Low protein + yellow grease	Low protein + 120 ppm copper	Low protein + 240 ppm copper
<u>15 Week Hen Turkeys</u>							
Myristic, %	1.37	0.55	0.91	0.63	1.29	0.92	0.94
Palmitic, %	21.55	16.48	20.90	16.00	19.85	23.00	23.68
Palmitoleic, %	3.56	1.29	4.02	1.65	3.27	5.32	4.35
Stearic, %	10.45	6.75	9.09	6.53	10.27	7.97	9.61
Oleic, %	39.58	27.30	36.98	27.54	38.69	37.42	34.98
Linoleic, %	21.67	47.08	26.69	45.86	24.88	24.05	25.13
Linolenic, %	2.02	1.59	1.45	1.65	1.85	1.42	1.34
% Saturation	33.37	23.78	30.90	23.16	31.41	31.89	34.23
% Unsaturation	66.83	77.26	69.14	76.70	68.59	68.21	65.80
Cooked muscle fat on a moisture-free basis, %	12.7	13.9	21.0	26.9	15.8	24.0	14.8
<u>24 Week Tom Turkeys</u>							
Myristic, %	1.56	0.37	0.38	0.88	1.44	0.91	0.91
Palmitic, %	20.63	16.16	16.03	22.32	20.80	23.66	24.00
Palmitoleic, %	2.57	0.72	0.91	5.09	3.56	4.65	5.17
Stearic, %	12.98	6.45	7.06	9.44	11.23	9.46	9.59
Oleic, %	40.21	25.74	26.28	37.59	41.05	35.88	36.10
Linoleic, %	21.71	50.65	49.38	24.71	21.95	25.46	24.46
% Saturation	35.17	22.98	23.47	32.64	33.47	34.03	34.50
% Unsaturation	64.87	77.11	76.57	67.39	66.56	65.99	65.73
Cooked muscle fat on a moisture-free basis, %	16.6	23.6	18.3	17.0	17.8	15.6	18.8