

South Dakota State University

# Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

---

Electronic Theses and Dissertations

---

1972

## Influence of Weather On Movement and Habitat Use of Hen Pheasants During Brood Rearing

James M. Ruth

Follow this and additional works at: <https://openprairie.sdstate.edu/etd>



Part of the [Natural Resources and Conservation Commons](#)

---

### Recommended Citation

Ruth, James M., "Influence of Weather On Movement and Habitat Use of Hen Pheasants During Brood Rearing" (1972). *Electronic Theses and Dissertations*. 211.

<https://openprairie.sdstate.edu/etd/211>

This Thesis - Open Access is brought to you for free and open access by Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact [michael.biondo@sdstate.edu](mailto:michael.biondo@sdstate.edu).

INFLUENCE OF WEATHER ON MOVEMENT  
AND HABITAT USE OF HEN PHEASANTS  
DURING BROOD REARING

BY

JAMES M. RUTH

A thesis submitted  
in partial fulfillment of the requirements for the  
degree Master of Science, Major in  
Wildlife Biology, South Dakota  
State University

1972

INFLUENCE OF WEATHER ON MOVEMENT  
AND HABITAT USE OF HEN PHEASANTS  
DURING BROOD REARING

This thesis is approved as a creditable and independent investigation by a candidate for the degree, Master of Science, and is acceptable as meeting the thesis requirements for this degree, but without implying that the conclusions reached by the candidate are necessarily the conclusions of the major department.

Thesis Adviser

Date

Head, Department of Wildlife  
and Fisheries Sciences

Date

## ACKNOWLEDGEMENTS

I am very grateful to Dr. Donald Progulske for his assistance and guidance throughout this study and the preparation of this manuscript.

I would also like to thank Dr. W. Lee Tucker, who helped with the statistical analysis of data, and David Oschner who wrote the computer programs, for their contributions to this effort.

My thanks to Wayne Grundstrom and Robert Fedeler for their assistance in field work and especially to Lynn Hanson who collected the data used in the first two years of this study.

I would also like to thank the South Dakota Department of Game, Fish, and Parks for allowing me to trap and mark pheasants used in this study.

Finally, I would like to thank my wife, Karen. Her encouragement and help were invaluable throughout the study.

Financial support was provided by South Dakota Agricultural Experiment Station Project S-521.



## TABLE OF CONTENTS

INTRODUCTION . . . . .	1
DESCRIPTION OF STUDY AREA . . . . .	3
METHODS AND MATERIALS . . . . .	5
Capturing Birds . . . . .	5
Marking Systems . . . . .	5
Radio-tracking Systems . . . . .	6
Data Analysis . . . . .	7
RESULTS . . . . .	10
Capture Data . . . . .	10
Radio-Tracking Movement Data . . . . .	10
Location of Birds . . . . .	13
Habitat-Use Indices . . . . .	18
DISCUSSION AND CONCLUSIONS . . . . .	23
LITERATURE CITED . . . . .	24
APPENDIX . . . . .	27

LIST OF TABLES

TABLE	PAGE
1. General cover types on study area 1969, 1970, 1971 . . . . .	4
2. Summary of radio-tracking data . . . . .	11
3. Summary of brood data . . . . .	12
4. Percentage of locations in cover types, 1969 . . . . .	14
5. Percentage of locations in cover types, 1970 . . . . .	16
6. Percentage of locations in cover types, 1971 . . . . .	17
7. Cover-use index based on changes in barometric pressure . .	19
8. Cover-use index based on changes in wind velocity . . . . .	20
9. Cover-use index based on occurrence of precipitation . . .	22

## INTRODUCTION

Pheasants survive in a wide range of yearly weather extremes but can be limited by certain short-term weather factors (Graham and Hesterburg 1948). Study of short-term weather conditions can therefore be important in understanding pheasant survival.

Many studies have been done on yearly and seasonal effects of weather on pheasant production and survival. Bennett and Terrill (1940) reported on possible temperature factors in pheasant management in the north central area of the United States. Effects of weather on pheasant reproduction were studied by Buss and Swanson (1950) and Kozielsky et al. (1955). MacMullan and Eberhardt (1953) studied the tolerance of incubating eggs and young chicks to exposure. Labisky et al. (1964) discussed the influence of land use and weather on pheasant distribution.

A few studies have reported the daily effects of weather on movement and habitat use of pheasants. The ability of pheasants to withstand climatic elements was measured by Latham (1947). Restriction of pheasant movement by weather was reported by Shick (1952) and Weston (1954). Cover type use was related to weather by Grondahl (1953).

The previous studies were all done by visual observation. The development of radio-telemetry systems (Lefkunan et al. 1959, Marshall 1960, Cochrane and Lord 1963) and techniques (Marshall and Kupa 1963) have made possible the collection of more information on daily movement and habitat use. The use of computers to analyze these data has allowed more detailed studies to be done (Siniff and Pester 1965). Movement and behavior of hen pheasants during the nesting season were reported by

Kuck et al. (1970). Carter (1971) studied seasonal movement of pheasants. General habitat use and movement of hen pheasants during brood-rearing were studied by Hanson (1971).

Survival of broods may have as much influence on pheasant population as hatching success (Trautman and Dahlgren 1965). Since little is known about the influence of daily weather variations on hens during the brood-rearing period, the objective of this study was to determine the effect of selected weather conditions on the movement and habitat use of hen pheasants during the brood-rearing period.

## DESCRIPTION OF STUDY AREA

Field studies were conducted on private farmland (W  $\frac{1}{2}$  and SE  $\frac{1}{4}$  of Section 22 and SE  $\frac{1}{4}$  of Section 21, Township 110 N, Range 49 W) 3 miles east of Brookings, South Dakota.

The topography is flat to gently sloping. Soils are glacial in origin with a loess overlay (Westin et al. 1958).

Climate is continental. Annual precipitation is 20 inches, generally occurring as frontal precipitation falling at a slow rate over a long period of time. About 80 percent of the precipitation comes during the growing season. Temperatures range between -30 F and 110 F with average monthly summer temperatures varying between 49 F and 74 F. Wind velocity averages about 10 MPH from the south during the summer (U. S. Dept. Commerce 1969, 1970, 1971). Barometric pressure can be highly variable but averaged 29.94 inches, corrected to sea level, for the period from 1931-1960 (U. S. Dept. of Commerce, Climatic Atlas of the U. S. 1970). Weather data for this study were obtained from the South Dakota State University Experiment Station located 3 miles west of the study area (Appendix Table 1).

Agriculture is the major activity on the area, with corn and sown small grain the major crops. The proportions of cover types changed slightly during the 3 years of the study, with the greatest change the last year (Table 1). In 1971, 100 acres that had been retired under the U. S. D. A. Feed Grain Program were put into small grain production. Even with this change, brood-rearing cover remained approximately 60 percent of the area each year of the study.

Table 1. General cover types on study area 1969, 1970, 1971.

Cover Type	1969		1970		1971	
	Acres	Percent of Total	Acres	Percent of Total	Acres	Percent of Total
Corn	188	29	205	32	247	38
Small grain	158	25	155	24	202	32
Residual cover	122	19	126	20	30	5
Pasture	56	9	56	9	54	8
Summer fallow	41	6	34	5	44	7
Alfalfa	37	6	26	4	25	4
Treecrow and farmstead	17	3	17	3	17	3
Ditches	16	2	16	2	16	2
Spoil pits	4	1	5	1	5	1

## METHODS AND MATERIALS

### Capturing Birds

A pickup truck with floodlights (Labisky 1968) and a backpack generator (Drewien et al. 1967) were used to capture birds at night for marking. The backpack generator was used most because of its portability and usefulness for searching small areas of roosting cover.

### Marking Systems

Aluminum leg bands were used to mark all birds large enough to carry them. For individual field identification, backtags and/or radio-transmitters were used on all birds over 6 weeks old. Neck tags were used on younger birds. Backtags made of vinyl plastic (U. S. Naugahyde) were similar to those described by Labisky and Mann (1962). Neck tags resembling those described by Taber (1949) were made of model airplane covering (Super Monocote). Both types of tags were marked with numbers painted with vinyl paint (Ramacote). Backtags and radio-transmitters were mounted on the center of the back between the wings by a loop of narrow material around the base of each wing securing them in place. Elastic sewing braid, cured buckskin, and No. 18 automotive wire were used at different times to secure the backtags and radio-transmitters. The automotive wire proved most satisfactory as it caused no abrasion and was strong but remained flexible enough to allow freedom of movement. Necktags were placed on the dorsal side of the neck,

midway between the head and the body. A short loop of monofilament line was threaded under a half-inch wide section of skin to attach necktags.

Duck et al. (1970), Carter (1971), and Hanson (1971), using similar telemetry equipment, reported no adverse effect on pheasant mobility or behavior. During this study, radio-equipped hens nested and raised broods successfully. Radio-equipped hens were observed moving with non-radioed hens in harems and breeding with cocks with no apparent adverse effect from the radios.

#### Radio-tracking Systems

The system for locating birds consisted of transmitters, a stationary receiving station and mobile receiving unit (Hanson 1971) and a hand-held receiving unit (Kuck et al. 1970).

Transmitters were built by the Electrical Engineering Department at South Dakota State University. The transmitters emitted a continuous signal on a frequency in the range of 150.0-151.1 MHz. The units had a life of about 100 days and a range of approximately 0.5 mile to the stationary antenna and 0.25 mile to the mobile unit.

A Drake tunable receiver with a converter was used on both the stationary and mobile receiving units. This fine tuning capability of the receiver allowed transmitters on close frequencies to be used with no problem in identifying individual birds.

The stationary tower was fitted with two yagi antennas matched to create a null in the signal when pointing at the transmitter's location. The direction or azimuth was read directly from the Telrex rotation indicator.



The mobile unit had a single yagi antenna. Determination of the azimuth was made by listening for nulls in the audible signal on either side of the loudest point. The degrees between these nulls was halved to find the midpoint direction.

Source of error in the location of birds have been reported as reading and recording the azimuth (Hoezen and Tester 1967), the distance of bird from receiver (Cochrane et al. 1965), and antenna misalignment and bird movement (Kuck et al. 1970). Simultaneous locations from two stations eliminated bird movement as a source of error in this study. Distance from the transmitter to the mobile receiver was kept as small as possible by having 27 alternate sites scattered throughout the study area and always using the site closest to the bird for the recorded location. But distance to the central tower, antenna misalignment and data recording may have contributed to errors in readings. Hanson (1971) reported an error of 40-50 feet at a distance of 0.5 mile with this system. As most readings were made from shorter distances this error was considered acceptable.

#### Data Analysis

Siniff and Tester (1965) described the use of computers in analyzing movement of animals. In this study, distances between locations on a given day were used to reflect activity. Weather conditions were averaged for the period of each movement and entered on IBM cards as were azimuths for each bird location. A computer program was developed to determine the distance between locations and the direction of bird movement.

Locations of birds to be used were selected by the following method. Only days between June 16 and October 15 with at least four locations for a given bird were selected. The four locations used were chosen by dividing the daylight hours (one-half hour before sunrise to one-half hour after sunset) into four equal periods, then selecting the location nearest in time to the midpoint of each of these periods. This selection criterion was used to insure a minimum number of locations per day for each bird to compare in the movement analysis.

Multiple regression was used to analyze movement data. Wind velocity, barometric pressure, precipitation and distance of bird movement were entered in the regression. Distance of bird movement was considered the dependent variable. Linear correlation was used to analyze both the direction of bird movement and wind. A probability level of 0.05 was used in all tests.

Cover-use indices (Robel et al. 1970) were compiled by dividing the percentage of the selected locations for all birds in a given cover type by the percentage of the study area occupied by that cover type. This was done to reduce bias in cover preference that might result from more readings being taken at a particular time of day (Hanson 1971). An index was prepared separately for three divisions (less than 28.12, 28.12-28.29, more than 28.29) under barometric pressure, two divisions (0-8 mph, over 8 mph) under wind velocity, and two divisions (0, any amount) under precipitation. These divisions were made based on the average values for these weather elements during the period of study. An index value of greater than 1.0 indicated greater than expected use.

Barometric pressure used in this study refers to the pressure recorded at the Brookings station not corrected to sea level. All weather values were averages for the period of bird movement to which they refer, except precipitation, which was recorded only as occurring or not. Direction of wind was the actual wind direction measured to the nearest of the eight major compass points to conform to the interpretation of bird direction by the computer.

## RESULTS

### Capture Data

A total of 202 birds was captured and marked in 1971. The majority of birds was captured in areas of undisturbed cover during the months of May and September. Larger numbers of hens were captured in the spring, while almost equal numbers of both sexes were caught in the fall. Greatest success in capturing birds was on calm, dark nights. During the summer months, heavy cover and dispersal of the birds made capture difficult.

### Radio-tracking Movement Data

Data from 21 hens were analyzed: Six adult and two juvenile hens in 1971 and eleven adult and two juvenile hens in 1969 and 1970 (Table 2). Of the adult hens, 13 had broods during the period that they were tracked (Table 3). The average age of these broods and the juvenile hens was 8 weeks. A total of 3,216 locations was obtained on these birds of which 1,068 were selected for analysis. The use of only four locations per day and the elimination of locations of incubating hens account for most of this large reduction in locations.

Data from all 3 years were combined for the movement study. Multiple regression showed no meaningful correlations between distance of bird movement and any of the weather factors tested. The amount of variance explained was only 0.008 or less of 11.3. These results compare with Duever and Fatora's (1968) study on bobwhite quail. They found no correlation between similar weather elements and daily activity patterns

Table 2. Summary of radio-tracking data.

Bird No.	Age	Date On/Off	No. of Days Tracked	No. of Days Used	No. of Locations	No. of Locations Used
215	A	6/26 - 10/20/69	106	36	400	144
216	A	6/30 - 7/23/69	21	8	72	32
218	A	7/10 - 8/1/69	20	7	62	28
273	J	9/25 - 12/4/69	20	1	33	4
308	A	9/23 - 10/17/69	20	5	53	20
312	J	8/24 - 12/6/69	22	2	45	8
332	A	6/21 - 8/27/70	24	4	57	16
334	A	7/16 - 9/1/70	36	11	113	44
336	A	8/10 - 9/4/70	21	9	68	34
350	A	8/25 - 9/21/70	20	12	75	48
425	A	9/4 - 9/11/70	5	4	25	16
426	A	9/4 - 10/6/70	24	9	79	36
427	A	9/16 - 9/29/70	12	1	26	4
509	A	5/7 - 9/1/71	118	21	368	84
551	A	5/11 - 7/29/71	78	27	290	108
562	A	5/22 - 8/13/71	83	10	332	40
567	A	7/30 - 8/16/71	17	7	65	28
569	A	8/10 - 12/31/71	143	52	668	208
574	J	9/10 - 10/3/71	23	13	93	52
581	A	8/11 - 9/5/71	25	16	193	64
621	J	9/8 - 10/1/71	<u>23</u>	<u>14</u>	<u>99</u>	<u>56</u>
Totals			861	269	3216	1068

Table 3. Summary of brood data.

Bird No.	Adult or Juvenile	Dates Tracked	Brood	Age of Brood*
215	A	8/ 5 - 10/15/69	Yes	6 weeks
216	A	7/ 6 - 7/18/69	Yes	5 weeks
218	A	7/11 - 7/28/69	Yes	4 weeks
273	J	9/25/69	No	14 weeks
308	A	9/25 - 10/15/69	No	
312	J	9/25 - 10/15/69	No	10 weeks
332	A	6/25- 8/24/70	No	
334	A	7/22 - 8/24/70	Yes	3 weeks
336	A	8/12 - 8/24/70	Yes	10 weeks
350	A	8/22 - 9/18/70	Yes	6 weeks
425	A	9/ 8 - 9/11/70	Yes	8 weeks
426	A	9/ 8 - 10/ 6/70	No	
427	A	9/ 8 - 10/ 6/70	Yes	14 weeks
509	A	8/ 2 - 9/ 1/71	Yes	0 week
551	A	6/18 - 7/28/71	Yes	0 week
562	A	7/30 - 8/13/71	No	
567	A	8/ 2 - 8/16/71	Yes	0 week
569	A	8/10 - 10/15/71	Yes	5 weeks
524	J	9/13 - 10/ 3/71	No	9 weeks
581	A	8/12 - 9/ 5/71	Yes	5 weeks
621	J	9/ 8 - 9/28/71	No	8 weeks

\*Age at start of tracking period

of quail. The results do not agree with the observations of Shick (1952) or Weston (1954). They reported restriction of pheasant movement by high wind velocity. However, their studies were both conducted in the winter.

Distance birds moved between locations was found to be highly variable. The average distance moved was 0.10 mile with a standard deviation of 0.12 mile. The inclusion of hens with broods of various ages probably accounts for much of this variation. Hanson (1971) found movements of hens increased directly with the age of broods. Also, no correlation was found between direction of bird movement and wind direction.

#### Location of Birds

Because acreages of cover types varied during this study, percentage-of-bird-location data and habitat-use indices were compiled for each year separately (Appendix Tables 2, 3, 4).

In 1969, three cover types showed the widest variations between the categories of barometric pressure (Table 4). The percentage of bird locations in corn and small grain decreased by one-half as the pressure went from the lowest pressure category to the highest. Bird locations in alfalfa showed exactly an opposite trend, almost doubling as the pressure increased. Use in treecrow and farmstead also doubled with an increase in barometric pressure from the less than 28.12 inch category to the over 28.29 category. The percentage of bird locations in corn and residual cover were the only values that changed greatly between the divisions of wind velocity. Bird locations in corn increased at wind

Table 4. Percentage of locations in cover types, 1969.

	Number of Locations	Corn	Residual Cover	Small Grain	Pasture	Alfalfa	Treerow & Farmstead	Ditches	Spoil Pits	Summer Fallow
<u>Barometric pressure</u>										
Less than 28.11	55	14.5	21.8	34.5	3.6	14.5	5.5	0.0	1.8	3.6
28.11-28.29	85	21.2	25.9	21.2	1.3	21.2	8.2	0.0	1.3	0.0
28.30 or higher	92	8.7	28.3	16.3	1.1	30.4	10.9	1.1	1.1	2.0
<u>Wind velocity</u>										
0-8 mph	128	8.6	33.6	19.5	0.0	26.6	7.8	0.8	0.0	3.1
Over 8 mph	104	22.1	16.3	26.0	3.8	19.2	9.6	0.0	2.9	0.0
<u>Precipitation</u>										
None	206	16.0	21.4	24.8	1.9	25.7	7.3	0.5	1.5	1.0
Any amount	26	3.8	61.5	3.8	0.0	3.8	19.2	0.0	0.0	7.7



velocities over 8 mph, while locations in residual cover fell sharply. Large differences occurred in almost all cover types between the precipitation categories, but the small number of locations (26) in the zero precipitation division is probably responsible for these differences.

In 1970, no large changes in percentage of locations were noted in any cover type (Table 5). Locations in corn accounted for over 50 percent of the locations in all weather categories. The percentage of locations in residual cover and alfalfa were much lower than the 1969 figures.

The percentage of bird locations in each cover type remained close to the same levels in 1971 as they were in 1970 (Table 6). Fifty percent or more of all locations were in corn. The only changes in use occurred between the wind-velocity categories. Bird locations in corn increased by one-third between the low and high wind-velocity categories while locations in residual cover and alfalfa decreased by one-half.

For the three-year period, the highest percentage of locations were in corn, residual cover, alfalfa, and small grain.

Table 5. Percentage of Locations in cover types, 1970.

	Number of Locations	Corn	Residual Cover	Small Grain	Pasture	Alfalfa	Treerow & Farmstead	Ditches	Spoil Pits	Summer Fallow
<u>Barometric pressure</u>										
Less than 28.11	74	60.8	10.8	12.2	8.1	4.0	0.0	4.0	0.0	0.0
28.11-28.29	66	65.2	0	16.7	10.6	6.0	0.0	1.5	0.0	0.0
28.30 or higher	56	66.1	14.3	14.3	1.8	3.6	0.0	0	0.0	0.0
<u>Wind velocity</u>										
0-8 mph	83	56.6	9.6	13.9	7.2	7.2	0.0	1.2	0.0	0.0
Over 8 mph	113	69.0	7.1	11.5	7.1	2.7	0.0	2.7	0.0	0.0
<u>Precipitation</u>										
None	179	64.2	7.3	14.0	7.8	5.0	0.0	1.7	0.0	0.0
Any amount	17	58.8	17.6	17.6	0.0	0.0	0.0	5.9	0.0	0.0

Table 6. Percentage of locations in cover types, 1971.

	Number of Locations	Corn	Residual Cover	Small Grain	Pasture	Alfalfa	Treerow & Farmstead	Ditches	Spoil Pits	Summer Fallow
<u>Barometric pressure</u>										
Less than 28.11	156	63.5	9.0	3.8	5.8	16.0	1.3	0.6	0.0	0.0
28.11-28.29	232	56.5	6.0	5.2	6.5	21.1	2.2	2.6	0.0	0.0
28.30 or higher	252	63.5	10.3	3.6	3.2	16.7	1.2	1.6	0.0	0.0
<u>Wind velocity</u>										
0-8 mph	442	54.3	10.0	4.8	5.0	21.7	2.0	2.0	0.0	0.0
Over 8 mph	198	75.8	5.1	3.0	5.1	10.1	0.5	.5	0.0	0.0
<u>Precipitation</u>										
None	555	59.8	8.6	4.5	5.2	18.4	1.6	1.8	0.0	0.0
Any amount	85	68.2	7.1	2.3	3.4	16.5	1.2	1.2	0.0	0.0

### Habitat Use Indices

While the percentage of locations in various cover types provides a general indication of habitat use under varying weather conditions and overall use, habitat-use indices were compiled to determine habitat preferences. An index for each weather class was prepared for each of the 3 years of study. These indices were then combined and a weighted mean index value calculated for each weather class.

Alfalfa was the preferred cover type under all conditions (mean index 2.2-4.8). Corn was the only other cover type for which a preference was shown under all conditions (mean index 1.2-1.8).

Under low barometric pressure conditions, alfalfa, corn and residual cover were the preferred habitat types with indices over 1.0 (Table 7). Pasture and ditches had indices of 0.7 while all other cover types were 0.5 or less. In the mid-range class of pressures alfalfa, corn, residual cover, and treerow and farmstead were the preferred cover types. Preference for alfalfa and treerow and farmstead increased markedly in this class. Use of other cover types remained at the same levels. Under high pressure conditions the same four types remained preferred, but residual cover was preferred over corn, increasing from a mean index of 1.0 to 1.8. No changes in preference were noted in any other cover type.

Analysis of wind velocity showed three major shifts in preference (Table 8). At low wind velocity the mean index value for alfalfa was at the highest point (4.8). Other preferred cover types at low wind

Table 7. Cover-use index based on changes in barometric pressure.

Year	Number of Locations	Corn	Residual Cover	Small Grain	Pasture	Alfalfa	Treecrow & Farmstead	Ditches	Spoil Pits	Summer Fallow
<u>Station pressure 28.11 or less</u>										
1969	55	0.5	1.1	1.4	0.4	2.4	1.5	0.0	1.3	0.6
1970	74	1.9	0.5	0.5	0.9	1.0	0.0	2.0	0.0	0.0
1971	156	1.7	1.8	0.1	0.7	4.0	0.4	0.3	0.0	0.0
Weighted mean index		1.5	1.3	0.5	0.7	2.9	0.5	0.7	0.4	0.1
<u>Station pressure 28.11-28.29</u>										
1969	85	0.7	1.4	0.8	0.1	3.5	2.7	0.0	1.3	0.0
1970	66	2.0	0.0	0.7	1.2	1.5	0.0	0.8	0.0	0.0
1971	232	1.5	1.2	0.2	0.8	5.3	0.7	1.3	0.0	0.0
Weighted mean index		1.4	1.0	0.4	0.7	4.3	1.0	0.9	0.3	0.0
<u>Station pressure 28.30 or higher</u>										
1969	92	0.3	1.5	0.7	0.1	5.1	3.6	0.6	1.1	0.3
1970	56	2.1	0.7	0.6	0.2	0.9	0.0	0.0	0.0	0.0
1971	252	1.7	2.1	0.1	0.4	4.2	0.4	0.8	0.0	0.0
Weighted mean index		1.4	1.8	0.3	0.4	4.0	0.1	0.6	0.3	0.1

Table 3. Cover-use index based on changes in wind velocity.

Year	Number of Locations	Corn	Residual Cover	Small Grain	Pasture	Alfalfa	Treerow & Farmstead	Ditches	Spoil Pits	Summer Fallow
<u>Wind velocity 0-8 mph</u>										
1969	128	0.3	1.8	0.8	0.0	4.4	2.6	0.4	0.0	0.5
1970	83	1.8	0.5	0.6	0.8	1.8	0.0	0.6	0.0	0.0
1971	442	<u>1.2</u>	<u>1.8</u>	<u>0.4</u>	<u>0.5</u>	<u>4.8</u>	<u>1.0</u>	<u>0.8</u>	<u>0.0</u>	<u>0.1</u>
Weighted mean index		1.2	1.8	0.4	0.5	4.8	1.0	0.8	0.0	0.1
<u>Wind velocity over 8 mph</u>										
1969	104	0.8	0.9	1.0	0.4	3.2	3.2	0.0	2.9	0.0
1970	113	2.2	0.4	0.5	0.8	0.7	0.0	1.4	0.0	0.0
1971	198	<u>2.0</u>	<u>1.0</u>	<u>0.1</u>	<u>0.6</u>	<u>2.5</u>	<u>0.2</u>	<u>0.3</u>	<u>0.0</u>	<u>0.0</u>
Weighted mean index		1.8	0.8	0.4	0.6	2.2	0.9	0.5	0.7	0.0

velocities were residual cover, corn and treerow and farmstead. When wind velocity increased to over 8 mph the only cover types still preferred by the birds were alfalfa and corn. The highest preference for corn (1.8) was reached under this condition while the lowest levels of 0.8 and 2.2 were reached for residual cover and alfalfa respectively.

During precipitation, preference for residual cover and treerow and farmstead increased markedly with the mean index for treerow and farmstead reaching 1.6, the highest level recorded (Table 9). Preference for alfalfa dropped rather sharply, going from 3.9 to 2.8 during periods of precipitation. Most other cover type preferences remained at the same level or dropped slightly during precipitation.

Table 9. Cover-use index based on occurrence of precipitation.

Year	Number of Locations	Corn	Residual Cover	Small Grain Pasture	Alfalfa	Treerow & Farmstead	Spoil Pits	Summer Fallow		
<u>Precipitation none</u>										
1969	206	0.6	1.1	1.0	0.2	4.3	2.4	0.3	1.5	0.2
1970	179	2.0	0.4	0.6	0.9	1.3	0.0	0.8	0.0	0.0
1971	555	<u>1.6</u>	<u>1.7</u>	<u>0.1</u>	<u>0.7</u>	<u>4.6</u>	<u>0.5</u>	<u>0.9</u>	<u>0.0</u>	<u>0.0</u>
Weighted mean index		1.5	1.3	0.4	0.6	3.9	0.8	0.8	0.3	0.0
<u>Precipitation any amount</u>										
1969	26	0.1	3.2	0.2	0.0	0.6	6.4	0.0	0.0	1.3
1970	17	1.8	0.9	0.7	0.0	0.0	0.0	2.9	0.0	0.0
1971	85	<u>1.8</u>	<u>1.4</u>	<u>0.1</u>	<u>0.4</u>	<u>4.1</u>	<u>0.4</u>	<u>0.6</u>	<u>0.0</u>	<u>0.0</u>
Weighted mean index		1.5	1.7	0.2	0.3	2.8	1.6	0.8	0.0	0.3



## DISCUSSION AND CONCLUSION

No significant relationships were found between distance of bird movement and any of the weather factors tested ( $P < 0.05$ ). Other studies such as that by Klonglan (1955), have shown weather to have an effect on daily activity of pheasants. In this study the weather parameters tested were assumed to reflect other weather elements. Further testing of additional weather factors may prove this assumption false. Normal daily fluctuations of weather elements did not affect pheasant activities to a large degree.

Mean index values for alfalfa and corn were high (greater than 1.0) for all weather conditions; mean index values for residual cover and treerows and farmstead were high under most conditions. While the preference for alfalfa was high at all times, it did vary. Under conditions of precipitation, high wind velocity, and low barometric pressure the preference index for alfalfa was much lower. With these same conditions an increase in use of corn and residual cover was noted. This seems to be a logical shift in usage because these two cover types offered more protection from the elements.

Index values for residual cover remained high the last year of the study despite the large reduction in amount of residual cover available, indicating a strong preference for this type of cover.

This was a quantitative study of habitat preference. Variations in quality of different cover types suggest that a qualitative study of habitat use could provide answers to questions about habitat preference.

## LITERATURE CITED

- Bernitt, R., and H. V. Terrill. 1940. Possible temperature factors in north central pheasant management. *Trans. N. Am. Wildl. Conf.* 5:428-432.
- Buss, I. O., and C. V. Swanson. 1950. Some effects of weather on pheasant reproduction in southeastern Washington. *Trans. N. Am. Wildl. Conf.* 15:365-378.
- Carter, A. V. 1971. Seasonal movements and behavior of ring-necked pheasants in eastern South Dakota. M. S. thesis, S. Dak. State Univ. 44 p.
- Cochrane, W. W., and R. D. Lord, Jr. 1963. A radio-tracking system for wild animals. *J. Wildl. Mgmt.* 27(1):9-24.
- \_\_\_\_\_, D. W. Warner, J. R. Tester, and V. B. Kuechle. 1965. Automatic radio-tracking system for monitoring animal movements. *BioScience* 15(2):98-106.
- Drewien, R. C., H. M. Reeves, P. F. Springer, and T. L. Kuck. 1967. Backpack unit for capturing waterfowl and upland game by night-lighting. *J. Wildl. Mgmt.* 31(4):778-783.
- Duever, M. J., and J. R. Fatora. 1968. Daily and seasonal activity patterns of bobwhite quail on the AEC Savannah River Plant. *Proceedings of the 22nd Annual Southeastern Association of Game and Fish Commissioners, Baltimore, Md., pp. 189-195.*
- Graham, S. A., and G. Hesterburg. 1948. The influence of climate on the ring-necked pheasant. *J. Wildl. Mgmt.* 12(1):9-15.
- Grondahl, C. R. 1953. Winter behavior of the ring-necked pheasant as related to winter cover in Winnibago County, Iowa. *Iowa State Coll. J. Sci.* 27(4):447-465.
- Hanson, L. E. 1971. Movements and habitat use by hen pheasants during brood rearing. M. S. thesis, S. Dak. State Univ. 54 pp.
- Heezen, H. L., and J. R. Tester. 1967. Evaluation of radio-tracking by triangulation with special reference to deer movements. *J. Wildl. Mgmt.* 31(1):124-141.
- Klonglar, E. D. 1955. Factors influencing the fall roadside pheasant census in Iowa. *J. Wildl. Mgmt.* 19(2):254-262.
- Kozlicky, E. L., C. O. Hendrickson, P. G. Hoveyer, and R. Nansen. 1955. Weather and fall pheasant populations in Iowa. *J. Wildl. Mgmt.* 19(1):163-172.

- Kuck, T. L., R. B. Dahlgren, and D. R. Progulske. 1970. Movements and behavior of hen pheasants during the nesting season. *J. Wildl. Mgmt.* 34(3):626-630.
- Labisky, R. F. 1968. Nightlighting: its use in capturing pheasants, prairie chickens, bobwhites, and cottontails. *Illinois Nat. Hist. Biol. Survey Notes No. 62.* 12 pp.
- \_\_\_\_\_, J. A. Harper, and F. Greely. 1964. Influence of land use, calcium, and weather on the distribution and abundance of pheasants in Illinois. *Ill. Nat. Hist. Sur. Biol. Notes No. 51.* 19 pp.
- \_\_\_\_\_, and S. H. Mann. 1962. Backtag markers for pheasants. *J. Wildl. Mgmt.* 26(4):393-399.
- Latham, R. M. 1947. Differential ability of male and female game birds to withstand starvation and climatic extremes. *J. Wild. Mgmt.* 11:139-149.
- Lemnyan, C. D., W. White, E. Nyberg, and J. J. Christian. 1959. Design of a miniature radio transmitter for use in animal studies. *J. Wildl. Mgmt.* 25(1):107-110.
- MacMullan, R. A., and L. L. Eberhardt. 1953. Tolerance of incubating pheasant eggs to exposure. *J. Wild. Mgmt.* 17:322-330.
- Marshall, W. H. 1960. Development and use of short wave radio transmitters to trace animal movements. *Prog. Rept. to Nat. Sci. Foundation*, 27 pp. multilith. Univ. of Minn.
- \_\_\_\_\_, and J. J. Kupa. 1963. Development of radio-telemetry techniques for ruffed grouse studies. *Trans. N. Am. Wildl. and Nat. Resources Conf.* 28:443-456.
- Robel, R. J., J. N. Briggs, J. J. Cebula, W. J. Silvy, C. E. Viers, and P. G. Watt. 1970. Greater prairie chicken ranges, movements, and habitat use in Kansas. *J. Wildl. Mgmt.* 34(2):286-306.
- Shick, C. 1952. A study of pheasants on the 9,000 acre prairie farm, Saginaw County, Michigan. *Game Division, Michigan Dept. of Cons. Bulletin.* 134 pp.
- Siniff, D. B., and J. R. Tester. 1965. Computer analysis of animal movement obtained by telemetry. *BioScience.* 15(2):104-108.
- Tabor, R. D. 1949. A new marker for game birds. *J. Wildl. Mgmt.* 13(2):228-231.

- Trautman, C. G., and R. B. Dahlgren. 1965. Causes and mechanisms of pheasant population fluctuations. Game Div. South Dakota Dept. of Game, Fish and Parks. 30 pp. (Mimeo).
- U. S. Dept. of Commerce. 1969. Climatological data. Vol. No. 74.
- U. S. Dept. of Commerce. 1970. Climatological data. Vol. No. 75.
- U. S. Dept. of Commerce. 1971. Climatological data. Vol. No. 76.
- U. S. Dept. of Commerce, 1970. Climatic atlas of the United States, Environmental Sciences Services Adm., Environmental Data Service. U. S. Govt. Print. Off., Washington, D. C.
- Westin, F. C., G. J. Buntley, F. E. Shubeck, L. F. Puhr, and N. E. Bergstresser. 1958. Soil survey, Brookings County, South Dakota. U. S. Govt. Print. Off., Washington D. C., 87 pp.
- Weston, H. G., Jr. 1954. Winter-spring movements of the ring-necked pheasant in northern Iowa. Iowa State Coll. J. Sci. 29:39-60.

APPENDIX

Appendix Table 1. Summary of weather information.

	<u>Barometric Pressure</u>			<u>Wind Velocity</u>		<u>Precipitation</u>	<u>Temperature</u>		
	High	Low	Ave	High	Ave	Amount	+ Normal	Amount	+ Normal
<u>1969</u>									
June	28.54	27.39	28.10	24	10.0	7.20	+3.25	58.5	-8.6
July	28.54	27.96	28.25	17	8.3	3.58	+1.33	69.6	-3.6
Aug.	28.45	27.88	28.21	24	10.1	1.53	-1.44	69.7	-1.5
Sept.	28.56	27.87	28.30	22	9.0	1.32	-0.65	59.8	-1.5
Oct.	28.63	27.68	28.26	23	10.0	1.87	+0.65	42.0	-7.5
<u>1970</u>									
June	28.51	27.79	28.17	23	10.9	4.22	+0.27	66.8	-0.3
July	28.56	28.00	28.24	20	8.5	2.35	+0.20	71.0	-2.2
Aug.	28.44	27.96	28.23	18	7.6	1.24	-1.73	68.5	-2.7
Sept.	28.56	27.69	28.22	24	9.4	1.13	-0.84	58.2	-3.1
Oct.	28.72	27.75	28.18	24	10.2	2.72	+1.50	47.7	-1.8
<u>1971</u>									
June	28.41	27.36	28.18	17	6.7	5.16	+1.21	68.6	+1.5
July	28.51	27.98	28.26	15	6.8	1.13	-1.02	65.9	-7.3
Aug.	28.57	28.04	28.28	18	7.0	3.00	+0.03	68.6	-2.6
Sept.	28.63	27.82	28.23	24	8.0	0.88	-1.09	57.7	-3.6
Oct.	28.62	27.52	28.22	24	7.7	3.56	+2.34	48.7	-0.8

Appendix Table 2. Number of locations in cover types, 1969.

	Corn	Residual Cover	Small Grain	Pasture	Alfalfa	Tree row & Farmstead	Ditches	Spoil Pits	Summer Fallow	Total Locations
<u>Station pressure</u>										
Less than 28.11	8	12	19	2	8	3	0	1	2	55
28.11-28.29	18	22	18	1	18	7	0	1	0	85
28.30 or higher	8	26	15	1	28	10	1	1	2	92
<u>Wind velocity</u>										
0-8 mph	11	43	25	0	34	10	1	0	4	128
Over 8 mph	23	17	27	4	20	10	0	3	0	104
<u>Precipitation</u>										
None	33	44	51	4	53	15	1	3	2	206
Any amount	1	16	1	0	1	5	0	0	2	26

Appendix Table 3. Number of locations in cover types, 1970.

	Corn	Residual Cover	Small Grain	Pasture	Alfalfa	Tree row & Farmstead	Ditches	Spoil Pits	Summer Fallow	Total Locations
<u>Station pressure</u>										
Less than 28.11	45	8	9	6	3	0	3	0	0	74
28.11-28.29	43	0	11	7	4	0	1	0	0	66
28.30 or higher	37	8	8	1	2	0	0	0	0	56
<u>Wind velocity</u>										
0-8 mph	47	8	15	6	6	0	1	0	0	83
Over 8 mph	78	8	13	8	3	0	3	0	0	113
<u>Precipitation</u>										
None	115	13	25	14	9	0	3	0	0	179
Any amount	10	3	3	0	0	0	1	0	0	17



Appendix Table 4. Number of locations in cover types, 1971.

	Corn	Residual Cover	Small Grain	Pasture	Alfalfa	Tree row & Farmstead	Ditches	Spoil Pits	Summer Fallow	Total Locations
<u>Station pressure</u>										
Less than 28.11	99	14	6	9	25	2	1	0	0	156
28.11-28.29	131	14	12	15	49	5	6	0	0	232
28.30 or higher	160	26	9	8	42	3	4	0	0	252
<u>Wind velocity</u>										
0-8 mph	240	44	21	22	96	9	10	0	0	442
Over 8 mph	150	10	6	10	20	1	1	0	0	198
<u>Precipitation</u>										
None	332	48	25	29	102	9	10	0	0	555
Any amount	58	6	2	3	14	1	1	0	0	85