

South Dakota State University

Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

Agricultural Experiment Station Circulars

SDSU Agricultural Experiment Station

12-1992

1992 South Dakota Corn Performance Trials

R. G. Hall

South Dakota State University, robert.hall@sdstate.edu

Follow this and additional works at: http://openprairie.sdstate.edu/agexperimentsta_circ

Recommended Citation

Hall, R. G., "1992 South Dakota Corn Performance Trials" (1992). *Agricultural Experiment Station Circulars*. Paper 291.
http://openprairie.sdstate.edu/agexperimentsta_circ/291


This Circular is brought to you for free and open access by the SDSU Agricultural Experiment Station at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in Agricultural Experiment Station Circulars by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

C 253
(revised annually)

1 9 9 2
South Dakota

CORN
PERFORMANCE
TRIALS

Agricultural Experiment Station
South Dakota State University
U.S. Department of Agriculture



1 9 9 2

South Dakota

CORN

PERFORMANCE TRIALS

Robert G. Hall

Assoc. Professor/Extension Agronomist
Manager, Crop Performance Testing

Plant Science Department
Agricultural Experiment Station
South Dakota State University
Brookings, SD 57007-1096

The relative performance of corn hybrids grown under similar environmental conditions in 1992 is evaluated in this report.

Information in the tables includes both 1992 and 1991-92 grain yields in bushels per acre (bu/a); and 1992 test weight, moisture percentages of shelled corn at harvest, final plant populations per acre, and stalk lodging percentages.

The test trials reported in this publication were conducted by the Plant Science Department Crop Performance Testing Program, Agricultural Experiment Station, South Dakota State University.

Location, Weather, and Climatic Conditions

Test trials locations and their nearest post office designations are in Table 1.

Climatic data (Table 2) for the 1992 corn growing season, April-September, are based upon U.S. Monthly Climatological Data (NOAA) recorded at a weather sta-

The assistance of the following individuals is appreciated: Dwayne Beck, Loyal Evjen, Delbert Robbins, and Dale Sorensen of the Stations; technician Kevin Kirby and Darin Huber; Robert Clark, Eldon Schmidt, and Steve Masat, farmer-cooperators.

tion reasonably near each trial site. The Milbank recording station is closest to the field north of Gary in Deuel County. The Watertown and Pierre airport data are used for the NE Farm and Dakota Lakes Research Farm trials, respectively. Stations are located at or near the other trial sites. Precipitation quantities would vary from the actual site to the recording station, but temperatures are similar over a much wider area and considered applicable to the trial area.

Field conditions varied in eastern South Dakota during most of the growing period. Spring was early and fairly dry, so seeding was completed by May 8. On Memorial Day weekend a severe freeze hit most of eastern South Dakota. The early trial at Pierre was reseeded on June 10.

April temperatures (Table 2) were 2 to 5° F below normal but were slightly above normal by the end of May. Temperatures dropped below normal about 3° in June, 9° in July, and 6° in August. Precipitation was fairly uniform for June, July, and August at most locations except Milbank. Precipitation began falling in mid-June and was fairly uniform except at Milbank.

Cool temperatures in June, July, and

August slowed growth which later gave pollination problems. The delayed growth caused silking and tasseling stages to be out of sync; some silks emerged after pollen shed was complete, resulting in some unfertilized silks and reduced yields.

In some cases, cool weather slowed silk growth. Short silks did not emerge before ear sheaths enclosed the ear; consequently, silks were trapped or formed silk balls under the ear sheath, resulting in unfertilized silks and reduced yields.

Hybrid Entry Procedure

Participating companies designated the locations where their entries were to be grown. Beginning in 1986, the entries were placed into early or late trials based upon maturity information supplied by the company. The arbitrary breaks at each site were 95 days for Watertown, Gary, and Brookings; 100 days for Frankfort and Pierre; and 110 days for Beresford and Armour. A fee was charged for every entry at each location. A listing of the participating firms and their hybrids is on the last pages of this circular.

Experimental Procedure

Entries in each trial were seeded in three replications. Plots of individual hybrids were located at random within each replication. Each plot consisted of two 30-inch rows, 26 feet long. The target seeding population per acre for each location was, in seeds per acre:

Watertown	21,440
Gary	21,440
Frankfort	19,000
Brookings	21,440
Beresford	23,450
Armour	19,000
Pierre, irrigated	33,500

Soil types, starter fertilizer applications, and land preparation at each test trial site are outlined in Table 3. No insecticides were used for corn root-worm control this year. A recommended short-residue preemergence herbicide was banded over the row at seeding at most sites.

A 31-cell cone drill seeder was used for all plots. Cone units were mounted

above commercial maxi-merge units. Seeding rate was 15% more than the desired number of plants harvested per plot. Seedbed preparation was good at all locations.

Final stands in most of the trials were near desired levels, except for six hybrids in the early trial at Watertown, five hybrids in the late trial at Frankfort, and seven hybrids in the early trial at Armour. These hybrids can be identified by comparing their final populations per acre to the minimum best value population at the bottom of the performance tables at these locations.

Measurements of Performance

Yield. Yields in all corn test trials are averaged from three replications and expressed as bushels per acre at 15.5% moisture and a bushel weight of 56 lb. Varieties of equal potential may yield differently because of variations in slope, soil fertility, and stand. Statistical tests were conducted to determine whether differences were caused by variations in environment or were true varietal differences.

The coefficient of variation (CV) for yield was greater than desired for the replanted early test trial at Pierre. Ideally, the CV value, which is a measure of experimental error, should not exceed 16%. **If the CV value exceeds 16%, the data should not be used to make hybrid comparisons because of the high degree of experimental error measured in the test trial.**

Experimental error may be the result of several factors, including test methods, or environmental factors such as moisture, temperature, soil variations, or agronomic factors like seeding date, reseeding, or seed quality factors, all of which may or may not be controllable in a given year.

To convert data in these tables to the metric system use the following methods. (The factor 1.121 converts from lb/a to kg/ha.)

I. 1 lb = 0.454 kilograms; 1 hectare = 2.471 a, so
 $56 \times 0.454 \times 2.471 = 62.8 \text{ x bu/a} = \text{kilograms per hectare.}$

II. Or, assuming a yield of 62.8 bu/a from the tables;

Step 1 = 62.8 bu/a \times 56 lb/bu = 3516.8 lb/a
Step 2 = 3516.8 lb/a \times 1.121 = 3942.3 kilograms/hectare or 39.4 quintals/hectare.

Moisture Content. The moisture content of each entry is expressed as the percentage of moisture in the shelled corn at the time of harvest. Moisture content is inversely related to maturity.

Because maturity is of prime importance in South Dakota, moisture figures are of considerable importance in the evaluation of the trial entries. Hybrids that provide satisfactory yields and can be stored without additional drying are desirable.

Stalk lodging. The stalk lodging percentage for each entry is expressed as the number of stalks lodged below the ear divided by the number of plants in each plot at harvest. The lodging percentages in both trials at Pierre were high for 1992. This was caused by extremely high winds on December 3 which broke many stalks below the ear.

Use of tables. First check for the "least significant difference" (LSD) value at the bottom of each column of data averages. The LSD value indicates how much a variable like yield must differ between two hybrids for there to be a real yield difference. An LSD value is given at the bottom of every column where there is significant difference among the averages within that column. If there is no real differences among the averages within a given column a "nonsignificant" (NS) difference designation is indicated.

The LSD value for a variable can be used in two ways. First, you can compare whether two hybrids really differ in yield. In this case the yield difference between two hybrids in the early trial at Watertown must be more than 11.3 bu/a; if not, there is no real yield difference between the two hybrids.

Second, you can use the same LSD value to identify which hybrids are the best yielders at a given location.

For example, in the early maturity test trial at Watertown the highest yield was 73.7 bu/a for Cargill 3427. To find whether Cargill 3427 is the only top

yielder at Watertown, use the LSD value of 11.3 bu/a at the bottom of the 1992 yield column. If hybrids are in the best yield group they will yield within 11.3 bu/a of Cargill 3427 (73.7 - 11.3), or yield 62.4 bu/a or more.

Therefore, any hybrid yielding 62.4 bu/a or higher is in the top yielding group, or we can say 62.4 bu/a is the minimum best value for top yielding hybrids at Watertown in 1992. In this case the hybrids Cargill 3427 down through and including Pioneer 3921 are in the best yield group because they all yielded higher than 62.4 bu/a.

Likewise, the best minimum value for bushel weight is 43.5 lb and for final population per acre is 20,725. Only those hybrids with a final acre population of 20,725 plants should be compared with one another because there is no significant difference in population. If a hybrid's final acre population is less than 20,725, its yields should not be compared with other hybrids unless the difference between their final acre populations is equal to or less than 1,050 plants (the final acre population LSD value).

In the case of harvest moisture content and stalk lodging percentages, look for the maximum best value at the bottom of those table columns. For moisture content and stalk lodging percentages the appropriate LSD values are added to the lowest numerical value within a column to obtain the maximum best values. At Watertown the best harvest moisture percentages within the harvest moisture column are 20.4% or less. Since there were no significant difference among the hybrids for stalk lodging percentage no maximum best value was calculated.

In summary, in the early maturity test trial at Watertown, the best hybrids will have a yield higher than 62.4 bu/a, a bushel test weight of 43.5 lb or higher, a harvest moisture of 20.4% or less, and a final acre population of 20,725 plants or higher.

Published in accordance with an act passed in 1881 by the 14th Legislative Assembly, Dakota Territory, establishing the Dakota Agricultural College and with the act of re-organization passed in 1887 by the 17th Legislative Assembly, which established the Agricultural Experiment Station at South Dakota State University.

3,000 printed by AES at a cost of 53¢ each.
AX062 1/93.

Table 1. Test trial locations, seeding dates, and harvest dates.

LOCATION	COUNTY	POST OFFICE	DATE SEEDED	DATE HARVESTED
N.E. RES. FARM, 15N	CODINGTON	WATERTOWN	MAY 5	DEC. 8
ELDON SCHMIDT FARM	DEUEL	GARY	MAY 8	OCT. 28
STEVE MASAT FARM, 6S, 2W	SPINK	FRANKFORT	APRIL 30	NOV. 11
SDSU AGRONOMY FARM, 2NE	BROOKINGS	BROOKINGS	APRIL 29	NOV. 18
S.E. RES. FARM, 7W, 3S	UNION	BERESFORD	MAY 1	NOV. 16
ROBERT CLARK FARM, 4W, 1S	DOUGLAS	ARMOUR	MAY 1	NOV. 11
DAK. LAKES RES. FARM, 17E, LATE	HUGHES	PIERRE	APRIL 30	DEC. 7
DAK. LAKES RES. FARM, 17E, EARLY	HUGHES	PIERRE	JUNE 10*	DEC. 7

*REPLANTED FOLLOWING LATE MAY FREEZE.

Table 2. Temperature and precipitation data for the 1992 South Dakota corn performance trials.

LOCATION	TYPE OF DATA	APRIL	MAY	JUNE	JULY	AUG.	SEPT.	TOTAL
WATERTOWN (N.E. FARM)	PRECIPITATION*	1.28	0.41	6.67	3.21	2.16	2.89	16.62
AIRPORT	AVER. TEMP.**	40.4	58.0	62.7	62.3	63.0	58.4	
	TEMP. DIFF.	-4.4	+0.9	-4.1	-10.2	-7.3	-1.2	
	FIRST FREEZE			SEPT. 22, 30				DEGREES, F.
MILBANK (GARY) 2 SSW	PRECIPITATION	2.53	1.13	6.74	5.74	0.96	1.48	18.58
	AVER. TEMP.	40.4	59.9	64.3	64.2	64.7	58.8	
	TEMP. DIFF.	-3.2	+3.0	-2.5	-8.0	-5.1	-0.4	
	FIRST FREEZE			SEPT. 28, 28				DEGREES, F.
REDFIELD (SPINK CO.) 2 NE	PRECIPITATION	0.58	0.88	7.28	2.93	3.30	1.18	16.15
	AVER. TEMP.	42.4	58.9	63.6	63.9	64.5	58.1	
	TEMP. DIFF.	-2.6	+1.9	-3.3	-9.4	-6.2	-1.8	
	FIRST FREEZE			SEPT. 28, 23				DEGREES, F.
BROOKINGS 2 NE	PRECIPITATION	1.79	1.19	7.98	3.69	4.67	1.87	21.19
	AVER. TEMP.	38.7	58.3	62.8	62.0	62.8	56.9	
	TEMP. DIFF.	-5.3	+2.4	-2.8	-8.7	-5.2	-1.3	
	FIRST FREEZE			SEPT. 28, 32				DEGREES, F.
CENTERVILLE (S.E. FARM) 6 SE	PRECIPITATION	1.92	2.32	1.89	6.74	4.69	4.17	21.73
	AVER. TEMP.	42.2	60.0	66.4	65.7	65.2	59.4	
	TEMP. DIFF.	-5.3	+1.3	-2.7	-8.2	-6.2	-1.9	
	FIRST FREEZE			SEPT. 28, 30				DEGREES, F.
ARMOUR (DOUGLAS CO.)	PRECIPITATION	2.76	0.53	1.97	6.12	4.67	2.78	19.83
	AVER. TEMP.	47.0	62.1	67.1	67.4	67.9	62.7	
	TEMP. DIFF.	-2.0	+1.8	-2.9	-8.8	-5.6	-0.5	
	FIRST FREEZE			SEPT. 28, 29				DEGREES, F.
PIERRE (DLR FARM) AIRPORT	PRECIPITATION	0.47	0.54	4.74	4.52	1.70	0.93	12.90
	AVER. TEMP.	45.7	60.4	65.2	66.5	67.3	62.6	
	TEMP. DIFF.	-1.4	+2.1	-3.5	-9.5	-6.6	-0.2	
	FIRST FREEZE			SEPT. 28, 26				DEGREES, F.

*PRECIPITATION = INCHES, TEMPERATURE = DEGREES FAHRENHIET.

Table 3. Soil classification, fertilizer applied and land preparation.

LOCATION	SOIL TYPE	STARTER, 2 X 2 FERTILIZER APPLIED (ACTUAL PER ACRE)			LAND PREPARATION		
		37	-	18	-	00	
BROOKINGS	BRANDT SIL*	37	-	18	-	00	CONVENTIONAL
WATERTOWN	KRANZBURG SIL	37	-	18	-	00	CONVENTIONAL
FRANKFORT	BEOTIA SICL	37	-	18	-	00	NO-TILL, WHEAT STUBBLE
CENTERVILLE	EGAN SIL	37	-	18	-	00	CONVENTIONAL
ARMOUR	EAKIN SIL	37	-	18	-	00	NO-TILL, WHEAT STUBBLE
PIERRE	LOWRY SIL	37	-	18	-	00	NO-TILL, SOYBEAN STUBBLE
GARY	FORMAN SICL	37	-	18	-	00	CONVENTIONAL

SIL= SILT LOAM, SICL=SILTY CLAY LOAM.

Table 4. 1992 corn hybrid performance trial results--Watertown, SD, N.E. Research Farm, early maturity--95 days or less.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
CARGILL 3427	73.7	108.4	24.7	35.8	21440	0	1X
DEKALB DK462	71.7	110.2	24.5	36.7	20882	0	1X
DEKALB DK401	71.2	101.4	19.2	42.9	21328	0	1X
ISC-PAYCO 531	70.6	•	25.1	36.9	20770	0	2X
PIONEER 3861	68.9	•	21.1	41.6	20770	0	1X
DAHLGREN D5962	66.8	•	25.0	35.1	21217	1	2X
SEXAUER SX450	66.4	•	24.5	38.1	20770	0	2X
ICI 8777	66.2	98.7	25.9	35.0	20993	0	2X
AGRIGENE AG3500	65.8	•	25.3	34.9	20993	0	2X
TOP FARM SX1194A	65.0	•	21.1	42.2	21440	0	2X
PIONEER 3921	62.6	92.0	19.5	46.9	20658	0	1X
DAIRYLAND ST1189	61.8	•	22.6	42.0	21217	0	2X
DEKALB DK381	61.0	•	18.8	41.3	21217	0	1X
NC+ 1991	60.1	•	25.2	35.2	21105	0	2X
JOHNSON SX417	59.5	•	25.1	37.7	20212	0	2X
ASGROW RX337	59.3	92.4	22.2	43.2	20993	1	2X
CARGILL 3637	59.2	103.7	24.9	36.6	21775	0	1X
N. KING N3808	58.6	•	24.5	37.1	20882	0	2X
AGRIGENE AG3920	57.3	•	22.8	40.3	19207	0	2X
NC+ 2190	57.3	•	25.9	36.2	20212	0	2X
PIONEER 3787	57.0	91.8	23.0	42.9	21217	0	1X
CIBA 4070	56.6	92.2	19.7	39.6	20993	0	2X
CIBA 4172	56.3	•	22.0	39.3	20993	0	2X
AGRIPRO AP 194	56.0	•	25.3	33.0	20658	0	2X
JOHNSON SX3952	55.1	•	20.3	40.6	20993	0	2X
ISC-PAYCO 402	55.0	•	23.6	36.9	21328	0	2X
CARGILL 2927	54.9	98.1	25.1	35.4	21663	0	1X
AGRIPRO AP 162	54.9	•	20.3	41.9	19988	0	2X
ASGROW RX406	54.4	95.7	22.9	40.1	20547	0	2X
ICI 8883	50.9	•	23.3	40.7	21105	0	2X
SEXAUER SX350	49.3	•	21.6	42.8	21775	0	2X
CIBA 4120	48.4	89.4	23.3	38.7	20770	0	2X
TOP FARM SX1195A	46.2	89.7	23.0	37.9	20993	1	2X
KAYSTAR KX-555	45.8	•	24.4	38.4	21105	1	2X
ISC-PAYCO 448	45.3	•	21.9	45.3	21328	0	2X
AGRIGENE AG3860	44.9	•	22.5	41.7	21217	0	M2X
CIBA 4202	40.4	•	22.6	38.9	20882	0	2X
TEST AVERAGE:	58.2	97.2	23.0	39.2	20963	0	
TEST LSD (5%) VALUE*:	11.3	NS**	1.6	3.4	1050	NS**	
MINIMUM BEST VALUE*:	62.4			43.5	20725		
MAXIMUM BEST VALUE*:			20.4				
TEST C.V.#:	11.9	7.6					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

Table 5. 1992 corn hybrid performance trial results--Watertown, SD, N.E. Research Farm, late maturity--96 days or more.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
STINE 992	77.6	.	28.1	40.6	21328	0	2X
N. KING N4242	77.5	.	27.2	43.1	21663	0	2X
DEKALB DK485	74.7	107.1	29.5	39.4	20882	0	1X
ISC-PAYCO 687	70.0	.	27.2	41.2	21328	0	2X
KAYSTAR KX-610	69.6	.	27.6	41.6	20993	0	2X
JACQUES 5270	69.4	.	27.0	44.6	21217	1	1X
JOHNSON SX417A	69.2	.	27.8	41.2	21552	0	2X
TOP FARM SX1097A	65.1	.	28.1	41.2	21217	0	2X
G. HARVEST H-2404	64.6	.	27.8	45.4	21105	0	1X
SIGCO 1200	64.2	.	26.5	45.1	21552	0	2X
DAHLGREN D5002	63.5	.	28.0	41.3	21328	0	2X
ASGROW RX497	62.6	.	27.7	37.9	20770	0	2X
CARGILL 4327	62.2	110.9	29.3	39.6	20658	0	1X
ISC-PAYCO 531	61.9	.	28.6	40.6	20882	0	2X
SIGCO 1301	60.2	.	28.6	43.7	20993	0	2X
G. HARVEST H-2390	59.8	102.9	27.4	43.9	20658	0	1X
STINE 1033	59.1	.	26.4	42.9	20882	0	2X
CARGILL 5327	58.0	104.7	29.0	41.0	21440	0	1X
G. HARVEST H-2322	54.7	.	28.4	39.3	21552	0	1X
ASGROW RX510	53.7	.	29.2	39.2	20100	0	2X
JACQUES 5600	53.5	.	28.2	40.5	21105	0	1X
SIGCO 1799	50.0	97.1	24.9	44.3	21440	0	3X
PIONEER 3563	49.8	97.7	25.2	43.6	21887	0	1X
DEKALB DK535	49.6	.	29.7	38.7	20658	0	1X
DAIRYLAND ST1198	49.2	.	29.2	39.8	21217	0	2X
PIONEER 3699	47.2	.	29.1	41.0	21663	0	1X
TOP FARM SX1101A	42.5	88.1	29.2	40.5	20993	0	2X
ICI 8740	40.0	.	28.3	38.0	21440	0	2X
CIBA 4282	29.2	76.9	27.9	38.3	21440	0	2X
TOP FARM SX1199	18.8	.	26.0	39.3	21663	0	2X
TEST AVERAGE:	57.6	98.2	27.9	41.2	21187	0	
TEST LSD (5%) VALUE*:	8.9	17.8	NS**	2.5	NS**	NS**	
MINIMUM BEST VALUE*:	68.7	93.1		42.9			
MAXIMUM BEST VALUE*:							
TEST C.V.#:	9.5	6.8					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

Table 6. 1992 corn hybrid performance trial results--Gary, SD, Eldon Schmidt farm, early maturity--95 days or less.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
CARGILL 3477	131.0	.	18.0	56.3	21663	0	1X
DEKALB DK462	125.4	.	17.0	55.1	21552	2	1X
ICI 8777	123.5	102.4	16.8	54.2	21552	1	2X
AGRIGENE AG5480	119.7	.	19.4	57.3	21328	1	2X
DEKALB DK381	118.3	.	14.8	56.3	21440	1	1X
PIONEER 3787	116.1	.	17.1	59.3	21552	1	1X
G. HARVEST H-2295	115.8	110.3	16.6	57.6	21440	0	1X
JACQUES 5270	114.0	.	17.9	57.6	21328	3	1X
ISC-PAYCO 687	114.0	.	18.1	57.2	21440	3	2X
SEXAUER SX540	112.8	.	17.1	55.7	21440	2	2X
PIONEER 3861	112.7	.	15.7	58.4	21440	1	1X
TOP FARM SX1102	112.5	.	17.5	59.2	21440	4	2X
N. KING N3808	111.7	.	15.8	56.9	21440	1	2X
TOP FARM SX1195A	111.6	105.7	16.6	56.8	21552	4	2X
JACQUES 4440	111.4	.	16.6	56.4	21440	1	1X
SEXAUER SX450	110.2	.	17.1	54.4	21217	1	2X
G. VALLEY GVS298	109.7	.	17.7	59.5	21440	3	2X
DEKALB DK421	109.4	.	16.2	57.7	21328	0	1X
JACQUES 5170	109.3	108.1	16.9	58.9	21328	1	1X
CARGILL 3627	108.8	102.0	18.5	58.4	21440	1	1X
ISC-PAYCO 448	108.4	.	15.8	59.8	20658	4	2X
AGRIGENE AG3920	107.2	.	17.0	57.9	21105	1	2X
ISC-PAYCO 402	106.4	.	15.0	56.8	21440	1	2X
CIBA 4172	104.3	.	14.9	54.9	21440	2	2X
KAYSTAR KX-555	103.8	.	18.4	57.9	20882	4	2X
G. VALLEY GVS282	97.0	105.4	14.9	61.4	21328	4	2X
HORIZON 4545	93.8	.	15.3	60.1	21440	1	1X
HORIZON 8095	87.2	.	15.9	59.0	20770	3	1X
TEST AVERAGE:	110.9	105.6	16.7	57.5	21352	2	
TEST LSD (5%) VALUE*:	13.0	NS**	0.8	2.5	NS**	2.5	
MINIMUM BEST VALUE*:	118.0			58.9			
MAXIMUM BEST VALUE*:			15.6			2.5	
TEST C.V.#:	7.2	8.3					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

Table 7. 1992 corn hybrid performance trial results--Gary, SD, Eldon Schmidt farm, late maturity--96 days or more.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
N. KING N4242	129.7	*	17.1	60.8	21440	0	2X
DEKALB DK485	127.6	*	18.4	57.9	21552	0	1X
PIONEER 3733	127.5	112.8	19.1	62.9	21440	0	1X
DEKALB DK512	126.6	*	18.6	57.4	21663	3	1X
CARGILL 4327	122.7	120.5	20.7	58.0	21663	1	1X
KAYSTAR KX-610	121.4	*	17.3	59.8	20882	1	2X
SIGCO 1301	121.2	*	18.4	60.9	21440	0	2X
ICI N9702	119.3	*	19.3	61.1	21440	1	2X
ISC-PAYCO 531	118.8	*	16.9	58.5	21440	1	2X
PIONEER 3563	118.0	114.1	18.3	61.6	21552	0	1X
PIONEER 3699	117.3	*	18.3	61.9	21440	0	1X
ISC-PAYCO 648	116.7	*	19.5	61.2	21440	2	2X
SIGCO 1200	116.6	*	18.1	61.1	21440	1	2X
ASGROW RX497	115.9	*	18.2	57.4	21440	1	2X
CARGILL 5327	115.5	109.1	21.9	58.2	21552	1	1X
SIGCO 1799	115.5	*	17.6	61.8	21440	3	3X
ASGROW RX510	115.1	*	20.1	57.3	20993	0	2X
TOP FARM SX1097A	113.5	*	16.3	60.4	21663	1	2X
G. HARVEST H-2390	113.3	118.4	19.8	58.1	20882	1	1X
TOP FARM SX1101A	107.0	*	18.1	61.1	20993	1	2X
G. HARVEST H-2322	103.3	*	17.6	57.8	21217	4	1X
TOP FARM SX1199	92.1	*	19.6	60.1	21440	0	2X
TEST AVERAGE:	117.0	115.0	18.6	59.8	21384	1	
TEST LSD (5%) VALUE*:	14.8	NS**	1.0	2.1	NS**	2.3	
MINIMUM BEST VALUE*:	114.9			60.8			
MAXIMUM BEST VALUE*:			17.3			2.3	
TEST C.V.#:	7.7	7.9					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

Table 8. 1992 corn hybrid performance trial results (no-till)--Frankfort, SD, Steve Masat farm, early maturity--100 days or less.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
DEKALB DK462	104.2	113.7	22.6	48.2	18760	0	1X
DEKALB DK485	99.7	115.6	27.0	43.5	18313	0	1X
ISC-PAYCO 531	96.5	.	23.6	48.0	18313	0	2X
SIGCO 1301	96.3	.	23.9	47.7	18983	0	2X
CARGILL 3477	94.8	.	22.9	46.8	19207	0	1X
N. KING N4242	93.5	.	21.1	48.6	18648	0	2X
SIGCO 1200	93.1	.	23.5	47.1	18537	0	2X
G. HARVEST H-2322	92.3	.	22.7	45.4	18872	0	1X
KAYSTAR KX-610	91.8	.	24.3	44.0	18760	0	2X
SIGCO 1799	91.5	110.4	22.4	49.4	18425	0	3X
CARGILL 3427	91.2	102.5	21.7	45.8	18313	0	1X
PIONEER 3733	89.5	103.8	23.4	48.6	19207	0	1X
DAIRYLAND ST1198	89.0	.	24.2	44.1	17978	0	2X
LEGEND 8301	88.9	.	23.0	46.5	18537	0	2X
TOP FARM SX1102	87.7	104.4	22.9	48.9	18537	0	2X
ASGROW RX510	86.7	.	28.0	42.5	18425	1	2X
STINE 1121X	85.6	.	23.2	46.4	18537	0	2X
ASGROW RX497	85.2	.	27.4	40.7	18872	0	2X
KAYSTAR KX-555	84.2	99.6	23.9	47.5	18313	0	2X
ISC-PAYCO 687	83.2	.	24.2	46.6	17978	0	2X
ISC-PAYCO 448	80.9	.	21.4	49.4	18760	0	2X
LEGEND 8203	79.4	.	25.6	45.3	18760	0	2X
CIBA 4282	79.4	97.1	24.7	44.5	18760	0	2X
PIONEER 3563	76.2	103.7	24.6	45.1	19207	0	1X
TOP FARM SX1101A	70.7	90.5	24.1	45.5	17867	0	2X
TEST AVERAGE:	88.5	104.1	23.9	46.2	18595	0	
TEST LSD (5%) VALUE*:	12.4	NS**	1.4	2.8	NS**	NS**	
MINIMUM BEST VALUE*:	91.8			46.6			
MAXIMUM BEST VALUE*:			22.5				
TEST C.V.#:	8.5	7.6					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

Table 9. 1992 corn hybrid performance trial results (no-till)--Frankfort, SD, Steve Masat farm, late maturity--101 days or more.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
DEKALB DK512	98.1	.	26.6	46.0	19318	0	1X
CARGILL 4327	92.4	107.7	29.4	44.6	18760	0	1X
PIONEER 3578	90.8	108.6	27.3	46.5	18090	0	1X
TOP FARM SX1097A	89.9	.	25.4	47.4	18648	0	2X
ISC-PAYCO 711	86.9	.	28.0	46.4	18872	0	2X
G. HARVEST H-2404	86.6	103.2	25.7	49.2	18425	0	1X
MCCURDY 5222	85.2	.	27.2	46.2	18537	0	2X
CARGILL 5327	84.5	103.0	28.8	44.7	18537	0	1X
ISC-PAYCO 648	84.2	.	25.8	49.6	18872	0	2X
TOP FARM SX1104A	84.1	105.2	26.8	48.9	18202	0	2X
G. HARVEST H-2390	83.9	101.6	24.1	49.4	18202	0	1X
HORIZON 5151	83.8	.	27.7	46.3	18648	0	1X
ASGROW RX623	82.8	.	28.1	45.6	19430	0	2X
MCCURDY 6660	82.6	.	29.5	44.2	18983	0	2X
N. KING N6560	82.5	.	29.4	45.0	18983	0	2X
DEKALB DK554	82.1	99.8	29.5	41.9	17755	0	1X
HORIZON 6101	81.1	.	26.0	48.9	17532	0	1X
PIONEER 3525	77.4	.	27.5	45.5	18872	0	1X
CIBA 4372	77.4	.	29.5	44.4	18983	0	2X
LEGEND 8104	74.7	.	26.9	47.9	18090	0	2X
N. KING N4545	73.8	88.0	28.3	42.8	18648	0	2X
SIGCO 1306	70.0	.	29.7	45.2	18313	0	2X
KAYSTAR KX-685	67.3	.	28.4	44.5	18760	0	2X
LEGEND 8205	67.3	.	28.8	44.2	17978	0	2X
STINE 1033	66.9	.	24.8	46.2	16415	0	2X
PIONEER 3503	65.5	100.7	27.4	47.7	17532	0	1X
CIBA 4393	65.5	94.6	28.7	45.6	17643	0	2X
LEGEND 9310	63.6	.	31.5	40.9	18983	0	2X
ASGROW RX653	62.0	.	28.8	43.3	18537	0	2X
DAIRYLAND ST1203	59.9	.	27.5	43.8	15075	0	2X
TOP FARM SX1199	44.7	.	25.8	47.2	18313	0	2X
TEST AVERAGE:	77.3	101.2	27.7	45.8	18321	0	
TEST LSD (5%) VALUE*:	10.5	NS**	1.3	2.3	1634	NS**	
MINIMUM BEST VALUE*:	87.6			47.3	17796		
MAXIMUM BEST VALUE*:			25.4				
TEST C.V.#:	8.3	7.1					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

**Table 10. 1992 corn hybrid performance trial results--Brookings, SD, SDSU
Agronomy Farm, early maturity--95 days or less.**

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
DEKALB DK462	108.0	131.1	23.2	46.2	21552	0	1X
DEKALB DK512	107.7	.	23.5	46.5	21552	0	1X
TERRA TR1010	104.3	120.1	24.2	51.4	21440	0	2X
CROW'S 414	97.9	.	24.5	48.9	21552	0	2X
KRUGER K9203	97.6	.	22.9	50.1	20770	0	1X
SIGCO 1200	97.5	.	23.0	51.5	21440	0	2X
ISC-PAYCO 531	97.1	.	23.5	49.6	21440	0	2X
AGRIGENE AG3920	96.9	.	21.7	51.5	21328	0	2X
PIONEER 3699	96.5	.	21.9	52.3	21663	0	1X
ISC-PAYCO 648	96.2	.	24.3	50.4	21440	0	2X
TERRA TR910	95.6	.	20.5	52.5	21552	0	2X
SEXAUER SX540	95.3	.	22.9	48.0	21217	0	2X
CARGILL 3427	94.5	114.4	24.0	47.2	21440	0	1X
KRUGER K9304	94.4	.	21.9	47.3	21552	0	1X
G. HARVEST H-2322	93.9	.	23.8	48.0	21552	0	1X
CARGILL 3637	92.8	119.4	23.6	49.4	21440	0	1X
AGRIGENE AG3980	92.6	122.3	24.4	50.5	21440	0	2X
CURRY 4411	92.4	.	22.4	48.5	20770	0	2X
SEXAUER SX620	92.0	.	21.7	49.8	21217	0	2X
SANDS SOI9991	91.9	.	23.5	46.4	21440	0	1X
YIELD KING K9303	91.5	.	25.0	44.6	21328	0	1X
DAIRYLAND ST1198	91.2	120.2	23.1	49.2	21663	0	2X
LEGEND 8301	90.3	.	21.8	48.8	21328	0	2X
PIONEER 3733	89.5	120.6	24.5	51.5	21440	0	1X
TOP FARM SX1102	89.5	.	23.3	50.5	21552	0	2X
NC+ 2844	89.4	.	25.6	47.8	21440	0	2X
DEKALB DK485	89.0	111.5	25.8	46.0	21328	0	1X
ISC-PAYCO 687	88.7	.	23.3	49.7	20882	0	2X
ICI 8777	88.5	112.2	23.3	48.8	21440	0	2X
NC+ 1991	87.6	.	23.5	48.5	21440	0	2X
KAYSTAR KX-610	86.4	.	22.0	49.2	21440	0	2X
CENEX/LOL 429	86.3	.	22.1	48.9	21440	0	1X
SIGCO 1799	86.2	112.0	22.2	50.9	21440	0	3X
AGRIPRO AP 250	86.1	.	23.7	46.2	21105	0	2X
AGRIGENE AG3860	86.0	.	21.6	52.7	21328	0	M2X
KRUGER K9204	85.7	.	24.5	51.6	21440	0	1X
JACQUES 4440	85.5	.	23.2	47.9	21440	0	1X
SIGCO 1301	85.0	.	24.2	49.4	21440	0	2X
TOP FARM SX1195A	84.8	110.9	21.5	53.6	21328	0	2X
NC+ 2190	84.0	.	23.8	47.7	21440	0	2X
HOEGEMEYER 2564	83.8	104.5	24.0	49.6	21440	0	1X
ASGROW RX497	81.7	.	24.7	45.1	21440	0	2X
ASGROW RX510	81.5	.	27.2	45.0	21328	0	2X
N. KING N3808	80.5	.	20.4	50.6	21552	0	2X
MCCURDY 4800	79.6	.	21.9	50.8	21217	0	2X
JOHNSON SX417	76.1	.	24.3	49.9	21328	0	2X
HOEGEMEYER 2583	76.1	.	27.5	44.8	21440	0	1X
ICI 8740	75.8	.	24.0	48.2	21328	0	2X
CIBA 4202	75.8	.	21.0	48.8	21217	0	2X
LEGEND 7392	74.2	.	19.8	51.3	21440	0	2X
CROW'S 237	74.1	.	27.6	43.4	21440	0	2X
JOHNSON SX3952	71.3	.	19.6	51.8	20993	0	2X
CIBA 4282	68.3	106.4	24.0	48.8	20993	0	2X
CENEX/LOL 438	59.0	.	26.4	48.4	21440	0	1X
TEST AVERAGE:	87.9	115.8	23.4	49.0	21372	0	
TEST LSD (5%) VALUE*:	13.8	NS**	1.1	2.1	NS**	0	
MINIMUM BEST VALUE*:	94.2			51.5			
MAXIMUM BEST VALUE*:			20.7			0	
TEST C.V.#:	9.7	9.0					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

**Table 11. 1992 corn hybrid performance trial results--Brookings, SD, SDSU
Agronomy Farm, late maturity--96 days or more.**

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
CARGILL 4327	110.5	130.5	27.2	48.2	21440	0	1X
ASGROW RX623	106.2	*	28.9	47.7	21440	0	2X
TOP FARM SX1097A	104.8	*	23.1	50.0	21440	0	2X
JACQUES 5270	104.3	*	23.5	50.7	21440	0	1X
AGRI-PRO AP 364	104.1	*	25.4	47.5	21552	0	2X
G. HARVEST H-2404	102.9	122.9	25.8	50.2	21440	0	1X
YIELD KING K9310	102.5	*	29.5	46.8	21440	0	1X
KRUGER K9007	102.2	129.2	26.0	50.9	21440	0	1X
CIBA 4372	101.3	*	27.7	46.0	21552	0	2X
CURRY 2142	101.1	*	26.1	48.5	21328	0	2X
G. HARVEST H-2390	99.2	124.5	26.1	49.8	21552	0	1X
SANDS SOI9061	99.0	*	28.5	47.5	21552	0	1X
N. KING N4242	98.7	*	24.3	50.0	21440	0	2X
JACQUES 4770	98.5	124.4	24.8	45.3	21440	0	1X
LEGEND 8102	98.1	125.2	25.5	50.8	21440	0	2X
AGRI-PRO AP 328	97.5	*	25.0	48.3	21440	0	2X
CURRY 4458	97.5	*	27.6	48.4	21440	0	2X
ICI 8574	97.1	105.6	28.3	46.2	21440	0	2X
HORIZON 5151	96.8	124.2	29.3	46.0	21440	0	1X
JOHNSON SX417A	96.6	*	22.7	48.1	21440	0	2X
ISC-PAYCO 711	96.1	*	28.1	47.6	21440	0	2X
PIONEER 3563	96.0	121.7	23.8	49.2	21440	0	1X
CARGILL 6227	94.3	*	30.1	44.8	21663	0	1X
TERRA TR1030	94.2	*	27.3	46.5	21440	0	2X
CARGILL 5327	93.7	126.8	28.4	48.3	21440	0	1X
CROW'S 449	93.3	121.1	30.4	44.8	21775	0	2X
NC+ 3795	93.3	*	28.3	48.6	21328	0	2X
N. KING N4545	92.6	114.1	26.6	45.2	21663	0	2X
CROW'S 370	92.4	*	26.1	46.1	21217	0	2X
YIELD KING K9306	91.5	*	27.6	47.5	21440	0	1X
TERRA TR1090	91.4	129.9	30.6	44.3	21440	0	2X
NC+ 4275	91.1	*	28.6	44.3	21105	0	2X
CIBA 4393	90.9	115.3	28.4	45.2	21328	0	2X
HORIZON 6101	90.3	*	24.7	51.8	21440	0	1X
TERRA TR1101	88.7	*	31.7	43.0	21552	0	2X
MCCURDY 4925	88.6	*	29.1	45.1	21328	0	2X
PIONEER 3525	88.3	*	25.9	49.6	20770	0	1X
LEGEND 8205	88.3	*	28.6	48.0	21440	0	2X
HOEGEMEYER 2608	88.1	*	28.7	45.5	21105	0	1X
KRUGER K9104	87.4	121.5	26.4	50.8	21440	0	1X
DEKALB DK554	87.1	*	28.5	42.3	21440	0	1X
KAYSTAR KX-685	86.3	*	28.2	46.7	21440	0	2X
ICI N9702	86.2	*	25.9	50.6	21440	0	2X
CURRY 2165	86.0	119.1	28.8	47.3	21328	0	2X
YIELD KING K9308A	85.2	*	30.5	45.5	21440	0	1X
CROW'S 401	84.3	*	31.3	42.4	21440	0	2X
DEKALB DK535	83.1	119.8	28.7	43.9	21552	0	1X
HOEGEMEYER 2611	80.4	*	27.5	46.9	21440	0	1X
ISC-PAYCO 802	79.8	*	29.0	44.7	21552	0	2X
DAIRYLAND ST1203	79.7	112.9	27.8	45.7	21440	0	2X
LEGEND 8203	75.8	*	27.3	48.1	21552	0	2X
TOP FARM SX1101A	74.6	92.5	24.5	50.2	21552	0	2X
YIELD KING K9307	70.5	*	26.4	47.8	21217	0	1X
TOP FARM SX1199	65.6	*	26.8	49.8	21105	0	2X
TEST AVERAGE:	92.1	120.1	27.3	47.3	21423	0	
TEST LSD (5%) VALUE*:	8.5	NS**	1.2	2.1	NS**	0	
MINIMUM BEST VALUE*:	102.0			49.7			
MAXIMUM BEST VALUE*:			23.9			0	
TEST C.V.#:	5.7	6.2					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

**Table 12. 1992 corn hybrid performance trial results--Beresford, SD,
S.E. Research Farm, early maturity--110 days or less.**

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
JACQUES 6970	178.5		22.8	53.1	23450	##	1X
DEKALB DK512	177.8		19.7	53.8	23673		1X
SEXAUER SX750	177.2		23.2	51.4	23338		2X
CURRY 4458	177.1		21.9	55.7	23562		2X
PIONEER 3525	176.9		20.5	57.9	23562		1X
ISC-PAYCO 802	176.7		20.9	54.7	23450		2X
NC+ 3795	174.0		22.1	55.9	23338		2X
TOP FARM SX1107	173.8		21.7	54.5	23338		2X
KAYSTAR KX-685	172.7		21.4	54.0	23450		2X
FONTANELLE 4232	172.5		22.9	51.6	23338		1X
CIBA 4372	171.9		21.0	55.8	23450		2X
ASGROW RX510	168.6		21.8	54.7	23562		2X
ASGROW RX623	167.2		21.4	56.2	23450		2X
PIONEER 3563	166.0		20.2	58.7	23338		1X
CARGILL 4327	164.3		21.9	55.4	23450		1X
CIBA 4393	163.7	120.6	21.2	54.1	23562		2X
YIELD KING K9308A	163.6		23.1	50.2	23338		1X
G. HARVEST H-2390	163.3	130.2	20.9	58.9	23450		1X
TERRA TR1030	162.8		21.5	56.7	23450		2X
HOEGEMEYER 2611	162.5		22.3	56.0	23338		1X
YIELD KING K9307	161.7		21.9	56.0	23450		1X
DEKALB DK584	161.4	129.2	22.6	55.3	23450		1X
ISC-PAYCO 711	161.4		21.8	52.6	23562		2X
NC+ 4616	161.3		23.8	53.0	23785		2X
CENEX/LOL 556	160.3		21.7	54.4	23450		1X
AGRIGENE AG6450	160.0	129.8	24.8	54.1	23450		2X
LEGEND 8205	159.9		21.8	55.2	23562		2X
HOEGEMEYER 2608	159.3		21.5	55.2	23450		1X
CARGILL 5877	158.3		21.4	53.0	23450		1X
WILSON 1140	158.1	129.3	19.7	55.4	23450		2X
YIELD KING K9306	157.8		21.0	56.8	23450		1X
N. KING N4242	157.6		19.6	55.4	23450		2X
CROW'S 237	156.2	132.9	20.9	53.2	23450		2X
DEKALB DK554	155.8	132.3	20.8	54.4	23562		1X
KAYSTAR KX-755	155.7		23.4	50.8	23450		2X
CURRY 4411	155.7		19.9	55.5	23562		2X
KRUGER K9310	155.2		22.2	57.3	23115		1X
DAIRYLAND ST1207	154.8		21.8	55.5	23450		2X
ICI N9702	154.8		20.5	58.2	23227		2X
CROW'S 179	154.6	123.1	19.8	57.4	23450		2X
NC+ 4275	154.6		22.8	50.7	23562		2X
LEGEND 8106	154.2	119.8	22.0	54.6	23450		2X
N. KING N4545	153.9		21.3	55.2	23562		2X
TERRA TR1010	153.9	128.5	20.4	57.5	23562		2X
SANDS SOI9991	152.2		19.1	55.8	23450		1X
JACQUES 7770	152.2		24.6	54.8	23338		1X
CROW'S 414	150.4		20.9	55.4	23338		2X
KRUGER K8107++	148.3	128.3	21.3	55.9	23338		1X
DAHLGREN D5057	148.3	124.5	20.3	58.4	23227		2X
FONTANELLE 3992	148.2		19.5	54.9	23450		1X
STINE 2071X	147.5		24.0	55.4	23562		2X
MCCURDY 5222	147.4		22.0	55.8	23338		2X
SEXAUER SX720	147.4		21.9	55.5	23450		2X
SIGCO 1306	143.7		21.0	54.3	23338		2X
G. HARVEST H-2404	142.8	128.0	20.2	59.0	23562		1X
CENEX/LOL 522	141.4		21.1	53.9	23338		1X
AGRIPRO AP 420	140.7		20.8	56.0	23338		2X
CURRY 2142	139.4		21.8	54.9	23562		2X
FONTANELLE 4180	137.6	116.3	20.9	57.1	23562		1X
SEXAUER SX777	137.1		25.3	52.9	23227		2X
TOP FARM SX1104A	131.1		21.7	56.1	23338		2X
KALTENBERG K5909	127.0	110.8	21.6	54.7	23450		2X
KRUGER K9007	126.1		21.4	55.2	23450		1X
TEST AVERAGE:	157.1	125.6	21.6	55.1	23443		
TEST LSD (5%) VALUE*:	16.9	NS**	0.7	2.5	NS**		
MINIMUM BEST VALUE*:	161.6			56.5			
MAXIMUM BEST VALUE*:			19.8				
TEST C.V.#:	6.6	8.0					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

##HIGH WINDS ON JULY 22, 1992 CAUSED MUCH LODGING WHICH MADE IT TOO DIFFICULT TO MAKE VALID STALK LODGING DETERMINATIONS.

**Table 13. 1992 corn hybrid performance trial results--Beresford, SD,
S.E. Research Farm, late maturity--111 days or more.**

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
PIONEER 3362	184.6	147.8	23.5	55.2	23450	##	1X
LEGEND 9113	182.5		26.2	53.8	23450	.	2X
CARCILL 7997	178.3	144.2	25.6	58.0	23450	.	1X
PIONEER 3357	176.7	143.0	25.7	54.9	23450	.	1X
WILSON E4332	175.9		22.4	57.5	23450	.	1X
HOEGEMEYER SX2628	175.7		24.1	56.1	23673	.	1X
CURRY 2187	175.5		24.2	56.0	23450	.	2X
YIELD KING K9315A	175.0		26.3	55.8	23450	.	1X
N. KING N6330	174.1		23.4	56.7	23450	.	2X
YIELD KING K9314A	173.9		24.7	55.5	23338	.	1X
CIBA 4472	173.6		24.3	56.6	23450	.	2X
KALTENBERG K7109	172.1		22.9	55.6	23338	.	2X
DEKALB DK646	171.2		24.6	56.8	23562	.	1X
ASGROW RX707	171.2		23.5	56.4	23338	.	2X
ASGROW RX653	171.1		24.0	58.0	23450	.	2X
KALTENBERG K7500	170.7	133.8	25.1	59.1	23338	.	2X
SANDS SOI9081	170.5		23.8	56.2	23450	.	1X
TERRA TR1101	168.4		24.7	55.5	23450	.	2X
STINE 1115	168.2		24.9	55.5	23450	.	2X
PIONEER 3417	166.9	133.1	22.9	57.1	23450	.	1X
PIONEER 3398	166.8	137.2	24.9	57.2	23450	.	1X
CURRY 2176	166.7	136.8	23.3	55.2	23450	.	2X
DEKALB DK623	166.7		24.8	54.4	23338	.	1X
TERRA TR1090	166.6	120.8	24.7	54.9	23450	.	2X
ICI 8532IT	166.4	130.9	24.3	57.9	23450	.	2X
ISC-PAYCO 829	164.8		24.6	56.3	23562	.	2X
G. HARVEST H-2485	164.3	128.5	23.0	56.8	23338	.	1X
TERRA TR1125	163.8	127.3	24.6	57.4	23450	.	2X
TOP FARM SX1112	163.0		24.9	57.1	23562	.	2X
ICI 8543	162.9	125.8	23.1	56.5	23562	.	2X
KRUGER K9315B	162.4		25.3	56.2	23450	.	1X
ISC-PAYCO 909	161.5		25.1	56.6	23562	.	2X
LEGEND 9310	161.3		23.2	54.9	23450	.	2X
LEGEND 9112	161.2	135.0	24.6	58.8	23450	.	2X
WILSON 1540	160.5		24.7	54.6	23338	.	
DAIRYLAND ST1116	158.5		23.4	57.5	23450	.	2X
CIBA 4432	158.3		23.4	58.8	23450	.	2X
STINE 1130	158.2		24.5	57.7	23450	.	2X
MW GENETIC G84601	157.6		26.6	56.0	23338	.	2X
CARGILL 7177	156.0		25.2	58.2	23338	.	1X
ICI 8539	155.8		22.4	58.5	23450	.	2X
FONTANELLE 1339	154.7		22.3	59.9	23338	.	1X
ISC-PAYCO 802	153.5		21.5	58.0	23450	.	2X
TOP FARM SX1113	153.0		24.3	55.6	23450	.	2X
CROW'S 498	152.8	124.7	24.9	56.5	23562	.	2X
SIGCO 1814	152.5		24.1	59.5	23450	.	1X
MW GENETIC G77501	152.2		24.0	58.3	23450	.	2X
SANDS SOI9100	151.7	125.5	25.3	56.0	23450	.	1X
G. HARVEST H-2540	151.4	127.5	24.5	58.0	23562	.	1X
HORIZON 7115	150.5	123.5	24.6	57.3	23562	.	1X
HORIZON 7878	150.3		25.4	54.2	23450	.	1X
KRUGER K8111A	150.3	126.3	24.8	57.0	23338	.	1X
CURRY 1481	150.2		24.9	58.7	23450	.	2X
HORIZON 9110	149.8	114.4	24.3	56.1	23338	.	1X
MCCURDY 6660	149.6		24.6	57.0	23338	.	2X
CROW'S 488	149.4		23.3	56.3	23562	.	2X
N. KING N6560	149.0		25.0	57.0	23450	.	2X
CIBA 4490	148.4	121.6	25.3	57.9	23450	.	2X
FONTANELLE 4280	143.7	121.1	25.0	57.1	23562	.	1X
HOEGEMEYER SX2632	141.0	123.0	25.6	57.5	23338	.	1X
DAHLGREN DC-541	139.9	120.1	27.0	56.7	23562	.	2X
TEST AVERAGE:	161.9	129.2	24.4	56.8	23448	.	
TEST LSD (5%) VALUE*:	18.1	NS**	1.2	2.3	NS**	.	
MINIMUM BEST VALUE*:	166.5			57.6		.	
MAXIMUM BEST VALUE*:			22.7			.	
TEST C.V.#:	6.9	8.1				.	

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS

16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

##HIGH WINDS ON JULY 22, 1992 CAUSED MUCH LODGING WHICH MADE IT TOO DIFFICULT TO MAKE VALID STALK LODGING DETERMINATIONS.

Table 14. 1992 corn hybrid performance trial results--Armour, SD, Robert Clark farm, early maturity--110 days or less.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR					
PIONEER 3525	159.0	.	21.6	53.6	18983	0	1X
CARGILL 4327	148.4	81.2	24.3	51.8	18760	0	1X
ISC-PAYCO 711	148.0	.	22.8	52.1	18983	0	2X
KAYSTAR KX-685	143.5	.	22.6	52.1	18537	0	2X
KRUGER K9310	143.3	.	24.0	52.2	18872	0	1X
DEKALB DK512	141.7	.	20.4	52.6	18872	0	1X
PIONEER 3563	139.7	82.0	21.6	55.7	18760	0	1X
ASGROW RX623	137.4	.	23.1	54.8	19318	0	2X
ASGROW RX510	135.2	.	23.0	51.6	17420	0	2X
CARGILL 5877	133.9	.	23.6	49.3	18983	0	1X
G. HARVEST H-2404	133.8	80.8	22.1	53.4	18872	0	1X
DEKALB DK554	133.7	83.5	24.0	48.8	18872	0	1X
HOEGEMEYER 670	132.2	.	21.4	54.0	18760	0	3X
G. HARVEST H-2390	130.5	77.1	22.4	54.0	18872	0	1X
ISC-PAYCO 648	130.0	.	22.2	54.0	19095	0	2X
KRUGER K9308A	127.3	.	23.8	51.0	18760	0	1X
TOP FARM SX1097A	127.0	75.1	20.3	52.9	18760	0	2X
CROW'S 237	126.9	81.8	22.6	50.6	18760	0	2X
ICI N9702	126.6	.	22.3	55.4	18983	0	2X
WILSON 1140	126.1	76.8	21.8	54.0	18648	0	2X
N. KING N4545	124.5	.	22.6	50.8	18760	0	2X
CROW'S 414	124.4	.	20.8	52.3	18760	0	2X
SANDS SOI9040	123.9	77.4	21.5	54.7	18760	0	1X
CIBA 4172	123.8	.	18.9	55.4	18760	1	2X
HOEGEMEYER 2571	123.5	.	19.5	53.8	18760	0	1X
ICI 8599	123.2	.	21.1	51.1	18872	0	2X
CROW'S 179	122.8	70.7	20.7	54.6	18872	0	2X
YIELD KING K9308	122.4	.	24.5	50.4	18760	0	1X
YIELD KING K9306	122.3	.	23.3	51.2	18760	0	1X
YIELD KING K9307	121.7	.	22.6	52.9	18313	0	1X
STINE 1068	119.9	.	20.9	56.0	18537	0	2X
DAHLGREN D5057	118.3	.	20.6	56.4	18425	0	2X
TOP FARM SX1199	111.6	.	21.0	56.3	18537	0	2X
HOEGEMEYER 2583	106.8	.	23.2	50.5	18760	0	1X
KAYSTAR KX-680	105.0	68.5	21.8	53.8	16415	0	2X
TEST AVERAGE:	129.1	77.7	22.1	53.0	18683	0	
TEST LSD (5%) VALUE*:	19.2	NS**					
MINIMUM BEST VALUE*:	139.8			53.8	18609		
MAXIMUM BEST VALUE*:			20.4				
TEST C.V.#:	9.1	14.6					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

Table 15. 1992 corn hybrid performance trial results--Armour, SD, Robert Clark farm, late maturity--111 days or more.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
WILSON 1540	148.4	.	25.6	49.5	19095	0	1X
ASGROW RX653	145.1	.	24.6	51.1	18760	0	2X
PIONEER 3357	137.9	85.9	25.9	51.8	19095	0	1X
ASGROW RX623	137.7	.	25.1	51.1	18760	0	2X
CARGILL 6927	136.6	85.6	25.8	52.9	18983	0	1X
WILSON E4332	136.6	.	23.7	51.9	18760	0	1X
PIONEER 3398	136.5	83.8	25.1	51.6	18872	0	1X
ISC-PAYCO 711	136.4	.	24.5	51.2	18648	0	2X
HOEGEMEYER 2681	135.6	.	27.6	49.2	18537	0	1X
YIELD KING K9315A	135.6	.	26.7	50.1	18983	0	2X
TOP FARM SX1107	134.8	80.3	23.3	51.7	18983	0	2X
KRUGER K9315B	134.5	.	25.8	50.0	18872	0	1X
STINE 1100	133.0	.	26.7	49.9	18983	0	2X
CIBA 4472	132.8	.	26.5	50.2	18983	0	2X
HORIZON 7878	129.6	.	27.8	47.8	19318	0	1X
PIONEER 3417	129.3	93.7	23.7	53.0	18983	0	1X
N. KING N6330	126.4	.	26.8	48.9	19095	0	2X
HOEGEMEYER SX2632	126.2	76.5	27.2	49.3	18983	0	1X
ISC-PAYCO 829	125.7	.	25.9	49.5	17755	0	2X
DEKALB DK623	125.6	.	26.2	50.3	18760	0	1X
YIELD KING K9314A	125.5	.	27.1	48.4	18983	0	2X
KAYSTAR KX-755	125.3	.	26.0	48.7	18648	0	2X
N. KING N6560	125.2	.	26.5	49.6	18872	0	2X
CARGILL 7177	124.9	.	26.3	49.7	18872	0	1X
G. HARVEST H-2540	124.1	.	26.1	50.1	18760	0	1X
DEKALB DK646	123.9	.	28.6	49.1	18760	0	1X
CIBA 4490	123.7	80.6	26.3	51.2	18872	0	2X
SANDS SOI9081	123.5	.	26.4	49.8	18760	0	1X
KRUGER K9316	122.4	.	27.3	52.0	18760	0	1X
TOP FARM SX1104A	121.6	.	24.7	52.9	18983	0	2X
ISC-PAYCO 802	121.4	.	24.1	49.8	18872	0	2X
STINE 1130	120.1	.	26.8	50.5	18872	0	2X
HORIZON 7115	119.0	79.2	26.4	49.7	18872	0	1X
DEKALB DK584	118.2	79.5	26.2	51.7	18872	0	1X
DAHLGREN DC-541	118.0	87.9	27.0	50.9	19095	0	2X
SANDS SOI9080	117.1	84.9	25.5	51.8	19207	0	M1X
KRUGER K9118	115.6	.	27.0	50.4	18537	0	1X
TEST AVERAGE:	128.5	83.5	26.0	50.5	18860	0	
TEST LSD (5%) VALUE*:	NS**		1.5	2.3	NS**	NS**	
MINIMUM BEST VALUE*:				50.7			
MAXIMUM BEST VALUE*:			24.8				
TEST C.V. #:	9.2	15.7					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS 16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

Table 16. 1992 corn hybrid performance trial results (irrigated no-till)--Pierre, SD, Dakota Lakes Research Farm, early maturity--100 days or less.

---- BRAND & HYBRID ----	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
N. KING N4242	77.1	.	18.4	48.1	13177	24 ##	2X
JACQUES 6970	58.5	.	24.2	39.6	3350	35	1X
TOP FARM SX1097A	56.2	120.4	20.8	44.2	13177	44	2X
DAHLGREN D5002	56.0	.	21.3	41.5	9380	34	2X
DEKALB DK512	52.3	.	25.3	34.3	20993	15	1X
ISC-PAYCO 531	50.2	.	21.7	39.3	3350	22	2X
LEGEND 8301	48.6	.	21.6	41.2	11837	45	2X
LEGEND 8104	48.2	119.7	23.5	43.2	28252	36	2X
DEKALB DK535	47.6	113.0	24.8	36.6	3350	8	1X
DEKALB DK554	46.7	116.1	25.2	35.5	12172	17	1X
G. HARVEST H-2390	44.4	107.3	22.5	40.2	12842	75	1X
ISC-PAYCO 687	44.3	.	22.7	40.5	12953	23	2X
KAYSTAR KX-610	43.6	.	22.9	38.7	11167	31	2X
CIBA 4202	41.9	.	19.7	40.5	29703	9	2X
ISC-PAYCO 711	40.7	.	24.9	38.2	3350	3	2X
PIONEER 3563	36.3	112.3	23.8	38.3	11167	16	1X
WILSON 1140	36.2	.	22.7	36.4	20658	21	1X
SEXAUER SX540	36.2	.	23.1	36.0	11725	30	2X
CARGILL 5327	35.7	119.3	25.3	37.1	21998	5	1X
JACQUES 5600	35.6	.	22.1	38.0	21887	9	1X
PIONEER 3525	34.8	.	25.1	37.9	3350	44	1X
LEGEND 8205	34.8	122.8	25.0	39.3	10943	16	2X
ASGROW RX510	34.5	.	24.8	36.7	25460	5	2X
DAIRYLAND ST1198	32.6	.	23.0	34.3	30820	18	2X
N. KING N4545	31.5	111.6	23.5	34.4	12283	16	2X
CIBA 4282	31.5	109.3	22.9	36.7	12283	18	2X
TOP FARM SX1101A	26.7	101.4	22.5	40.7	12842	39	2X
SEXAUER SX750	24.1	.	25.9	35.2	3350	34	2X
LEGEND 8203	22.6	111.0	24.1	38.3	3350	23	2X
TEST AVERAGE:	41.7	113.7	23.2	38.6	13489	25	
TEST LSD (5%) VALUE*:	16.2	NS**	1.6	4.7	NS**	32.1	
MINIMUM BEST VALUE*:	60.9			43.4			
MAXIMUM BEST VALUE*:			20.0			35.4	
TEST C.V.#:	23.8	12.3					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - THE 1992 YIELDS CONTAIN ALOT OF EXPERIMENTAL ERROR;
AND SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS. HIGH EXPERIMENTAL ERROR
WAS LIKELY THE RESULT OF REPLANTING THIS TEST ON JUNE 10TH FOLLOWING THE
SEVERE FREEZE OF MAY 25, 1992.

##EXTREMELY HIGH WINDS HIT THIS TEST TRIAL ON DECEMBER 3 RESULTING IN HIGH
STALK LODGING PERCENTAGES.

Table 17. 1992 corn hybrid performance trial results (irrigated no-till)--Pierre, SD, Dakota Lakes Research Farm, late maturity--101 days or more.

--- BRAND & HYBRID ---	YIELDS AT 15.5% MOIST.		1992 HARVEST MOIST. (%)	BUSHEL TEST WT. (LB)	FINAL POP. (AC)	STALKS LODGED (%)	TYPE CROSS
	1992 (BU/AC)	2-YR (BU/AC)					
G. HARVEST H-2493	130.5	:	21.9	49.8	12953	19 ##	1X
MCCURDY 6222	121.0	:	21.3	52.5	21887	23	2X
N. KING N6873	114.3	:	20.7	52.6	12283	86	2X
ISC-PAYCO 802	106.5	:	21.8	49.2	12395	15	2X
DEKALB DK646	103.6	:	25.2	46.8	29703	57	1X
ICI 8513	103.4		21.9	51.5	20547	71	2X
CARGILL 6227	101.9	143.6	22.2	52.3	12842	20	1X
CARGILL 5877	101.5	:	17.2	52.0	20882	76	1X
CARGILL 7697	101.3	:	25.3	50.6	3350	88	1X
ICI 8543	99.8	137.2	22.2	52.0	20212	37	2X
G. HARVEST H-2485	97.7	139.6	21.4	51.3	3350	42	1X
N. KING N6560	97.0	145.5	21.7	54.8	29815	67	2X
CIBA 4393	96.3	143.0	20.7	52.7	29145	24	2X
LEGEND 8106	96.2	:	21.1	52.0	29703	6	2X
CARGILL 6927	95.9	136.9	25.2	52.3	18983	62	1X
HORIZON 6101	92.0	:	20.6	54.5	27247	64	1X
DEKALB DK623	91.9	:	25.4	47.4	29480	50	1X
TOP FARM SX1104A	91.7	137.0	22.5	54.3	20100	34	2X
PIONEER 3357	90.6	140.6	24.0	54.0	21887	65	1X
SIGCO 1814	89.8	134.1	22.0	54.3	20658	81	2X
KAYSTAR KX-755	89.5	:	21.1	50.7	24567	48	2X
CIBA 4372	88.8		21.8	52.7	25348	46	2X
PIONEER 3398	87.4	132.7	23.0	54.2	26912	14	1X
ISC-PAYCO 841	86.7	:	18.9	51.4	25125	6	2X
HORIZON 5151	86.7	:	20.1	52.2	16303	32	1X
DAHLGREN D5092	86.3	:	19.1	52.5	24567	13	2X
KAYSTAR KX-685	85.1	131.4	20.0	53.0	15410	43	2X
ASGROW RX623	84.8	:	22.4	52.1	28587	49	2X
WILSON E4332	81.0		19.4	52.6	23450	16	1X
PIONEER 3417	80.1	133.3	19.3	53.0	22557	27	1X
DAIRYLAND ST1205	77.8	:	21.4	51.6	14963	11	2X
SIGCO 1306	77.6	:	19.9	51.7	14628	57	2X
SEXAUER SX620	73.8	:	17.4	51.5	13288	14	2X
ASGROW RX653	68.6	:	18.6	51.5	15745	4	2X
TOP FARM SX1199	64.1	:	16.5	55.7	25683	16	2X
TEST AVERAGE:	92.6	137.9	21.2	52.1	20416	39	
TEST LSD (5%) VALUE*:	20.1	NS**	3.7	2.7	NS**	37.0	
MINIMUM BEST VALUE*:	110.4			53.0			
MAXIMUM BEST VALUE*:			20.2			41.0	
TEST C.V.#:	13.3	10.2					

*SEE EXPLANATION ON USE OF TABLE ON PAGE 3.

**NS - INDICATES HYBRID DIFFERENCES WITHIN A COLUMN ARE NOT SIGNIFICANT.

#COEF. OF VARIATION - A MEASURE OF EXPERIMENTAL ERROR; IF VALUE EXCEEDS

16.0% DATA SHOULD NOT BE USED TO MAKE HYBRID COMPARISONS.

##EXTREMELY HIGH WINDS HIT THIS TEST TRIAL ON DECEMBER 3 RESULTING IN HIGH STALK LODGING PERCENTAGES.

Entries in the 1992 South Dakota corn performance trials.

COMPANY (BRAND)	HYBRID	COMPANY (BRAND)	HYBRID	COMPANY (BRAND)	HYBRID
AGRIGENE	AG3980 AG6450 AG3500 AG3860 AG3920 AG5480	DAHLGREN	DC-541 D5057 D5002 D5092 D5962	ICI (CONTINUED)	8740 8883 N9702 ISC-PAYCO
AGRIPRO	AP 364 AP 250 AP 162 AP 194 AP 420 AP 328	HEFTY SEED CO. "DAIRYLAND"	ST-1189 ST-1198 ST-1203 ST-1116 ST-1205 ST-1207		448 531 687 711 802 829 841 909
ASGROW	RX337 RX406 RX497 RX510 RX623 RX653 RX707	DEKALB	DK535 DK485 DK584 DK421 DK401 DK462 DK554 DK381 DK512	JACQUES	7770 4440 4770 5170 5270 5600 6970
CARGILL	3477 6227 6927 3627 4327 3427 3637 5327 2927 5877 7697 7977 7177	FONTANELLE JC ROBINSON "GOLDEN HARVEST"	4280 4180 3992 4232 H-2404 H-2295 H-2540 H-2390 H-2485 H-2322 H-2493	JOHNSON KALTENBERG	SX417 SX417A SX3952 K7500 K5909 K7109 KAYSTAR
CENEX/LOL	522 429 438 556	STENGEL SEED "GOLDEN VALLEY"	GVS282 GVS298	KRUGER	KX-555 KX-680 KX-685 KX-610 KX-755 K8107++ K811A
CIBA-GEIGY "CIBA"	4393 4490 4070 4120 4472 4282 4172 4202 4372 4432	HOEGEMEYER	SX2632 SX2628 2564 670 2571 2583 2608 2611 2681		K9118 K9203 K9204 K9304 K9308A K9310 K9315B K9316 LEGEND
CROW'S	488 414 237 449 498 179 370 401	HORIZON	6101 7115 8095 4545 5151 9110 7878		8102 8106 9112 7392 8203 8205 8301 9310 9313
CURRY	1481 2165 2176 2142 2187 4411	ICI	8332IT 8599 8574 8543 8777 8513 8539	MC CURDY	4925 6660 5222 6222 4800

Entries in the 1992 South Dakota corn performance trials (continued).

COMPANY (BRAND)	HYBRID	COMPANY (BRAND)	HYBRID	COMPANY (BRAND)	HYBRID
MIDWEST GENETICS	G8460 G7750	TOP FARM	SX1199 SX1102 SX1112		
NC+	4616 1991 2190 2844 3795 4275		SX1195A SX1101A SX1097A SX1104A SX1107 SX1113		
NORTHRUP KING	N-4545 N-6560 N-3808 N-4242 N-6330 N-6873	WILSON	SX1194A 1140 1540		
PIONEER	3362 3578 3788 3921 3787 3503 3417 3563 3398 3357 3525 3861 3699	YIELD KING	K9303 K9306 K9307 K9308 K9310 K9314B K9315A		
SANDS "SOI"	9040 9080 9100 9061 9081 9991				
SEXAUER	SX350 SX450 SX540 SX620 SX720 S 92-73				
SIGCO	1799 1814 1200 1301 1306				
STINE	992 1033 1068 1100 1115 1121X 1130 2071X				
TERRA	TR1125 TR1010 TR1090 TR910 TR1030 TR1101				