

1980

# Effects Of Vitamin D Metabolites On Egg Shell Characteristics Of Aged Hens

Rose Marie Wright  
*South Dakota State University*

C. W. Carlson

R. J. Emerick


Follow this and additional works at: [http://openprairie.sdstate.edu/sd\\_poultry\\_1980](http://openprairie.sdstate.edu/sd_poultry_1980)

---

## Recommended Citation

Wright, Rose Marie; Carlson, C. W.; and Emerick, R. J., "Effects Of Vitamin D Metabolites On Egg Shell Characteristics Of Aged Hens" (1980). *South Dakota Poultry Field Day Proceedings and Research Reports, 1980*. Paper 10.  
[http://openprairie.sdstate.edu/sd\\_poultry\\_1980/10](http://openprairie.sdstate.edu/sd_poultry_1980/10)

This Report is brought to you for free and open access by the Animal Science Reports at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in South Dakota Poultry Field Day Proceedings and Research Reports, 1980 by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact [michael.biondo@sdstate.edu](mailto:michael.biondo@sdstate.edu).



---

## **Effects Of Vitamin D Metabolites On Egg Shell**

### **Characteristics Of Aged Hens**

Rose Marie Wright, C. W. Carlson, And R. J. Emerick<sup>1</sup>

DEPT. OF ANIMAL SCIENCE REPORT

POULTRY 80-9

---

One hundred twenty-seven 18-month old hens were placed on a 16% protein layer diet containing no supplemental vitamin D<sub>3</sub>. Egg weight, breaking strength, and shell thickness were measured on one egg per hen each week. After 19 weeks, the hens were divided into six treatment groups. Groups 1 and 2 were fed the original diet plus 24 and 48 µg D<sub>3</sub> per kg of diet, respectively. Treatment groups 3 and 4 were supplemented with 6 and 12 µg of 1,25-(OH)<sub>2</sub>-D<sub>3</sub> per kg of diet, and groups 5 and 6 were treated with 6 and 12 µg of 24,25-(OH)<sub>2</sub>-D<sub>3</sub>. Treatment was maintained for 16 weeks. During the first 7 weeks, one egg per hen per week was taken for weight, breaking strength and shell thickness measurements and thereafter every egg was measured.

During the first 7 weeks of dietary treatment, there was an appreciable increase in all egg parameters across all treatments as shown in Table 1. Egg weight dropped through the final 9 weeks to nearly the same levels shown during the weeks on the D<sub>3</sub>-free diet. Overall, breaking strength and shell thickness levels were maintained satisfactorily throughout the duration of the experiment.

When treatment groups were compared with each other, Diets 2 and 4, the higher levels of D<sub>3</sub> and 1,25-(OH)<sub>2</sub>-D<sub>3</sub>, showed consistently significantly superior results across all three egg characteristics. However, this was only during the first 7 weeks. In the last 9 weeks, the breaking strength of the eggs from hens on Diet 2 dropped to among the lowest of the six diets, and the shell thickness of eggs from hens on Diet 4 increased slightly and remained among the highest values. The other parameters for Diets 2 and 4 dropped to moderate levels when compared to the four remaining diets.

The two "extreme" diets, the lower level of D<sub>3</sub> (Diet 1) and the higher level of 24,25-(OH)<sub>2</sub>-D<sub>3</sub> (Diet 6), showed generally poor effectiveness on improving shell strength and thickness for the first 7 weeks as compared to Diets 2 through 5. Diet 1 was effective in maintaining egg weights which compared favorably with egg weights from hens fed Diets 2 and 4.

Based on these results, there does appear to be some advantage for the use of the vitamin D metabolites in egg calcification with aged hens. Further evaluation of their effectiveness awaits analyses of the enzymes involved in calcification.

---

<sup>1</sup> Graduate Assistant in Animal Science; Professor and Leader, Poultry Research and Extension, Department of Animal Science, and Professor of Chemistry.

Table 1. Weight, breaking strength and shell thickness as affected by age and vitamin D<sub>3</sub> and its metabolites

Period <sup>1</sup>	Diet <sup>2</sup>	Egg weight g	Breaking strength kg	Shell thickness X.01 mm
A	1	63.28a	2.25a	27.97b
	2	61.44b	2.21b	27.57c
	3	62.38b	2.17c	27.70c
	4	61.62b	2.17c	27.97b
	5	62.01b	2.22b	28.26a
	6	62.47b	2.21b	27.46c
	Avg	62.20	2.21	27.82
B	1	64.76a	2.42c	30.15b
	2	64.57a	2.55a	31.06a
	3	64.88a	2.48b	31.34a
	4	65.08a	2.55a	31.21a
	5	64.06b	2.46b	30.90a
	6	64.24b	2.40c	30.16b
	Avg	64.59	2.48	30.81
C	1	64.04a	2.35d	30.14b
	2	62.71b	2.34d	29.90b
	3	62.09b	2.64a	31.53a
	4	62.44b	2.50b	31.36a
	5	61.77b	2.51b	31.74a
	6	60.59c	2.44c	30.40b
	Avg	62.27	2.47	30.84

<sup>1</sup> Periods and eggs as follows:

A = group average of one egg per hen per week for 19 weeks of D<sub>3</sub>-free diet

B = group average of one egg per hen per week for first 7 weeks of treatment

C = group average of every egg per hen for last 9 weeks of treatment.

<sup>2</sup> Dietary treatments as follows:

1 = 24 µg D<sub>3</sub> per kg diet

2 = 48 µg D<sub>3</sub> per kg diet

3 = 6 µg 1,25-(OH)<sub>2</sub>-D<sub>3</sub> per kg diet

4 = 12 µg 1,25-(OH)<sub>2</sub>-D<sub>3</sub> per kg diet

5 = 6 µg 24,25-(OH)<sub>2</sub>-D<sub>3</sub> per kg diet

6 = 12 µg 24,25-(OH)<sub>2</sub>-D<sub>3</sub> per kg diet.

a,b,c,d Means with unlike subscripts within a group are significantly different (P<0.05).