

South Dakota State University

Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

Oak Lake Field Station Research Publications

Oak Lake Field Station

1996

A Biological Assessment of Landscape Disturbance Potential on the Littoral Zone Benthos of a Prairie Pothole Lake

John L. Foley

South Dakota State University

N. H. Troelstrup Jr.

South Dakota State University, nels.troelstrup@sdstate.edu

Follow this and additional works at: https://openprairie.sdstate.edu/oak-lake_research-pubs

Recommended Citation

Foley, John L. and Troelstrup, N. H. Jr., "A Biological Assessment of Landscape Disturbance Potential on the Littoral Zone Benthos of a Prairie Pothole Lake" (1996). *Oak Lake Field Station Research Publications*. 10.
https://openprairie.sdstate.edu/oak-lake_research-pubs/10

This Article is brought to you for free and open access by the Oak Lake Field Station at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in Oak Lake Field Station Research Publications by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

A BIOLOGICAL ASSESSMENT OF LANDSCAPE DISTURBANCE POTENTIAL ON THE LITTORAL ZONE BENTHOS OF A PRAIRIE POTHOLE LAKE.

John L. Foley and Nels H. Troelstrup, Jr.
Department of Biology and Microbiology
South Dakota State University
Brookings SD 57007.

ABSTRACT

Land-use practices are known to impact adjacent water bodies. Little information exists regarding the identification of areas of disturbance around a single lake basin. Thirty-six drainages were identified on the landscape surrounding Oak Lake, Brookings County, SD. Components of the Revised Universal Soil Loss Equation (RUSLE) and land-use practices were incorporated into a GIS to delineate high and low disturbance potential for each drainage point. Littoral zone macroinvertebrates were sampled at drainage entry points around the basin to determine if changes in community structure were correlated with the disturbance potential of adjacent land areas. Preliminary analysis indicates more variety in high disturbance potential sites than reference sites, as well as some unexpected results for several metrics. For example, some high disturbance potential sites exhibit higher species diversity. This may reflect an intermediate disturbance and support the concept of a subsidy-stress gradient. Mean Macroinvertebrate Biotic Index scores (7.28 vs 6.14) and mean EPT: Chironomidae ratios (0.30 vs 1.61) for high disturbance potential vs reference sites, respectively, indicate a shift in community structure toward more tolerant species at high disturbance potential sites. These data suggest a correlation between landscape disturbance potential and adjacent littoral zone benthic communities.