

South Dakota State University  
**Open PRAIRIE: Open Public Research Access Institutional  
Repository and Information Exchange**

---

Extension Extra

SDSU Extension

---

7-1-2002

## Early Weaning Beef Calves

Kent Tjardes

*South Dakota State University*

Cody Wright

*South Dakota State University*

Don Boggs

*South Dakota State University*

Follow this and additional works at: [http://openprairie.sdstate.edu/extension\\_extra](http://openprairie.sdstate.edu/extension_extra)

---

### Recommended Citation

Tjardes, Kent; Wright, Cody; and Boggs, Don, "Early Weaning Beef Calves" (2002). *Extension Extra*. Paper 56.  
[http://openprairie.sdstate.edu/extension\\_extra/56](http://openprairie.sdstate.edu/extension_extra/56)

This Other is brought to you for free and open access by the SDSU Extension at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in Extension Extra by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact [michael.biondo@sdstate.edu](mailto:michael.biondo@sdstate.edu).



# Extension Extra

ExEx 2031  
July 2002  
Animal &  
Range  
Sciences

---

COLLEGE OF AGRICULTURE & BIOLOGICAL SCIENCES / SOUTH DAKOTA STATE UNIVERSITY / USDA

## Early Weaning Beef Calves

*Kent Tjardes, Cody Wright, and Don Boggs  
Animal and Range Sciences Department, SDSU*

Weaning calves earlier than the normal 6-7 months of age is a management practice that should be considered when pastures are short or quality is poor. Calves have been weaned successfully at less than 2 months of age. However, normal rumen function is not developed until approximately 120 days of age. Research has demonstrated that calves older than 90 days have satisfactory gains without being put on milk replacers. Calves can be placed in a drylot on a medium energy diet (NE<sub>g</sub> ≥ 0.45 Mcal/lb), on a higher quality pasture, or a pasture with supplemental creep feed.

### **Advantages:**

- Lowers the nutrient requirement for the cow to more closely match forage quality and quantity
- Increases the carrying capacity of the pasture by reducing the cow's intake
- Gives cows a better chance to regain body condition before winter
- May improve reproductive performance of cows (both this year and next)
- Improves efficiency of gain for early weaned calf
- Improves quality grade of early weaned calf

### **Disadvantages:**

- Requires more labor
- Requires facilities and feed for early weaned calves
- Increases susceptibility of early weaned calves to stress and subsequent health problems
- Increases possibility that early weaned calves may have lower carcass weights or become fatter at lighter weights

### **Considerations:**

- Complete castration, dehorning, and branding at least 10-14 days prior to weaning.
- Vaccinate calves against clostridial and viral infections prior to weaning.
- Decide the destination for calves—such as selling calves straight off the cow, backgrounding, or retained ownership.
- Consider adapting calves to creep feed prior to early weaning to aid in starting calves on feed.
- If calves are placed on a self feeder and a high quality forage, use an intake limiter in the self feeder to prevent over consumption of grain.

More detailed information on early weaning beef calves can be found at <http://ars.sdstate.edu/extbeef/>  
or contact your local Extension educator or Extension beef specialists.

This publication can be accessed electronically from the SDSU College of Agriculture & Biological Sciences  
publications page at <http://agbiopubs.sdstate.edu/articles/ExEx2031.pdf>