

South Dakota State University

Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

Agronomy, Horticulture and Plant Science
Faculty Publications

Department of Agronomy, Horticulture, and
Plant Science

4-2020

Soil Sample Timing, Nitrogen Fertilization, and Incubation Length Influence Anaerobic Potentially Mineralizable Nitrogen

Jason Clark

Kristen S. Veum

Fabian G. Fernandez

Newell R. Kitchen

James J. Camberato

See next page for additional authors

Follow this and additional works at: https://openprairie.sdstate.edu/plant_faculty_pubs



Part of the [Agronomy and Crop Sciences Commons](#)

Authors

Jason Clark, Kristen S. Veum, Fabian G. Fernandez, Newell R. Kitchen, James J. Camberato, Paul R. Carter, Richard B. Ferguson, David W. Franzen, Daniel E. Kaiser, Carrie A.M. Laboski, Emerson D. Nafziger, Carl J. Rosen, John E. Sawyer, and John F. Shanahan

Soil sample timing, nitrogen fertilization, and incubation length influence anaerobic potentially mineralizable nitrogen

Jason D. Clark¹  | Kristen S. Veum³  | Fabián G. Fernández² | Newell R. Kitchen³ |
James J. Camberato⁴ | Paul R. Carter⁵ | Richard B. Ferguson⁷ | David W. Franzen⁶  |
Daniel E. Kaiser²  | Carrie A. M. Laboski⁸ | Emerson D. Nafziger⁹ | Carl J. Rosen² |
John E. Sawyer¹⁰  | John F. Shanahan¹¹

¹South Dakota State University, 1148 Medary Ave., Brookings, SD, 57007

²University of Minnesota, 1991 Upper Buford Circle, St. Paul, MN, 55108

³USDA-ARS Cropping Systems and Water Quality Research Unit, Columbia, MO, 65211

⁴Purdue University, 915 W. State St., West Lafayette, IN, 47907

⁵Independent Agronomist, 13801 Summit Dr., Clive, IA, 50325

⁶North Dakota State University, PO Box 6050, Fargo, ND, 58108

⁷University of Nebraska, Keim 367, Lincoln, NE, 68583

⁸University of Wisconsin-Madison, 1525 Observatory Dr., Madison, WI, 53706

⁹University of Illinois, 1102 S. Goodwin, Urbana, IL, 61801

¹⁰Iowa State University, 3208 Agronomy Hall, 716 Farm House Lane, Ames, IA, 50011

¹¹Soil Health Institute, 6807 Ridge Rd., Lincoln, NE, 68512

Correspondence

J.D. Clark, South Dakota State University,
1148 Medary Ave., Brookings, SD 57007
Email: Jason.D.Clark@sdstate.edu

Funding information

DuPont Pioneer

Abstract

Understanding the variables that affect the anaerobic potentially mineralizable N (PMN_{an}) test should lead to a standard procedure of sample collection and incubation length, improving PMN_{an} as a tool in corn (*Zea mays* L.) N management. We evaluated the effect of soil sample timing (preplant and V5 corn development stage [V5]), N fertilization (0 and 180 kg ha⁻¹) and incubation length (7, 14, and 28 d) on PMN_{an} (0–30 cm) across a range of soil properties and weather conditions. Soil sample timing, N fertilization, and incubation length affected PMN_{an} differently based on soil and weather conditions. Preplant vs. V5 PMN_{an} tended to be greater at sites that received < 183 mm of precipitation or < 359 growing degree-days (GDD) between preplant and V5, or had soil C/N ratios > 9.7:1; otherwise, V5 PMN_{an} tended to be greater than preplant PMN_{an} . The PMN_{an} tended to be greater in unfertilized vs. fertilized soil in sites with clay content > 9.5%, total C < 24.2 g kg⁻¹, soil organic

Abbreviations: AWDR, Abundant and well-distributed rainfall; GDD, Growing degree-day; PMN_{an} , Anaerobic potentially mineralizable N; SDI, Shannon diversity index; SOM, Soil organic matter.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Soil Science Society of America* published by Wiley Periodicals, Inc. on behalf of Soil Science Society of America

matter (SOM) < 3.9 g kg⁻¹, or C to N ratios < 11.0:1; otherwise, PMN_{an} tended to be greater in fertilized vs. unfertilized soil. Longer incubation lengths increased PMN_{an} at all sites regardless of sampling methods. Since PMN_{an} is sensitive to many factors (sample timing, N fertilization, incubation length, soil properties, and weather conditions), it is important to follow a consistent protocol to compare PMN_{an} among sites and potentially use PMN_{an} to improve corn N management.

1 | INTRODUCTION

Nitrogen mineralization can supply 20 to 100% of crop N needs depending on several factors (Khan, Mulvaney, & Hoefl, 2001; Ros, Temminghoff, & Hoffland, 2011; Yost, Coulter, Russelle, Sheaffer, & Kaiser, 2012). Knowledge of the N supplied from soil organic matter (SOM) mineralization may improve N fertilizer guidelines. The N mineralization estimated from the PMN_{an} test was used, along with preplant- and presidedress-nitrate tests, to improve N management decisions in Argentina (Orcellet, Reussi, Sainz Rozas, Wyn-gaard, & Echeverría, 2017; Sainz Rozas, Calvino, Echeverría, Barbieri, & Redolatti, 2008). The use of the PMN_{an} test also improved the predictability of N needs of winter wheat (*Triticum aestivum* L.) in the U.S. Pacific Northwest and corn in the U.S. Southeast (Christensen & Mellbye, 2006; Williams, Crozier, White, Sripada, & Crouse, 2007). Therefore in the U.S. Midwest, the use of the PMN_{an} test may also be able to improve N guidelines for corn. However, we need to consider various sampling and methodological conditions in order to determine a standardized PMN_{an} protocol that will optimize the utility of the PMN_{an} test in predicting corn N requirements in the U.S. Midwest.

First, most soil samples collected for PMN_{an} analysis are obtained early in the spring when limited mineralization has taken place. These mineralization rates increase through the spring as temperatures increase and change throughout the remainder of the growing season (Culman, Snapp, Green, & Gentry, 2013; Fernández, Fabrizzi, & Naeve, 2017; Kuzyakova, Turyabahika, & Stahr, 2006; Sierra, 1996). However, the differences between early and later season soil and weather conditions and their influence on PMN_{an} have not been investigated. Another important aspect of soil sample timing to consider in the U.S. Midwest is that N mineralized early in the season (April to approximately mid-June) is susceptible to loss (denitrification or leaching) because of greater spring precipitation and limited N uptake by young corn (Randall & Vetsch, 2005; Struffert, Rubin, Fernández, & Lamb, 2016). Moving PMN_{an} soil sampling to later in the season when N loss potential is less and corn N uptake is increasing may improve the accuracy of the N amount that will be available to the corn crop, potentially improving the ability to predict corn N needs.

Core Ideas

- Soil parameters and weather influence how sampling time, N fertilization, and incubation length affect N mineralization.
- Nitrogen mineralization at preplant > in-season timing 27% of the time; in-season timing > pre-plant 23% of the time.
- Nitrogen fertilization reduced N mineralization 31% of the time and increased it 7% of the time.
- Sites with fine-textured soils and higher SOM had the greatest change in N mineralization from extended incubations.

Second, most soil samples collected for PMN_{an} analysis are obtained before spring N fertilizer application. However, the application of N fertilizer before soil sampling results in greater variability of N mineralization (Fernández et al., 2017; Kuzyakova et al., 2006; Ma, Dwyer, & Gregorich, 1999). Understanding the influence of N fertilizer application on N mineralization has important practical implications because most agricultural fields receive some N fertilizer before or at planting to optimize corn yield. The greater variability of N mineralization after N fertilizer application in the spring may be partially attributed to the interaction of N fertilizer with the quality of soil organic matter (i.e., C to N ratio) (Chen et al., 2014; Conde et al., 2005; Hamer & Marschner, 2005). The rate of mineralization early in the season may be more influenced by N fertilization in soils with high C to N ratios compared to soils with C to N ratios that are already low enough to promote mineralization without additional N inputs. Because of the potential for N fertilizer to influence N mineralization, mineralization estimates obtained from soil before spring N fertilization might result in an inaccurate estimate of how much N the soil can supply to a crop. Therefore, the measurement of the effect of early-season N fertilization on PMN_{an} requires further research. Increasing our understanding of the effect of N fertilizer on PMN_{an} would also likely improve N management guidelines.

Third, the standard incubation length for the PMN_{an} test is 7 d. Extending the incubation allows for more mineralization and often results in greater PMN_{an} (Angus, Ohnishi, Horie, & Williams, 1994; Smith, McNeal, Owens, & Klock, 1981). Increasing the anaerobic incubation beyond 7 d may be difficult for commercial soil testing labs that prefer high-throughput analytical methods, unless the benefits outweigh the extra costs associated with longer incubations. Clark et al. (2019) showed that PMN_{an} from longer than 7-d incubations (e.g., 14 or 28 d) related better to soil properties such as SOM, total N, and clay content in soils that have been fertilized with N and other studies observed improved correlations with crop biomass and N uptake of rice (*Oryza sativa*) with PMN_{an} from longer than 7-d incubations (Russell, Dunn, Batten, Williams, & Angus, 2006). While limited at present, those studies hint that longer incubation lengths may be more representative of N mineralization in the field, which could improve the accuracy of fertilizer-N guidelines. The potential for longer incubations to explain the variability of N mineralization in relation to contrasting soil properties and weather conditions deserves further inquiry. Therefore, the objective of this paper was to evaluate the effect of soil sample timing, N fertilization, and incubation length on PMN_{an} across a range of soil properties and weather conditions in the U.S. Midwest. Specific research findings regarding the relationships between PMN_{an} from different sampling methodologies and PMN_{an} incubation lengths with plant available N, N uptake, and yields will come in future papers.

2 | MATERIALS AND METHODS

2.1 | Experimental design

This study was conducted as a coordinated effort with uniform treatments and measurement methodology across eight U.S. Midwestern states (Illinois, Indiana, Iowa, Minnesota, Missouri, Nebraska, North Dakota, and Wisconsin). Kitchen et al. (2017) contains information regarding general soil characteristics and precipitation and temperature patterns across the study region along with specific details of experimental site descriptions, agronomic practices, and research protocols. Briefly, two sites were selected in each state in 2014 and 2015 for 32 site-years total that varied in soil properties and weather conditions (Table 1). An unfertilized check and an N rate that was considered representative of the optimal N rate (180 kg N ha^{-1}) was selected in this study for measuring PMN_{an} . Ammonium nitrate (340 g N kg^{-1}) was broadcast applied on the soil surface at planting. As stated in Kitchen et al. (2017), ammonium nitrate was used because it was expected to perform more similarly across the environmental conditions represented in the study region,

provide a uniform broadcast application that would allow for soil NO_3-N and NH_4-N evaluation shortly after application, and be suitable for surface application.

2.2 | Soil sampling and analysis

Soil characterization was performed before spring tillage and planting at each experimental site by obtaining two, 120-cm deep soil cores (3.8 to 4.0 cm i.d.) from every replicate and dividing them by horizons to measure physical and chemical properties including a taxonomic description; bulk density (bulk density-measured), soil texture, total C, total organic C, SOM, total N, cation exchange capacity, and pH (1:1 soil/water) as described in Kitchen et al. (2017). Saxton bulk density (bulk density-Saxton) was also calculated using the soil texture and SOM measurements (Saxton & Rawls, 2006). Weighted averages were calculated for the top 30 cm using the depth of each horizon within the 0- to 30-cm soil depth.

The preplant soil samples were obtained each spring 2 to 4 wk before planting and fertilization using a ten core (1.9 to 4.0 cm i.d.) composite soil sample from each replication at 0- to 30-, 30- to 60-, and 60- to 90-cm soil depths. In addition, a six-core composite (1.9-cm i.d.) soil sample (0- to 30- and 30- to 60-cm depth) was obtained at the V5 corn development stage from the 0 and 180 kg N ha^{-1} treatments. All soil samples were dried ($\leq 32^\circ\text{C}$) and ground to pass through a 2-mm sieve. Soil NO_3-N was extracted using 0.2 mol L^{-1} KCl (Saha, Sonon, & Biswas, 2018) and quantified by the cadmium reduction method (Gelderman & Beegle, 2012) with a modified Technicon AutoAnalyzer (SEAL Analytical, Inc., Fareham, UK). For PMN_{an} analysis, only the surface soils (0–30 cm in this study) were analyzed to maintain consistency with depth used when the PMN_{an} test was originally calibrated (Bundy & Meisinger, 1994). Anaerobic potentially mineralizable N was quantified by combining 4.0 g of dried soil with 20 ml of ultrapure water in 50 ml Falcon tubes (Corning Inc., Corning, NY), capped, and subjected to an incubation length of 7, 14, and 28 d at 40°C (Keeney & Bremner, 1966). After incubation, 20 ml of 4 mol L^{-1} KCl was added for a final extractant concentration of 2 mol L^{-1} KCl and samples were shaken for 30 min. Next, the solution was passed through a washed $0.45\text{-}\mu\text{m}$ syringe filter disk and stored in a microtube at -80°C to await NH_4-N analysis. Extracted NH_4-N was determined by the Berthelot method (Rhine, Mulvaney, Pratt, & Sims, 1998) using a Glomax-Multi Detection System plate reader (Promega Biosystems, Inc., Sunnyvale, CA, USA). An initial NH_4-N value was determined for each soil sample following the above extraction procedure with 2 mol L^{-1} KCl and subtracted from the incubation results to obtain net NH_4-N produced or PMN_{an} .

TABLE 1 Minimum, maximum, mean, standard deviation, and coefficient of variation of soil properties and weather conditions across 32 site-years

Property ^a	Min.	Max.	Mean	SD	CV
Soil properties					
Sand, g kg ⁻¹	20	930	260	250	950
Silt, g kg ⁻¹	40	790	500	190	390
Clay, g kg ⁻¹	20	610	240	110	470
BD-measured, g cm ⁻³	1.0	1.7	1.4	0.1	9.8
BD-Saxton, g cm ⁻³	1.1	1.6	1.3	0.1	10.0
TC, g kg ⁻¹	4.4	55.5	14.6	7.6	51.8
TOC, g kg ⁻¹	4.4	47.8	14.2	6.9	48.5
SOM, g kg ⁻¹	7.7	71.0	25.7	10.0	38.9
TN, g kg ⁻¹	0.4	4.3	1.4	0.6	41.8
C to N ratio	7.2	12.7	10.0	1.0	10.4
CEC, cmol _c kg ⁻¹	3	44	20	9	46
pH-salt	4.4	7.8	6.1	0.8	13.6
pH-water	5.1	8.8	6.7	0.8	11.4
Soil-N at PP _{0N} , mg kg ⁻¹					
NH ₄ -N 0–30 cm	3	19	8	4	44
NO ₃ -N 0–30 cm	1	18	6	3	53
NO ₃ -N 0–60 cm	1	12	5	2	42
NO ₃ -N 0–90 cm	1	9	4	2	40
Soil-N at V5 _{0N} , mg kg ⁻¹					
NH ₄ -N 0–30 cm	1	14	7	3	47
NO ₃ -N 0–30 cm	3	27	8	4	58
NO ₃ -N 0–60 cm	2	21	7	4	49
Soil-N at V5 _{180N} , mg kg ⁻¹					
NH ₄ -N 0–30 cm	2	34	9	5	63
NO ₃ -N 0–30 cm	7	75	32	12	38
NO ₃ -N 0–60 cm	9	58	24	9	35
Precipitation, Preplant-V5					
Max precipitation, mm	19	95	39	18	46
Sum of precipitation, mm	85	331	175	68	39
Mean precipitation, mm	2	5	3	0.6	17
SDI	0.5	0.7	0.6	0.1	8
AWDR	47	242	110	47	43
Temperature, Preplant-V5					
Mean max temperature, °C	19	27	22	2	8
Mean min temperature, °C	6	13	10	2	1
Mean temperature, °C	13	20	16	2	10
GDD	228	543	347	84	24

^aBD, bulk density; TC, total carbon; TOC, total organic carbon; SOM, soil organic matter; TN, total nitrogen; CEC, cation exchange capacity; SDI, Shannon diversity index; AWDR, abundant and well-distributed rainfall; GDD, growing degree-day.

2.3 | Weather

Weather data was collected at each experimental site with a HOBO U30 automatic weather station (Onset Computer Corporation, Bourne, MA, USA). Precipitation and temperature measurements were recorded every five min. These measure-

ments were used to determine the daily minimum, maximum, and mean temperatures, and the daily cumulative precipitation. These daily weather measurements were quality checked by comparing the weather station measurements against interpolated temperature data from Multi-Radar/Multi-Sensor rainfall data (National Severe Storms Lab, NOAA). Outliers

TABLE 2 Weather variables used and their definitions

Weather parameter	Definition
Mean min temperature	Tmin = Minimum daily temperature
Mean max temperature	Tmax = Maximum daily temperature
Mean temperature	MeanTemp = (Tmax + Tmin)/2
Growing degree-days	GDD = [(Tmax + Tmin)/2] - 10°C, where Tmax = Tmax if 10 ≤ Tmax ≤ 30, if Tmax ≤ 10 then Tmax = 10, if Tmax ≥ 30 then Tmax = 30; Tmin = the minimum daily temperature if Tmin ≥ 10, if Tmin ≤ 10 then Tmin = 10; all temperatures were measured in degrees Celsius, °C
Sum of precipitation	SP = Σ(Rain), where rain is the daily precipitation (mm)
Mean precipitation	MP = SP/n, where n is the number of days in that period.
Max precipitation	MP = Maximum amount of rain in a single day in that period
Shannon diversity index	SDI = [-Σpi ln(pi)]/ln(n), where pi = rain/SP is the fraction of daily precipitation relative to the total precipitation in a given time period and n is the number of days in that period. SDI = 1 implies complete evenness (i.e., equal amounts of precipitation in each day of the period); SDI = 0 implies complete unevenness (i.e., all rain in 1 d)
Abundant and well-distributed rainfall	AWDR = SP(SDI)

TABLE 3 Minimum, maximum, mean, standard deviation, and coefficient of variation of anaerobic potentially mineralizable N (PMN_{an}) as influenced by soil sample timing, N fertilization and incubation length across 32 site-years

Property ^a	PMN _{an}				
	Min.	Max.	Mean	SD	CV
	mg kg ⁻¹				
PP _{0N} , 7 d	0.7	84.0	26.7	15.1	56.8
PP _{0N} , 14 d	2.4	94.5	37.8	18.9	50.0
PP _{0N} , 28 d	6.0	125.3	48.9	25.4	51.9
V5 _{0N} , 7 d	0.2	99.9	28.3	15.0	53.1
V5 _{0N} , 14 d	2.1	122.7	37.0	17.4	47.0
V5 _{0N} , 28 d	4.0	136.7	48.5	23.2	47.8
V5 _{180N} , 7 d	0.9	92.2	23.2	15.2	65.4
V5 _{180N} , 14 d	6.9	109.9	32.4	17.5	53.9
V5 _{180N} , 28 d	8.1	130.7	43.1	23.6	54.7

^aPP_{0N}, PMN_{an} from preplant soil sampling with 0 kg N ha⁻¹; V5_{0N}, PMN_{an} from V5 corn development stage with 0 kg N ha⁻¹; V5_{180N}, PMN_{an} from V5 corn development stage with 180 kg N ha⁻¹ applied at planting.

and/or missing values were replaced by the interpolated temperature or Multi-Radar/Multi-Sensor rainfall estimates (Kitchen et al., 2017). The daily measurements were then used to calculate growing degree-days (GDD), mean precipitation, Shannon diversity index (SDI) of daily cumulative precipitation following Bronikowski and Webb (1996), and abundant and well-distributed rainfall (AWDR) following Tremblay et al. (2012) for the time period between the two soil sample timings (preplant to V5). These weather parameters were calculated using equations contained in Table 2. Water provided as irrigation in four of the 32 experimental sites was treated as natural precipitation in these calculations. These weather measurements were used to evaluate the effect of weather on PMN_{an} from the two sample timings.

2.4 | Statistical analysis

The effect of soil sample timing, N fertilization, and incubation length on PMN_{an} were evaluated using the MIXED procedure of SAS (SAS Institute Inc.). The experimental design was a randomized complete block design with four replications (blocks). Residuals within each experimental unit showed normality and constant variance assumptions were met. Block was considered a random effect. Experimental site, sample timing and N rate, incubation length, and their interactions were considered fixed effects. Least squares means were calculated for each effect and their interactions using the LSMeans statement and the differences between them were determined using Tukey's adjustment for multiple comparisons when needed. Within the three sample timing and N fertilization treatments, contrasts were used to determine the significance ($P \leq .05$) of the effect of soil sample timing (preplant vs. V5 with no N fertilization), N fertilization (0 vs. 180 kg N ha⁻¹ applied at planting and soil sampled at V5), and their interaction with site on PMN_{an} (Crossa et al., 2015). When the site by fixed effects interactions were significant, sites were evaluated individually. Soil sample timing was evaluated at only 30 sites due to missing preplant soil samples. All 32 sites were used to evaluate the effect of N fertilization and incubation length (except at two sites where incubation length was evaluated only using the V5 soil samplings due to missing preplant soil samples).

The effect of soil properties and weather conditions on the site-year to site-year differences in the effect of soil sample timing, N rate, and incubation length on PMN_{an} were evaluated using covariate analysis in the MIXED procedure of SAS (SAS Institute Inc., Cary, NC). Soil properties, weather measurements, sample timing and N rate, incubation length, and their interactions were considered fixed effects with

TABLE 4 Statistical analysis of fixed and random effects and their interactions for anaerobic potentially mineralizable N (PMN_{an}) across 32 site-years

Covariance parameters	Fixed effects			
	Numerator df	Denominator df	F-value	Pr > F
Site	31	96	14.3	<.0001
Sample timing and N rate (STNR)	2	750	34.8	<.0001
Incubation length (Inc.)	2	750	383.8	<.0001
Site × STNR	60	750	6.6	<.0001
Site × Inc.	62	750	3.9	<.0001
STNR × Inc.	4	750	0.8	0.5200
Site × STNR × Inc.	120	750	0.5	1
Contrasts				
Preplant (PP) vs. V5	1	720	1.1	0.3000
0 kg N ha ⁻¹ vs. 180 kg N ha ⁻¹	1	750	45.8	<.0001
Site × (PP vs. V5)	29	720	8.2	<.0001
Site × (0 kg N ha ⁻¹ vs. 180 kg N ha ⁻¹)	31	480	4.7	<.0001
Random effects				
	Estimate	Standard error	Z value	Pr > Z
Block (Site)	44	8.1	5.4	<.0001
Residual	106	5.5	19.4	<.0001

block, site, and site by fixed effect interactions as random effects. This analysis method allowed us to determine what soil properties and weather conditions were likely responsible for the site-year to site-year variations of the effect of soil sample timing, N fertilization, and incubation length on PMN_{an} . The slope and intercept coefficients from regressing soil characteristics and weather measurements against each PMN_{an} treatment combination were also determined using this covariate analysis. These coefficients were then used to determine the critical value of the soil or weather variables at which PMN_{an} from the preplant sample timing became greater or less than the V5 sample timing where no N fertilizer was applied, and PMN_{an} at V5 from the unfertilized soil became greater or less than the soil fertilized with 180 kg N ha⁻¹. The intercept and slope coefficients were also used to compare the effect of soil and weather variables on PMN_{an} as incubation length increased from 7 to 14 and 28 d.

3 | RESULTS AND DISCUSSION

The wide range in soil properties and weather conditions (Table 1) across all sites prior to soil sample collection led to a wide range of PMN_{an} values (0.2 to 137 mg N kg⁻¹) (Table 3). The 7 d PMN_{an} incubation results of this study (0.7 to 100 mg N kg⁻¹) were similar to other reported PMN_{an} values (12 to 87 mg N kg⁻¹) in Pennsylvania and western Oregon, USA (Christensen & Mellbye, 2006; Fox & Piekielek, 1984) and generally lower than PMN_{an} values from Argentina (71

to 222 mg N kg⁻¹) (Reussi, Rozas, Echeverría, & Berardo, 2013). Lower mean PMN_{an} in our study may be related to our overall smaller mean SOM value (25.7 g kg⁻¹) from a greater range of lower SOM values (7 to 71 g kg⁻¹) or deeper soil samples (30 cm) relative to the SOM values (44 to 68 g kg⁻¹) and shallower sampling depth (20 cm) of the Argentina study.

3.1 | Soil sample timing effect on PMN_{an}

The effect of soil sample timing on PMN_{an} varied from site-year to site-year (Table 4). Time of soil sampling did not affect PMN_{an} in 15 of the 30 sites evaluated (50%) (Figure 1; Supplemental Table S1). In the 15 sites where PMN_{an} was affected by soil sample timing, eight sites (27%) had greater PMN_{an} at V5 than preplant while in the other seven sites (23%), PMN_{an} from preplant was greater than V5 (preplant vs. V5 contrast analysis, $P \leq .05$). Soil properties and early season weather conditions influenced the effect of soil sample timing on PMN_{an} , namely precipitation amount and evenness of distribution, temperature, C to N ratio, and V5 soil NO₃-N concentration (Figure 2). The strength of the relationships between PMN_{an} from preplant and V5 with soil properties and early weather conditions shown in Figure 2 were significant but not strong (mean $R^2 = 0.05$). However, these relationships help determine how soil properties and weather conditions likely influenced the effect of sample timing on PMN_{an} .

Preplant and V5 PMN_{an} were likely to be similar at sites that received approximately 183 mm of precipitation, rainfall distribution of 0.6 SDI or 115 AWDR, or accumulated




FIGURE 1 Percentage and number of sites where anaerobic potentially mineralizable N (PMN_{an}) was affected ($P \leq .05$) by soil sample timing in the 0 kg N ha⁻¹ treatment (preplant [PP] vs V5), fertilizer-N rate applied at planting and soil sampled at V5 (0 [0N] vs. 180 kg N ha⁻¹ [180N]), and incubation length (7, 14, and 28 d) when averaged across all treatments

359 GDD between the preplant and V5 sample timings (Figure 2a–d). These were the critical values where (1) above these threshold values, PMN_{an} from V5 tended to be greater than preplant, or (2) below these threshold values, PMN_{an} from preplant tended to be greater than at V5. Greater than normal early season temperatures and more evenly distributed precipitation between the preplant and V5 soil samplings likely increased the breakdown of organic materials into more easily decomposable materials by the V5 sample timing (Cabrera, Kissel, & Vigil, 2005; Culman et al., 2013; Fernández et al., 2017; Goulding et al., 1998; Kuzyakov, Friedel, & Stahr, 2000; Ma & Wu, 2008). This greater abundance of easily decomposable material available at V5 sampling likely led to the increase of V5 PMN_{an} over preplant PMN_{an} .

The C to N ratio and V5 soil NO_3-N concentration also influenced the effect of sample timing on PMN_{an} (Figure 2e,f). The critical values where PMN_{an} from preplant and V5 were similar were 9.7:1 for C to N ratio and 8.2 mg kg⁻¹ for V5 soil NO_3-N . Specifically, the PMN_{an} from preplant tended to be greater than V5 at C to N ratios and V5 soil NO_3-N values above these critical values and PMN_{an} from preplant tended to be greater than V5 below




FIGURE 2 Anaerobic potentially mineralizable N (PMN_{an}) from a 7-d incubation that was soil sampled before planting (PP) and at the V5 corn development stage as a function of soil properties (a) and weather conditions (b to e). Critical values represent the intersection point where PMN_{an} from the preplant and V5 sample timing became greater or less than the other. Only those weather conditions and soil properties that had a significant interaction ($P \leq .05$) with soil sample timing were included

these critical values. These results indicate PMN_{an} is not consistent throughout the growing season and that soil and weather conditions influence the effect soil sample collection timing has on PMN_{an} . Therefore, the timing of obtaining soil samples to complete PMN_{an} analysis should remain consistent from year to year to make appropriate comparisons. In addition, because PMN_{an} is sensitive to sample timing, further research is needed to determine the timing of soil sampling that best relates to crop N requirement before a standard protocol can be recommended.

3.2 | Nitrogen fertilization effect on PMN_{an} at the V5 corn development stage

The effect of N fertilization on PMN_{an} varied from site-year to site-year (Table 4). Nitrogen fertilization did not affect PMN_{an} in 20 of the 32 sites evaluated (63%) (Figure 2; Supplemental Table S1). In the 12 sites where N fertilization affected PMN_{an} , 10 sites (31%) had greater PMN_{an} from unfertilized compared to fertilized soil while in the other two sites (6%), PMN_{an} from fertilized soil was greater than unfertilized soil (0 vs. 180 kg N ha⁻¹ applied at planting and soil sampled at V5 contrast analysis, $P \leq .05$). These results indicate that N fertilization does not consistently influence PMN_{an} and when it does, it most often reduces PMN_{an} . The variable effect of N fertilization on PMN_{an} in this study is similar to the findings of others (Fernández et al., 2017; Kuzyakova et al., 2006; Ma et al., 1999). Furthermore, soil properties influenced the effect of N fertilization on PMN_{an} , namely total C, total organic C, SOM, C to N, clay content, and V5 soil NO₃-N concentration (Figure 3). The strength of the relationships between PMN_{an} from fertilized and unfertilized soil with soil properties shown in Figure 3 were significant but not strong (mean $R^2 = 0.16$). However, similar to the N timing evaluations, these relationships help determine how soil properties likely influenced the effect of N fertilization on PMN_{an} .

The PMN_{an} from unfertilized soil was generally greater than fertilized soil at sites with low amounts of total C, total organic C, SOM, or C to N ratio (Figure 3a–d). The reduction in PMN_{an} from fertilized relative to unfertilized soil is likely the result of the N fertilizer stimulating mineralization of the labile organic matter in the soil and depleting the amount of SOM available for mineralization by the V5 sample timing (Chen et al., 2014; Conde et al., 2005; Hamer & Marschner, 2005; Kuzyakov et al., 2000). The differences in PMN_{an} due to N fertilization became less pronounced as total C, total organic C, SOM, or C to N ratio increased toward the high end of the range measured across the sites. The similarity in PMN_{an} values from fertilized and unfertilized soil with these characteristics is likely the result of a reduction in the stimulation of N mineralization from the addition of N fertilizer as soil C content increased, as reported in other studies (Chen et al., 2014; Conde et al., 2005). Since only two sites

had statistically greater PMN_{an} from fertilized compared to unfertilized soils, it is difficult to establish what soil parameters or critical values may help explain this response. We observed only a trend, suggesting that PMN_{an} from fertilized relative to unfertilized soil became greater when total C, total organic C, SOM, or the C to N ratio increased above 24.2 g kg⁻¹, 21.1 g kg⁻¹, 37.9 g kg⁻¹, 11.0:1, respectively.

The clay content and V5 soil NO₃-N concentrations also influenced the similarities and differences between PMN_{an} from unfertilized and fertilized soil (Figure 3e,f). The PMN_{an} from unfertilized relative to fertilized soil was generally greater at those sites with the greatest amounts of clay content and V5 soil NO₃-N concentrations. The PMN_{an} from unfertilized and fertilized soils became similar as clay content and V5 soil NO₃-N decreased toward the low end of the range measured in our study. These results indicate N fertilization can affect PMN_{an} and that soil properties influenced the effect N fertilizer application has on PMN_{an} . Therefore, soil samples collected for PMN_{an} analysis should always be obtained before or after N fertilization from year to year to make appropriate comparisons. In addition, because PMN_{an} is sensitive to N fertilization, further research is needed to determine whether PMN_{an} from fertilized or unfertilized soil best relates to crop N requirements before a standard protocol can be recommended.

3.3 | Incubation length effect on PMN_{an}

Extending the incubation length beyond 7 d generally increased PMN_{an} at all sites (Figure 1; Supplemental Table S1), which is similar to the findings of Angus et al. (1994). The magnitude of the increase in PMN_{an} with longer incubations varied from site to site (Table 4), depending on soil properties such as silt and clay content, cation exchange capacity, total C, total organic C, SOM, total N, or preplant NH₄-N concentration (30-cm depth) (Table 5). The greater PMN_{an} from longer incubations increased (greater slope and intercept values) as these soil properties increased across the sites. In contrast to this result, PMN_{an} increased at a reduced rate (reduced slope but greater intercept values) with longer incubation lengths as sand content or bulk density values increased across the sites. Precipitation and temperature did not impact the effect of incubation length on PMN_{an} . Cation exchange capacity, total C, total organic C, SOM, total N, and bulk density were the soil measurements that interacted with incubation length and accounted for the greatest variation in PMN_{an} (mean F-value of 13) (Table 5). These soil properties also reduced the estimate of variance the most for site (mean decrease = 73) and the site by incubation length interaction (mean decrease = 12) (Supplemental Table S3). All other significant interactions between incubation length and soil variables had a weaker influence on PMN_{an} (mean F-value of 4.7) (Table 5). These results indicate that cation exchange




FIGURE 3 Anaerobic potentially mineralizable N (PMN_{an}) from a 7-d incubation that was soil sampled at the V5 corn development stage where 0 (0N) or 180 kg N ha^{-1} (180N) was applied as a function of soil properties (a to f). Critical values represent the intersection point where PMN_{an} from the unfertilized and fertilized soil became greater or less than the other. Only those soil properties that had a significant interaction ($P \leq .05$) with N fertilization were included

TABLE 5 Change in slope and intercept of anaerobic potentially mineralizable N (PMN_{an}) as a function of soil properties when incubation length increased from 7 to 14 and 7 to 28 d. Only those soil properties that had a significant interaction ($P \leq .05$) with incubation length were included

Variable ^a	Change in slope coefficient from 7-d incubation		Change in intercept from 7-d incubation		F-value
	14 d	28 d	14 d	28 d	
Sand, g kg^{-1}	-0.01 ^b	-0.02*	+12*	+27*	9*
Silt, g kg^{-1}	+0.01	+0.02*	+4	+10*	4*
Clay, g kg^{-1}	+0.02	+0.05*	+6	+10*	8*
BD-measured, g cm^{-3}	-8.23	-33.50*	+21	+67*	9*
BD-Saxton, g cm^{-3}	-19.89*	-51.73*	+36*	+90*	16*
TC, g kg^{-1}	+0.23	+0.62*	+6*	+12*	8*
TOC, g kg^{-1}	+0.28	+0.79*	+6*	+10*	11*
SOM, g kg^{-1}	+0.22	+0.60*	+4	+5	14*
TN, g kg^{-1}	+3.48	+9.28*	+5	+8*	11*
CEC, $cmol_c kg^{-1}$	+0.22	+0.63*	+5	+9*	10*
$PP_{0N} NH_4-N$, 0-30 cm	+0.14	+1.00*	+8*	+12*	4*

*Significant at $P \leq .05$.

^aBD, bulk density; TC, total carbon; TOC, total organic carbon; SOM, soil organic matter; TN, total nitrogen; CEC, cation exchange capacity.

^bChange in slope coefficient and intercept when moving from 7 to 14 or 28 d of incubation. (Sand content example: 7 d PMN_{an} = (slope coefficient)(sand content) + intercept. When PMN_{an} incubation length moves from 7 to 14 d, the slope coefficient decreases by 0.01 and the intercept increases by 12.

capacity, total C, total organic C, SOM, total N, and bulk density were likely the soil properties that were driving most of the differences in the increase of PMN_{an} with longer incubations from site to site.

4 | CONCLUSIONS

Soil properties (especially cation exchange capacity, soil C content, SOM, total N, and bulk density) and early season weather conditions (especially evenness of early season precipitation) had a large influence on the effect of soil sample timing, N fertilizer application, and incubation length on PMN_{an} . Therefore, careful consideration as to the time of soil sampling, N fertilization status, and incubation length should be made when comparing PMN_{an} values among sites. Producers and scientists should follow a consistent protocol when obtaining soil samples and analyzing them for PMN_{an} to make comparisons related to N mineralization capacity of soils and for guiding fertilizer-N rates. There are tradeoffs with the sampling methodologies and incubation lengths evaluated in this study. For example, commercial soil testing labs may not want to incubate soil samples for 28 d because they prefer high-throughput analytical methods that reduce costs and provide rapid results to producers. Therefore, a better understanding of the relationship between crop N availability, N uptake, yields, and PMN_{an} from these different soil sampling methodologies and incubation lengths are needed before we can determine the protocol that best relates to crop N management.

ACKNOWLEDGMENTS

We thank DuPont Pioneer for funding this research. The authors thank the supporting scientists (Matt Yost; Dan Barker [Iowa]; Lakesh Sharma, Amitava Chatterjee, and Norm Cattanaach [North Dakota]; Todd Andraski [Wisconsin]; and Tim Hart [DuPont Pioneer]), field technicians (Matt Volkmann [Missouri]; Jason Niekamp and Joshua Vonk [Illinois]; Glen Slater [Nebraska]; Andrew Scobbie, Thor Sellie, Nicholas Severson, Darby Martin, and Erik Joerres [Minnesota]), and cooperating farmers and research farm personnel for their help in completing this project. Mention of trade names or commercial products in this publication is solely for the purpose of providing information and does not imply recommendation or endorsement by the affiliated Universities or the U.S. Department of Agriculture.

ORCID

Jason D. Clark  <https://orcid.org/0000-0001-7793-6411>
 Kristen S. Veum  <https://orcid.org/0000-0002-6492-913X>
 David W. Franzen  <https://orcid.org/0000-0003-4862-8086>
 Daniel E. Kaiser  <https://orcid.org/0000-0002-8478-701X>
 John E. Sawyer  <https://orcid.org/0000-0003-4080-9616>

REFERENCES

- Angus, J. F., Ohnishi, M., Horie, T., & Williams, R. L. (1994). A preliminary study to predict net nitrogen mineralization in a flooded rice soil using anaerobic incubation. *Australian Journal of Experimental Agriculture*, *34*, 995–999.
- Bronikowski, A., & Webb, C. (1996). A critical examination of rainfall variability measures used in behavioral ecology studies. *Behavioral Ecology and Sociobiology*, *39*, 27–30. <https://doi.org/10.1007/s002650050263>
- Bundy, L. G., & Meisinger, J. J. (1994). Nitrogen availability indices. In R. W. Weaver (Ed.), *Methods of soil analysis: Biochemical and microbial properties* (pp. 951–984). SSSA Monogr. 5. Madison, WI: Soil Science Society of America.
- Cabrera, M. L., Kissel, D. E., & Vigil, M. F. (2005). Nitrogen mineralization from organic residues: Research opportunities. *Journal of Environmental Quality*, *34*, 75–79. <https://doi.org/10.2134/jeq2005.0075>
- Chen, R., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., Lin, X., ... Kuzyakov, Y. (2014). Soil C and N availability determine the priming effect: Microbial N mining and stoichiometric decomposition theories. *Global Change Biology*, *20*, 2356–2367. <https://doi.org/10.1111/gcb.12475>
- Christensen, N. W., & Mellbye, M. E. (2006). Validation and recalibration of a soil test for mineralizable nitrogen. *Communications in Soil Science and Plant Analysis*, *37*, 2199–2211. <https://doi.org/10.1080/00103620600817416>
- Clark, J. D., Veum, K. S., Fernández, F. G., Camberato, J. J., Carter, P. R., Ferguson, R. B., ... Shanahan, J. F. (2019). United States Midwest soil and weather conditions influence anaerobic potentially mineralizable nitrogen. *Soil Science Society of America Journal*, *83*, 1137–1147. <https://doi.org/10.2136/sssaj2019.02.0047>
- Conde, E., Cardenas, M., Ponce-Mendoza, A., Luna-Guido, M. L., Cruz-Mondragón, C., & Dendooven, L. (2005). The impacts of inorganic nitrogen application on mineralization of ¹⁴C-labelled maize and glucose, and on priming effect in saline alkaline soil. *Soil Biology & Biochemistry*, *37*, 681–691. <https://doi.org/10.1016/j.soilbio.2004.08.026>
- Crossa, J., Vargas, M., Cossani, C. M., Alvarado, G., Burgueño, J., Mathews, K. L., & Reynolds, M. P. (2015). Evaluation and interpretation of interactions. *Agronomy Journal*, *107*, 736–747. <https://doi.org/10.2134/agronj2012.0491>
- Culman, S. W., Snapp, S. S., Green, J. M., & Gentry, L. E. (2013). Short- and long-term labile soil carbon and nitrogen dynamics reflect management and predict corn agronomic performance. *Agronomy Journal*, *105*, 493–502. <https://doi.org/10.2134/agronj2012.0382>
- Fernández, F. G., Fabrizzi, K. P., & Naeve, S. L. (2017). Corn and soybean's season-long in-situ nitrogen mineralization in drained and undrained soils. *Nutrient Cycling in Agroecosystems*, *107*, 33–47. <https://doi.org/10.1007/s10705-016-9810-1>
- Fox, R. H., & Piekielek, W. P. (1984). Relationships among anaerobically mineralized nitrogen, chemical indexes, and nitrogen availability to corn. *Soil Science Society of America Journal*, *48*, 1087–1090. <https://doi.org/10.2136/sssaj1984.03615995004800050027x>
- Gelderman, R. H., & Beegle, D. (2012). Nitrate-nitrogen. Recommended chemical soil test procedures for the North Central Region. *North Central Regional Res. Publ. no. 221* (revised Oct 2012). Columbia, MO: Missouri Agricultural Experiment Station.
- Goulding, K. W. T., Bradbury, N. J., Hargreaves, P., Howe, M., Murphy, D. V., Poulton, P. R., & Willison, T. W. (1998). Nitrogen

- deposition and its contribution to nitrogen cycling and associated soil processes. *New Phytologist*, *139*, 49–58. <https://doi.org/10.1046/j.1469-8137.1998.00182.x>
- Hamer, U., & Marschner, B. (2005). Priming effects in different soil types induced by fructose, alanine, oxalic acid and catechol additions. *Soil Biology & Biochemistry*, *37*, 445–454. <https://doi.org/10.1016/j.soilbio.2004.07.037>
- Keeney, D. R., & Bremner, J. M. (1966). Comparison and evaluation of laboratory methods of obtaining an index of soil nitrogen availability. *Agronomy Journal*, *58*, 498–503. <https://doi.org/10.2134/agronj1966.00021962005800050013x>
- Khan, S. A., Mulvaney, R. L., & Hoefl, R. G. (2001). A simple soil test for detecting sites that are nonresponsive to nitrogen fertilization. *Soil Science Society of America Journal*, *65*, 1751–1760. <https://doi.org/10.2136/sssaj2001.1751>
- Kitchen, N. R., Shanahan, J. F., Ransom, C. J., Bandura, C. J., Bean, G. M., Camberato, J. J., ... Shafer, M. (2017). A public-industry partnership for enhancing corn nitrogen research and datasets: Project description, methodology, and outcomes. *Agronomy Journal*, *109*, 2371–2388. <https://doi.org/10.2134/agronj2017.04.0207>
- Kuzyakov, Y., Friedel, J. K., & Stahr, K. (2000). Review of mechanisms and quantification of priming effects. *Soil Biology & Biochemistry*, *32*, 1485–1498. [https://doi.org/10.1016/S0038-0717\(00\)00084-5](https://doi.org/10.1016/S0038-0717(00)00084-5)
- Kuzyakova, I. F., Turyabahika, F. R., & Stahr, K. (2006). Time series analysis and mixed models for studying the dynamics of net N mineralization in a soil catena at Gondelsheim (S-W Germany). *Geoderma*, *136*, 803–818. <https://doi.org/10.1016/j.geoderma.2006.06.003>
- Ma, B. L., Dwyer, L. M., & Gregorich, E. G. (1999). Soil nitrogen amendment effect on seasonal nitrogen mineralization and nitrogen cycling in maize production. *Agronomy Journal*, *91*, 1003–1009. <https://doi.org/10.2134/agronj1999.9161003x>
- Ma, B. L., & Wu, T. Y. (2008). Plant-available nitrogen in the soil: Relationships between pre-plant and pre-sidedress nitrate tests for corn production. *Journal of Plant Nutrition and Soil Science*, *171*, 458–465. <https://doi.org/10.1002/jpln.200700091>
- Orcellet, J., Reussi, Calvo, N. I., Sainz Rozas, H. R., Wyngaard, N., & Echeverría, H. E. (2017). Anaerobically incubated nitrogen improved nitrogen diagnosis in corn. *Agronomy Journal*, *109*, 291–298. <https://doi.org/10.2134/agronj2016.02.0115>
- Randall, G. W., & Vetsch, J. A. (2005). Nitrate losses in subsurface drainage from a corn-soybean rotation as affected by fall and spring application of nitrogen and nitrapyrin. *Journal of Environmental Quality*, *34*, 590–597. <http://www.ncbi.nlm.nih.gov/pubmed/15758112>
- Reussi, Calvo, N. I., Rozas, H. Sainz, Echeverría, H., & Berardo, A. (2013). Contribution of anaerobically incubated nitrogen to the diagnosis of nitrogen status in spring wheat. *Agronomy Journal*, *105*, 321–328. <https://doi.org/10.2134/agronj2012.0287>
- Rhine, E. D., Mulvaney, R. L., Pratt, E. J., & Sims, G.K. (1998). Improving the Berthelot Reaction for determining ammonium in soil extracts and water. *Soil Science Society of America Journal*, *62*, 473–480. <https://doi.org/10.2136/sssaj1998.03615995006200020026x>
- Ros, G. H., Temminghoff, E. J. M., & Hoffland, E. (2011). Nitrogen mineralization: A review and meta-analysis of the predictive value of soil tests. *European Journal of Soil Science*, *62*, 162–173. <https://doi.org/10.1111/j.1365-2389.2010.01318.x>
- Russell, C. A., Dunn, B. W., Batten, G. D., Williams, R. L., & Angus, J. F. (2006). Soil tests to predict optimum fertilizer nitrogen rate for rice. *Field Crops Research*, *97*, 286–301. <https://doi.org/10.1016/j.fcr.2005.10.007>
- Saha, U. K., Sonon, L., & Biswas, B. K. (2018). A comparison of diffusion-conductimetric and distillation-titration methods in analyzing ammonium- and nitrate-nitrogen in the KCl-extracts of Georgia soils. *Communications in Soil Science and Plant Analysis*, *49*, 63–75. <https://doi.org/10.1080/00103624.2017.1421647>
- Sainz Rozas, H., Calvo, P. A., Echeverría, H. E., Barbieri, P. A., & Redolatti, M. (2008). Contribution of anaerobically mineralized nitrogen to the reliability of planting or presidedress soil nitrogen test in maize. *Agronomy Journal*, *100*, 1020–1025. <https://doi.org/10.2134/agronj2007.0077>
- Saxton, K. E., & Rawls, W. J. (2006). Soil water characteristic estimates by texture and organic matter for hydrologic solutions. *Soil Science Society of America Journal*, *70*, 1569–1578. <https://doi.org/10.2136/sssaj2005.0117>
- Sierra, J. (1996). Nitrogen mineralization and its error of estimation under field conditions related to the light-fraction soil organic matter. *Australian Journal of Soil Research*, *34*, 755–767. <https://doi.org/10.1071/SR9960755>
- Smith, J. L., McNeal, B. L., Owens, E. J., & Klock, G. O. (1981). Comparison of nitrogen mineralized under anaerobic and aerobic conditions for some agricultural and forest soils of Washington. *Communications in Soil Science and Plant Analysis*, *12*, 997–1009.
- Struffert, A. M., Rubin, J. C., Fernández, F. G., & Lamb, J. A. (2016). Nitrogen management for corn and groundwater quality in Upper Midwest irrigated sands. *Journal of Environmental Quality*, *45*, 1557–1564. <https://doi.org/10.2134/jeq2016.03.0105>
- Tremblay, N., Bouroubi, Y. M., Bélec, C., Mullen, R. W., Kitchen, N. R., Thomason, W. E., ... Ortiz-Monasterio, I. (2012). Corn response to nitrogen is influenced by soil texture and weather. *Agronomy Journal*, *104*, 1658–1671. <https://doi.org/10.2134/agronj2012.0184>
- Williams, J. D., Crozier, C. R., White, J. G., Sripada, R. P., & Crouse, D. A. (2007). Comparison of soil nitrogen tests for corn fertilizer recommendations in the humid southeastern USA. *Soil Science Society of America Journal*, *71*, 171–180. <https://doi.org/10.2136/sssaj2006.0057>
- Yost, M. A., Coulter, J. A., Russelle, M. P., Sheaffer, C. C., & Kaiser, D. E. (2012). Alfalfa nitrogen credit to first-year corn: Potassium, regrowth, and tillage timing effects. *Agronomy Journal*, *104*, 953–962. <https://doi.org/10.2134/agronj2011.0384>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Clark JD, Veum KS, Fernández FG, et al. Soil sample timing, nitrogen fertilization, and incubation length influence anaerobic potentially mineralizable nitrogen. *Soil Sci. Soc. Am. J.* 2020;84:627–637. <https://doi.org/10.1002/saj2.20050>