South Dakota State University

Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

Electronic Theses and Dissertations

1986

Effects of Prairie Dogs on Mound Soils

Daniel C. Carlson

Follow this and additional works at: <https://openprairie.sdstate.edu/etd>

Recommended Citation

Carlson, Daniel C., "Effects of Prairie Dogs on Mound Soils" (1986). *Electronic Theses and Dissertations*. 4352.

<https://openprairie.sdstate.edu/etd/4352>

This Thesis - Open Access is brought to you for free and open access by Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

EFFECTS OF PRAIRIE DOGS ON MOUND SOILS

BY

DANIEL C. CARLSON

A thesis submitted in partial fulfillment
of the requirements for the degree
Master of Science
Major in Soil Science
South Dakota State University
1986

EFFECTS OF PRAIRIE DOGS ON MOUND SOILS

This thesis is approved as a creditable and independent investigation by a candidate for the degree, Master of Science, and is acceptable for meeting the thesis requirements for this degree. Acceptance of this thesis does not imply that the conclusions reached by the candidate are necessarily the conclusions of the major department.

Everett M. White
Major and Thesis Adviser

Date

Maurice L Horton
Dept. Head, Plant Sci.

Date

ACKNOWLEDGEMENTS

The author wishes to thank Dr. Everett M. White, Major and Thesis Adviser and Professor in Soil Science, for his continued interest in and support of this study and for the hours of time he spent directing and assisting my efforts. Without him, this study would not have been possible.

I am also grateful to the National Park Service, Wind Cave National Park, and the South Dakota State University Plant Science Department for financial and technical assistance provided for the study.

In addition, Research Biologist Richard Klukas of the National Park Service, Professors Doug D. Malo, Gary D. Lemme, Greg Carlson, and the faculty, staff, and research assistants at South Dakota State University are remembered for their part in an enjoyable tenure here pursuing the M.S. degree.

To my caring and loving wife, Laura, and family... Thank you for being there for me, even when I wasn't. I am indebted to all of you.

TABLE OF CONTENTS

	<u>PAGE</u>
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF FIGURES	vi
LIST OF TABLES	vii
INTRODUCTION	1
REVIEW OF LITERATURE	3
A. Geography and Climate	3
B. Vegetation	4
C. Geology	6
D. History of Fire and Extent of Forest in the Black Hills	7
E. Phosphorus Accumulation in Soil	10
F. Prairie Dogs	12
DESCRIPTION OF THE STUDY AREAS	15
METHODS AND MATERIALS	21
A. Sampling and Laboratory Procedures (1984)	21
B. Sampling and Laboratory Procedures (1985)	22
C. Statistical Methods	26
D. Mound Area Relationships	27
E. Phosphorus	29
RESULTS AND DISCUSSION	31
Soil Changes Within Mounds (1984)	31
A. Soil Color	31

B. Nitrogen	37
C. Soil pH	41
D. Total Phosphorus	45
Soil Changes Across Mounds (1985)	60
A. Soil Color	60
B. Soil Texture	71
C. Soil pH	77
D. Total Phosphorus in Surface Layers	84
Mound Formation Rate	91
General Relationships	97
SUMMARY AND CONCLUSIONS	106
BIBLIOGRAPHY	109

LIST OF FIGURES

<u>FIGURE</u>	<u>PAGE</u>
A. Location of Mound Sites in Wind Cave National Park	16
1. Subsample Locations Within Mound Sampling Rings	23
2. Soil Colors at Mound 1	32
3. Soil Colors at Mound 2	33
4. Soil Colors at Mound 3	35
5. Nitrogen Contents at Mound 1	38
6. Nitrogen Contents at Mound 2	39
7. Nitrogen Contents at Mound 3	40
8. pH Values at Mound 1	42
9. pH Values at Mound 2	43
10. pH Values at Mound 3	44
11. Total Phosphorus Contents at Mound 1	46
12. Total Phosphorus Contents at Mound 2	47
13. Total Phosphorus Contents at Mound 3	50
14. Relationship Between Total and Disturbed Soil Area Within Mounds	93
15. Scale of Relative Mound Ages Based on Total Area of Disturbed Soil at each Mound	96

LIST OF TABLES

<u>TABLE</u>	<u>PAGE</u>
1. Summary of Laboratory Analyses for Mound 1--ENE Trench (1984)	52
2. Summary of Laboratory Analyses for Mound 1--WSW Trench (1984)	53
3. Summary of Laboratory Analyses for Mound 2--ENE Trench (1984)	54
4. Summary of Laboratory Analyses for Mound 2--WSW Trench (1984)	55
5. Summary of Laboratory Analyses for Mound 3 (1984)	56
6. Summary of Laboratory Analyses for Mound 3 (1984)	57
7. Summary of Laboratory Analyses for Mound 3 (1984)	58
8. Ranges and Means of Soil Properties Measured for Mounds 1-3 (1984)	59
9. Classification of Soils Sampled for Prairie Dog Mound Analysis	61
10. Studied Soils and Mound Locations	62
11. Munsell Soil Colors from Mound and Adjacent Surface Soil	64
12. Soil Textures from Mound and Adjacent Surface Soil	72
13. Average 1:1 pH of the 0-10 cm layer of Prairie Dog Mounds with Increasing Distance from the Mound Center	78
14. Analysis of Variance for Average 1:1 pH of Mound Surfaces Compared to Adjacent Non-Mound Soil	79
15. Analysis of Variance for Average 1:1 pH values Between Selected Mounds From the Same Study Area	83

16.	Mean Total Phosphorus Content of the 0-10 cm layer of Prairie Dog Mounds with Increasing Distance from the Mound Center	85
17.	Statistical Analysis of Mean Total Phosphorus Contents of Mound Surfaces Compared to Non-Mound Soil	86
18.	Orthogonal Comparison of Mean Total Phosphorus Content Between Selected Mound and Undisturbed Surface Soils	88
19.	Total Mound Surface Area and Area of Disturbed Original Soil Surface	92

INTRODUCTION

Biological pedoturbation is a process of soil formation which has not been studied extensively.

Pedoturbation is the physical and/or biological mixing of soil material which tends to homogenize soil color and reduce soil morphology differences in the soil solum.

Pedoturbation is often a sub-surface or scarcely-noticeable soil surface activity, yet many studies dealing with soil mixing are based on surface observations of displaced soil without considering soil profile changes below the surface in determining the volume of soil disturbed by mixing.

Black-tailed prairie dogs (Cynomys ludovicianus) are burrowing rodents that alter soil development on gently-sloping, shortgrass areas of the Great Plains. The construction of "mounds" of soil is characteristic for this species, the result of burrowing into soil surfaces, subsoil tunneling, and concurrent soil excavation. The amount of excavated subsoil redeposited on the original soil surface increases with time as burrowing activity alters subsoil development and as new burrows are constructed in the same tunnel system.

Soils which are part of or underneath mound surface soils are chemically and physically different compared to soil derived from the same parent material in the absence of pedoturbation. The mound soil is generally lighter in

color, coarser-textured, higher in pH, lower in organic matter, variable in nitrogen content, and marked as a site of phosphorus accumulation by the addition of fecal, carcass, and plant residue wastes over time. Phosphorus accumulation in mound soil due to its relative chemical immobility would depend on the length of time a mound has been used by prairie dogs.

The objectives of this study were:

1. To examine soil development within selected prairie dog mounds compared to adjacent, non-mixed soil profiles.
2. To chemically analyze mound and adjacent non-mound soils in determining laboratory analyses which most consistently measure the length of time a mound has been used.
3. To estimate relative ages of studied mounds from mound surface area, underlying subsoil mixing, surface soil color and pH distribution, and from accumulation of total phosphorus in mound soils.

REVIEW OF THE LITERATURE

Geography and Climate

The Black Hills of west-central South Dakota and northeastern Wyoming encompass an irregularly-shaped area of about 5,650 square kilometers within 43-44 degrees North latitude and 103-104 degrees North longitude (Darton and Paige, 1925). The Hills rise up from surrounding grasslands to a maximum height of 2,207 meters at Harney Peak, with ridges and valleys vegetated largely by ponderosa pine (Pinus ponderosa).

The Black Hills are located in the semiarid Great Plains, but have a semiarid to subhumid continental climate. Precipitation ranges from high to low moving from the central part of the Hills towards the adjacent grasslands, with an average of 75 cm per year at Deadwood, SD. to 35-40 cm annually in the plains (Johnson, 1949). A majority of the precipitation falls in intermittent rains during the late spring and early summer months.

Temperatures typically fluctuate greatly from over 37 degrees C in summer down to near -40 degrees C in winter (Feldman and Heimlich, 1980). Thermal inversions during winter often moderate temperatures. Physical weathering processes have contributed to the development of sharp, angular landforms found in this climate.

Vegetation

The dense forest vegetation of the Hills gives them a "black" appearance when viewed from a distance (Newton and Jenny, 1880). Ponderosa pine forest makes up about 95 percent of forested area, with white spruce (Picea glauca) comprising about 4 percent of the coniferous timber. Spruce is the dominant tree on relatively moist sites (north aspects, higher elevations, and stream bottoms), especially in the northern reaches of the Limestone Plateau.

Isolated stands of Rocky Mountain juniper (Juniperus scopulorum), lodgepole pine (Pinus contorta var. latifolia), and limber pine (Pinus flexilis) occur as codominant or subordinate conifers (Thilenius, 1971; 1972). Important deciduous tree species include quaking aspen (Populus tremloides), often found on old burn sites either alone or in combination with paper birch (Betula papyrifera) or as a subordinate to ponderosa pine, and bur oak (Quercus macrocarpa), found along the northeastern foothills and on relatively dry slopes at moderate elevation in the northern Hills (Thilenius, 1971).

Numerous ridges occur where the outer edge of the Limestone Plateau merges with the Red Valley (Gartner and Thompson, 1973). The predominance of warm-season grasses there was termed a "mixed prairie...bunchgrass subclimax" by McIntosh (1949). Based on the degree of site

disturbance, a wide variety of grasses were found, including true prairie grasses and subclimax bunchgrasses. The former are represented by junegrass (Koeleria cristata), needlegrasses (Stipa comata, Stipa spartea, Stipa viridula), wheatgrasses (Agropyron smithii, Agropyron cristatum), gramagrasses (Bouteloua gracilis, Bouteloua curtipendula, Bouteloua hirsuta) and buffalo-grass (Buchloe dactyloides), while the latter classification was composed of little bluestem (Schizachyrium scoparium), big bluestem (Andropogon gerardii), indian-grass (Sorghastrum nutans), prairie dropseed (Sporobolus heterolepis), stoneyhills muhly (Muhlenbergia cuspidata), sedges (Carex spp.), and bluegrasses (Poa spp.).

McIntosh further noted that:

"The bunchgrass association forms a zone intermediate between the mixed prairie forest..."

and in the absence of natural disturbance (fire) that:

"...pines (Ponderosa pine) are invading this grassland and probably represent the climax vegetation of the area, but here and there are openings acres in extent covered by tall grasses."

Important forbs include white sage (Artemisia ludoviciana), silvery wormwood (Artemisia frigida), scurfpeas (Psoralea spp.), goldenrod (Solidago spp.), scarlet globe-mallow (Sphaeralea coccinea), and prairie dogweed (Dyssodia papposa), with other forbs and shrubs of local importance.

Bison (Bison bison), pronghorn antelope

(Antilocapra americana), mule deer (Odocoileus hemionus), whitetail deer (Odocoileus virginianus), and elk (Cervus canadensis) graze freely over large areas of Wind Cave National Park, the former two species commonly associated with prairie dog towns, although Lovaas (1972) concluded that antelope avoided heavily-grazed dog towns because the abundant forbs found there were a minor part of the antelope diet.

Geology

Primary references describing details of Black Hills geology are few in number (Feldman and Heimlich, 1980). Ludlow (1875) and Newton and Jenny (1880) compiled the earliest comprehensive geological records, but the work of Darton and Paige (1925) is generally accepted as being the most descriptive Hills geology reference (Feldman and Heimlich, 1980).

The interior Black Hills are a dome-shaped arrangement of exposed Pre-Cambrian age schists and slates with intrusive masses of granite (Feldman and Heimlich, 1980), referred to as the central crystalline core, dissected by scattered ridges and valleys. The core is encompassed by large limestone (Limestone Plateau) and sandstone outcrops which extend in wide bands on the western side of the Hills and are very narrow to the east. Encircling the Limestone Plateau near the margin of the

Hills is a relatively level basin of more readily-weathered, finer-textured reddish shale known as the Red Valley or "Racetrack" (Darton and Paige, 1925; Gartner and Thompson, 1973; Feldman and Heimlich, 1980). The perimeter of the Hills is marked by a long, narrow hogback ridge (Dakota Hogback) of Mesozoic-age shales and sandstones from 1,200 to 1,450 meters in elevation (Darton and Paige, 1925).

History of Fire and Extent of Forest in the Black Hills

The importance of wildfires caused by lightning in the Hills was referred to in the descriptions of early explorers in the area (Ludlow, 1875; Dodge, 1876; Newton and Jenny, 1880). Dodge (1876) related the occurrence of lightning-caused fires to the distribution of forest through his observations:

"The number of trees which bear the marks of the thunderbolt is very remarkable, and the strongest proof of the violence and frequent recurrence of these storms...forest openings, larger or smaller, sometimes several miles in extent and very irregular in shape...'parks'...caused by fires. A forest is destroyed (by fire). In a few years another fire destroys the young growth...this happens several times at intervals of a few years, effectually destroys both roots and seeds, and converts pine forests into parks."

Newton and Jenny (1880) also noted the interaction between fire and forest:

"Often...trees showing unmistakable marks of lightning would be seen...some portions of the parks and valleys, now destitute of trees, show by the presence of charred and decaying stumps that they were once covered by forest, but generally the pine springs up again as soon as it is burnt off,

though sometimes it is succeeded for a time by thickets of small aspens."

McIntosh (1927) recognized that "within the last 50 years man, fire, and the Black Hills (pine) beetle have been the most potent agents in altering the Black Hills flora..."

Black-tailed prairie dogs are among the mammals of the Hills region that are attracted to park-like areas of shortgrass vegetation which occur following fire or overgrazing (Schripsema, 1977; Garrett, 1982). The regrowth of desirable forbs and the elimination of tall grasses, shrubs, and trees for better visibility in locating predators aids prairie dogs in colonization of areas adjacent to existing dog towns. Repeated burning of an area will thin or eliminate woody plant species, thus maintaining grassland dominance in environments where woody vegetation would encroach into grass in the absence of fire (Humphrey, 1958). Fire also commonly favors forb growth (Biswell, 1973), which constitutes a sizeable share of the prairie dog diet (Kelso, 1939; Koford, 1958; Uresk, 1984).

Realistically, most modern literature dealing with forest-grassland interactions concerns expansion of forest into grassland due to the emphasis placed on fire control in forested public-use areas of the western United States (Biswell, 1973; Gartner and Thompson, 1973). Comparison

of photos from early Black Hills expeditions with those taken more recently allow for the study of vegetation changes (Progulske, 1972; Gruell, 1973), as do topographic maps of study areas (United States Geological Survey, 1957; 1979). Gartner and Thompson (1973) recorded ponderosa pine encroachment in the Red Valley on uplands where little bluestem was dominant, but not at sites where western wheatgrass proliferated. Differences in root growth habits and soil texture between the Red Valley and adjacent uplands were thought to be major factors in grassland-forest zonation there. White et al. (1969) examined soils in prairie and forested areas of the Black Hills and concluded that most areas of the Hills, regardless of present vegetation, were under forest at one time, and that soils were not the only factor limiting pine establishment in grasslands. Dense "doghair" stands of low-quality pine crowd out grass species, causing a reduction in quality and quantity of grassland production in areas of non-managed timber (Thompson and Gartner, 1971). The most active ponderosa pine encroachment in the Hills is in the transition area from the Limestone Plateau to the Red Valley and on the hogback ridge, with one landowner estimating that pine encroachment had extended about 2.5 kilometers from the outer slope to the outside of the hogback in 58 years (Gartner and Thompson, 1973).

Phosphorus Accumulation in Soil

Analysis of soils subject to non-mechanical pedoturbation has largely been undertaken as part of discovery of or investigation of abandoned human settlements. Chemical additions to soil by natural activities include nitrogen, phosphorus, potassium, and lesser amounts of calcium, magnesium, and sulfur. Nutrients are accumulated from excreta, burials, and other forms of refuse. Soil enrichment by such activity can increase phosphorus and nitrogen levels ten percent relative to natural soil concentrations (Eidt, 1984).

Phosphate analysis may be ideal in determining the location and estimated age of disturbed soil, provided the phosphate accumulates uniformly over the period of time the mound is inhabited and is fixed by the predominant soil cations at the site of deposition. Phosphorus is a primary part of plant and animal matter and is more available as phosphate in near-neutral pH soils (Waggamann, 1969; Eidt, 1977).

Prairie dog pedoturbation activity could be classified as propedisotropic, (Hole, 1961) or soil mixing that impedes soil formation by homogenizing two or more horizons of a soil profile. Body excreta, food caches, and dead plant and animal life enhanced soil phosphorus contents in Baxter and Hole's (1967) study of ant and

earthworm pedoturbation. Further differences between constructed mounds and surrounding soil have been found for many soil chemical and physical properties (Thorp, 1949; Buntley and Papendick, 1960; Hole, 1981 with an extensive bibliography), yet most observations were made concerning microfaunal activity (especially ants and earthworms). It is desirable to analyze soil samples in layers to estimate relative duration of soil phosphorus at a disturbed site, and also at intervals horizontally on a landscape to mark areal extent of pedoturbation and phosphorus accumulation (Eidt, 1984).

Soil phosphate levels at or near mound sites can be used as indicators of the age of mound materials. Puecker (1968) documented the work of J. G. Kohl, emphasizing Kohl's concentric ring theory of settlement analysis. The closer a sampling site is to the midpoint of a settlement, the greater is the density of built units and the degree of disturbance, and the older the structure built there. An older mound site would likely have a phosphorus content higher than adjacent non-mixed soil and higher than more recently-constructed mounds due to the cumulative effect phosphorus would have over time on the mounded soil. The interpretation of soil phosphate chemistry has emerged as a new tool in detection and analysis of abandoned settlements (Eidt, 1984), and has proven useful in studied comparisons between disturbed and non-disturbed soil profiles.

The highest animal percentages of phosphorus are found in bone. Animal wastes are likewise high in phosphates (Waggamann, 1969). For soil parent materials, igneous rocks contribute the greatest amount of total phosphate, about 1,000 mg Kg⁻¹ elemental phosphorus as an average (McKelvey, 1973). Sedimentary rocks range from 0 to 1,500 mg Kg⁻¹ elemental phosphorus in some sandstones and limestones, with metamorphic rocks containing similar amounts of phosphorus as their unaltered parent materials (Gotterman, 1973). Sources of phosphate addition present the most reliable data for identifying intensity of site use (Eidt, 1984). Although vegetation removal from prairie dog mound areas during habitation would decrease phosphate addition to soil from plant sources, large phosphate inputs to mound areas in food caches, fecal-urine, and carcass-bone additions could mark mounds over time as sites of phosphate accumulation.

Prairie Dogs

Black-tailed prairie dogs are herbivorous rodents of shortgrass prairie in the central Great Plains of North America (Costello, 1970). They occur as social units in colonies or "dog towns", using excavated subsoil and topsoil mixed together to form conical, asymmetrically-shaped mounds. Mounds of mixed subsoil and topsoil are referred to as crater mounds, while the larger dome mounds are

composed entirely of subsoil (Tileston and Lechleitner, 1966). Towns are typically located on gentle slopes of short grass interspersed with patches of forbs (Koford, 1958). Prairie dogs enhance forb growth by disturbing the soil at and near mound sites and by removing tall plants which hinder vision and movement, also creating favorable conditions for other prairie dog forage species including bluegrama and buffalograss (Summers and Linder, 1978; Uresk, 1984). Homeranges of prairie dogs are variable and depend on food resources, topography and climate within an area (Crocker-Bedford and Spillett, 1977).

Numerous attempts have been documented of prairie dog eradication with poisoning techniques (Taylor and Loftfield, 1924; Koford, 1958; Clark 1979, and others). Prairie dogs prefer overgrazed, depleted livestock ranges but are more an effect of such conditions than a cause (Koford, 1958). Prairie dogs readily abandon and reinhabit previously-used dog towns with environmental and population changes (Osborn, 1942; Osborn and Allen, 1949; R. Klukas, Wind Cave Biologist, personal communication, 1984), adding subsoil to existing mounds from further tunnel excavation (Longhurst, 1944).

Prairie dogs construct mounds with a frequency dependent on ease of subsoil excavation, degree of territory available for colony expansion, population

trends in a colony, and availability of food plant species in the colonized area. White and Carlson (1984) surveyed literature on prairie dog pedoturbation and found from 25 to 125 mounds constructed per hectare by prairie dogs within established towns, with 62 mounds representing an average number. Mounds created in optimum habitat are used annually by prairie dogs in the absence of predatory and environmental disturbance (Koford, 1958). Over time, addition of fecal, carcass, and plant residue materials chemically enrich mounded soil, particularly for phosphorus because of its immobility in soil. Prairie dogs limit daily activity to areas of grass and forbs which can be seen through or over; Often little bluestem and other tall perennial grasses mark edges of dog towns, serving as an effective barrier to colony expansion if stands are thick and tall enough (Koford, 1958). Shorter grasses and crops including western wheatgrass, bluegrama, annual grasses, brome-grasses (Bromus spp.), and cereal crops are less effective barriers, can be cut down rapidly, and are used as food by prairie dogs (King, 1955). Natural barriers may limit colony expansion, creating a situation where prairie dogs alter development of mound-area soils with time.

DESCRIPTION OF THE STUDY AREAS

Five separate areas of prairie dog mounds were selected for study in the park (Figure A). A continuum of degrees of mound construction, selected to represent differences in mound age, was included in sampled mounds.

Area 1 included mounds 1 and 2 and was located at the north border of the park in a creek valley, with prairie enclosed by ponderosa pine on adjacent slopes (see Results and Discussion for taxonomic and legal descriptions of all mounds). Soils were Bullflat-Cordeston silt loams: deep, well-drained soils derived from eroded silty alluvial-colluvial sediments of arkosic sandstone and limestone on 2 to 9 percent slopes. The area had 43 cm average annual precipitation and was at 1,330 meters elevation.

Mounds 3 through 5 were sampled from area 2, located on a nearly level terrace of an intermittent stream (Flynn creek) that flowed in a section of the Red Valley at the edge of the Hills. Vegetation was prairie, with a hogback ridge covered by ponderosa pine 1 kilometer to the east and scattered plains cottonwood (Populus deltoides var. occidentalis) and balsam poplar (Populus balsamifera) on the terrace-stream boundary. Annual precipitation averaged 38 cm. The deep, well-drained terrace soil was Tilford silt loam, formed in alluvium of reddish siltstone and shale on 0 to 5 percent slopes. The

Figure A. Location of Mound Sites in Wind Cave National Park.

soil had 25 to 50 cm of fluvial sediments overlying a boulder layer, with cobbles occupying interstices between boulders. The area was underlain by reddish Spearfish shale at depths of 1 to 2 meters. A tepee ring was found about 500 meters south of mound 3 and north of a gully created by the stream which deposited the terrace sediment. However, no evidence was discovered that area 2 mounds were constructed by human activity.

Area 3 was a large, active upland colony known as Bison Flats dog town. Mounds 6 through 12 chosen for analysis came from three zones of colonization within the town as determined by park personnel descriptions of colony expansion. Mounds 6 and 7 were located at the town center. These mounds were representative of the oldest mounds at the Bison Flats site. The colony expanded into non-colonized areas away from the town center after population increase and overgrazing of forage by prairie dogs in the town center made colony expansion necessary. Sparse vegetative cover between mounds consisted mainly of silvery wormwood and red threeawn (Aristida longiseta) with isolated thin stands of western wheatgrass. Tilford silt loam (0 to 3 percent slopes) was the dominant soil series within the inner colonization zone.

Mounds 8 and 9 were constructed southwest of the central colony at a depressional area of an old streambed

. Both mounds were inhabited at the time of sampling. Recent subsoil excavation, undecomposed scats, and a noticeable absence of vegetation around mounds indicated directional and areal extent of colony expansion from the town center. Prairie dog colonization of the streambed and surrounding landscape represented post-1976 colony expansion from the town center. Mound 8 soils were Hilger cobbly loam: deep, well-drained soils formed in alluvium on 0 to 40 percent slopes. Mound 9 (100 meters NNW of mound 8) was Winetti cobbly loam, a deep, very well-drained soil formed in alluvium from sedimentary rock on 2 to 10 percent slopes.

Mounds 10, 11, and 12 were part of the most recent colony expansion in Bison Flats (post-1982). Mounds in this zone were small in diameter, nearly circular, and composed of shallowly-excavated subsoil and topsoil. Mound 10 was located 600 meters WSW of mounds 8 and 9 and 50 meters east of U. S. highway 385 at the crest of a small ridge. Soil was Hilger cobbly loam, with excavated burrow material more reddish in color (2.5YR) than original surface soil. Mounds 11 and 12 were found on the footslope position of a ridge where terminated by the intersection of two drainage channels about 500 meters east of mound 10. The mounds were constructed 25 meters apart. Noticeable color differences between mounds indicated the nature of tilted beds of soil parent materials typical of the Black Hills. Soils there were part of the Hilger-

variant complex on 10 to 40 percent slopes. A majority of mounds observed nearby were in use, although mounds were generally located on level to gently-sloping surfaces and not on steep slopes. Sparse vegetation across the area was cropped close to the soil surface.

The fourth area chosen for study was part of an old, abandoned dog town on a high terrace between areas 1 and 2. Mounds 13 and 14 were constructed in Bullflat silt loam on level

sites 100 meters apart. Both mounds were large and overgrown by Kentucky bluegrass (Poa pratensis).

Scattered stands of little bluestem were present on mounds, while western wheatgrass grew at the edges of and adjacent to each mound. Non-mound areas were predominantly western wheatgrass but also were composed of green needlegrass and needle and thread. Few distinct mounds were identified in the area. Large, nearly circular mounds were predominant.

Two mounds were also sampled from a fifth area of the park, located west of the Rankin Ridge lookout and 400 meters west of state highway 87 near the northwest park border. Mounds 15 and 16 were part of post-1980 colony expansion and were constructed 60 meters apart on a strip of gently-sloping Hilger cobbly loam bordered by a dense stand of little bluestem to the south and a rocky ridge covered by ponderosa pine to the north. Close proximity of little bluestem to mound sites and small mound sizes across

the area confirmed recent mound construction there.

METHODS AND MATERIALS

Sampling and Laboratory Procedures (1984)

Mounds 1 and 2 from area 1 were trenched with tile spades and representative soil samples were collected from morphologically-distinct layers. A transect string was stretched across each mound 1 meter above the mound edges parallel to the trench as a reference point to identify positions of soil layers within the mound area. Vertical spade slices were placed horizontally side-by-side on polyethylene construction film to determine the layer sequence of soils as they existed in the mounded soil. From these measurements, cross-sectional diagrams of mound and adjacent non-mound soil profiles were drawn on graph paper.

Mound 3 in area 2 was sampled from holes dug with a spade and auger at 2 meter intervals along the mound transect. Trenching was not practical because of the size of the mound. Adjacent non-mixed soils were sampled with spade and auger 2 meters beyond both ends of all mound transects to represent soil development in the absence of prairie dog pedoturbation. These soils were used to evaluate soil changes in mounds. Samples were placed on polyethylene film and layers were identified. Soil structure and carbonate presence with hydrochloric acid were determined and recorded in the field. One sample

was collected from the original soil surface below the edge of the mounded soil and submitted to Teledyne Isotopes for C14-dating after removal of roots and other organic debris.

Samples of each soil layer were collected for chemical analysis and Munsell soil color determination (Soil Survey Staff, 1975). Dry and moist soil colors were read under uniform light intensity in the laboratory. Soil samples for laboratory analysis were crushed, screened (>2 millimeters), and oven-dried to constant weight before transfer to sample bags. Soil pH (saturated paste) and electrical conductivity were measured (Soil Conservation Service, 1972) using an Orion microprocessor ionalyzer and a Hach conductivity meter, respectively. Walkley-Black chromic acid and macro-Kjeldahl procedures were used, respectively, for organic matter and total nitrogen analyses.

Sampling and Laboratory Procedures (1985)

Mounds 4 through 16 were sampled systematically from the 0 to 10 cm surface layer (Figure 1). Mounds 1 through 3 studied in 1984 were resampled in 1985 because sampling procedure differed. Mounds were divided into zones of concentric circles which originated from the mound center and extended outward (radii of 0.5, 1, 2, 3 meters...) until the entire mound surface was encompassed

Figure 1. Subsample locations within mound sampling rings.

by sampling rings. Soil subsamples were collected midway between radii (0.25, 0.75, 1.5, 2.5 meters...). Numbers of subsamples collected within each ring increased with distance from the mound center to correspond with the increase in mound area (2, 3, 4, 5 subsamples...). Arc degrees between subsamples were 180, 120, 90, 72 degrees... for uniform spacing. Subsamples within each ring were mixed together to obtain a representative composite sample. A duplicate set of subsamples was composited from each ring at points midway between locations sampled for the first composite. The number of subsamples required was estimated by the variance found from previous determinations and methods outlined by Steel and Torrie (1980). Two sets of samples were taken 3 meters beyond each mound edge at 45 degree angles to each other (0, 90, 180, 270; 45, 135, 225, 315 degrees) to represent undisturbed soils. Arc positions for sample collection points were located to scale and marked on the polar coordinate paper. A string attached to a pin in the origin of the coordinate paper was rotated to the proper arc position and each subsample was collected at the proper distance from the mound center.

The boundaries beneath mounds where the original soil was still present and where soil mixing had occurred were located with a 1.9 cm-diameter soil auger. Total mound surface area and area of the mound where the original soil had been mixed by prairie dogs were plotted on polar

coordinate graph paper and each area was measured with a planimeter.

Collected samples were placed in paper bags for drying. After air-drying, soil was crushed, screened (>2 millimeters) and stored in plastic bags until analyzed for total phosphorus and for color. Carbonate effervescence with hydrochloric acid was determined in conjunction with color analysis. Soil pH (1:1 with deionized water) was also measured (McLean, 1982). Soil samples were hand-textured to identify texture differences between sampling rings. Previous studies (Mausbach et al., 1980) have shown a precision of about 3 to 5 percent in field texturing for clay content. The higher standard deviations for alluvial and residual soils reflect greater variations within parent materials rather than greater field texturing variability. A surface soil sample was also taken from the mound 1 surface for C14-dating so a chronological comparison could be made with the 1984 sample taken from the buried original soil surface.

Statistical Methods

Standard analysis of variance using a two-way design (sampling radii as rows and duplicate samples from each ring as columns) was used with the linear additive model as a basis for an F-test of two or more mean values for average total phosphorus contents:

- a). within each mound sampled
- b). between selected mounds

The linear additive model:

$$X_{ij} = u + T_i + B_j + E_{ij}$$

explained the variation between two populations (average phosphorus values within sampling rings of mounds and between different mounds in this case) as being composed of a mean (u), a component for the different sampling rings within and between each mound (T_i), a component for sampling error during collection of duplicate soil samples within each sampling ring (B_j), and a random element of variation (E_{ij}). In terms of the analysis of variance, the linear additive model represented:

$$\text{Total SS} = \text{Mean SS} + \text{Row SS} + \text{Column SS} + \text{Error SS}$$

where SS equals the sum of squares for each source of variation.

An HP-97 programmable calculator was used to input data. Calculated output included means of all rows and columns, complete analysis of variance including F-tests,

components of variance, and the coefficient of variation. Mean total phosphorus values for sampling rings and for mounds were compared to control means and significance at 90 and 95 percent confidence levels was determined by an F-ratio test (Steel and Torrie, 1960; 1980). A set of orthogonal comparisons was used to test for significance between average inner mound and non-mound total phosphorus contents across selected mounds.

Soil pH samples were subject to a two-way analysis of variance with subsampling, comparing selected mounds from the same study area. Two-way analysis of variance was also used in testing for significance between mound and undisturbed soils at the same mound site. Procedures, tests, and confidence levels established were similar to those used for mean total phosphorus except that the inner two sampling rings were compared to the non-mound sampling ring to maintain a uniform number of subsamples for analysis of mounds within the same study area.

Mound Area Relationships

After calculating total mound surface area and disturbed soil area beneath mounds, total mound surface area measurements obtained by planimeter readings from mound diagrams were presented as a logarithmic function of disturbed soil area to establish the correlation

between external and internal mound dimensions. Measurement of total mound surface area assumed mounds were of conical shape. Soil mixing boundaries located by auger underneath mounds best represented pedoturbation below the original soil surface.

Age estimates of mounds 1 and 3, based on C14 analyses, were used along with calculated area of disturbed soil beneath studied mounds to develop a linear scale of relative mound ages. Mound age was plotted as a function of disturbed original surface area to obtain a graphic chronology of mound construction.

Phosphorus

Soil samples were analyzed for total phosphorus utilizing a nitric-perchloric acid digestion (Jackson, 1958) and Murphy-Riley ascorbic acid procedure (Watanabe and Olsen, 1975). Half gram soil samples were digested with 10 milliliters of 50 percent nitric acid in 125 milliliter Erlenmeyer flasks. After the dry residue was moistened with distilled water, 10 milliliters of 70 percent perchloric acid were added and samples were digested to release phosphorus from organic and inorganic compounds in the residue. An aspirator was used to draw off perchloric acid fumes during the digestion. After samples were nearly dry, distilled water was added to each flask and the contents were filtered through 11 cm Whatman 50 filter paper and collected in 100 milliliter volumetric flasks. Each Erlenmeyer flask was washed three times to extract all phosphorus from the filtrate. After dilution to 100 milliliters and thorough shaking, a 5 milliliter aliquot of each solution was transferred from the 100 milliliter volumetric flask to a separate 50 milliliter volumetric flask.

Twenty milliliters of distilled water and 2 drops of P-nitrophenol indicator were added to each sample, with 5N sodium hydroxide then added dropwise until the solution turned yellow. An 8 milliliter volume of the following reagent mixture was added to each sample with an automatic

pipette. The resultant solution was diluted to 50 milliliters and shaken:

Reagents (single batch--about 30 samples)

1. 5N sulfuric acid--70 milliliters diluted to 500 milliliters.
2. 20 grams ammonium molybdate diluted to 500 milliliters.
3. 0.2743 grams potassium antimony tartrate in 100 milliliters distilled water.
4. 1.32 grams ascorbic acid in 75 milliliters distilled water.

Samples were analyzed colorimetrically with a Turner Model 330 spectrophotometer. A set of phosphorus color standards were prepared to represent 0, 400, 800, 1,200, 1,600, and 2,000 mg Kg⁻¹ phosphorus solutions. The percent transmittance was plotted against total phosphorus on semi-logarithmic paper for a standard curve. the percent transmittance determined for solutions made from each soil sample was used to determine total phosphorus content from the standard curve. The phosphorus content in the original sample was calculated with:

$$\text{mg Kg}^{-1} \text{ P in soil} = \frac{(\text{ml in vol. flask} \times \text{mg Kg}^{-1} \text{ from curve})}{\text{grams of soil used}}$$

$$\times 100/5 \text{ (a dilution factor)}$$

If all steps of the laboratory procedure were followed as outlined, the above equation could be simplified to:

$$\text{mg Kg}^{-1} \text{ P in soil} = \text{mg Kg}^{-1} \text{ P from curve} \times 2,000$$

RESULTS AND DISCUSSION

Soil Changes Within Mounds (1984)

Soil properties were studied vertically within each mound along a horizontal transect across the mound surface in 1984. Soil differences within mounds and between mounds and undisturbed adjacent soil are useful in estimating the duration and magnitude of soil alteration (Eidt, 1984).

Data collected for mounds 1 through 3 is presented in Tables 1 through 8 and Figures 2 through 13. Conductivity and organic matter data were left in tabular form due to variability of each property, although g Kg-1 organic matter analyses tended to parallel g Kg-1 nitrogen levels.

A. Soil Color

Soil colors of mound and non-mound soils were markedly different with depth. Mound 1 and 2 soils had lower values and chromas (ie., were darker and grayer) than the subsoil of adjacent non-mound soils (Figures 2 and 3). The cross-sectional diagram of mound 1 illustrates mound dimensions but not slope direction, which is from right to left on all mound 1, 2, and 3 figures. Downslope movement of excavated subsoil would require less energy, thus mound enlargement over time has probably proceeded downslope from soil E to soil B.

Moist soil colors generally decrease in value and

Figure 2. Munsell color (10YR hue) value and chroma for dry (left side) and moist (right side) soil samples from profiles A through F in mound 1.

Figure 3. Munsell color (10YR hue) value and chroma for dry (left side) and moist (right side) soil samples from profiles G through J in mound 2.

chroma as soil organic matter contents increase. Soil nitrogen contents tend to increase as soil organic matter increases. Subsoil deposited in the original soil surface, pedoturbation, and settling of soil layers above excavated tunnels and galleries (tunnels where plant residues and excreta are accumulated) could homogenize large color variations and eventually develop intermediate color properties of the undisturbed adjacent soils. Tunnels beneath mounds are often below the normal depth of wetting and drying in southwestern South Dakota (Sheets et al., 1971; Westin and Malo, 1978) so that they tend to remain open. Mound 1, in particular, contains remnants of forest vegetation soil colors (10YR 4/1, 5/1 dry) which have settled downward compared to undisturbed soils. Black Hills soils developed under forest vegetation have lower phosphorus contents (White et al., 1974b) and lighter (more grayish) chromas relative to prairie-derived soils (White et al., 1969). Soil colors of layers in mounds tended to be more similar than in adjacent non-mound soils because pedoturbation counteracted soil horizon formation.

Mound 3 soil colors exhibited wide variability (Figure 4). Hues ranged from 2.5YR to 7.5YR, with reddish (2.5YR) hues located in the mound above a subsidence in the boulder layer at soils N to P. Stream-deposited boulders mark the boundary between reddish Spearfish-formation

Figure 4. Munsell colors (A=7.5YR, B=5YR, C=2.5YR hues) value and chroma for dry and moist soil profile samples K through R from mound 3.

silty clay and the overlying alluvial-derived Tilford soil. Variable chemical and physical properties of alluvium and residuum have been documented (Mausbach et al., 1980), which in combination with different degrees of soil mixing would produce a range of soil colors and properties in mound constructed from these parent materials. Prairie dogs apparently were able to tunnel through a crevice in the boulder layer along an upward network of tunnels above the boulder layer subsidence. This subsidence was caused by settling of boulders downward into underlying tunnels.

Mound 3 surface hues were 5YR in comparison to the 7.5YR hues of adjacent non-mound surface layers. Mound values and chromas were higher and hues more reddish than non-mound soils. Both color differences could be caused by pedoturbation associated with tunneling and excavation. For mound 3 construction, there was extensive subsoil movement upward onto the original soil surface. Differences in the amount of mixing of materials with 2.5YR and 7.5YR hues could form intermediate 5YR hues within the mound. Except at the ends of the mound transect (soils K and R), the original soil surface layer was not identifiable and presumably was mixed with other mound soil materials. Pedoturbation homogenizes soil colors in older mounds to more closely resemble mound subsoil hues since a greater volume of subsoil would be used over time as a mound is enlarged. Younger mounds would have a soil color

more characteristic of the original soil surface due to prairie dog scraping of the ground surface in mound construction (Koford, 1958) and lack of deep subsoil tunneling present in younger, smaller-sized mounds.

B. Nitrogen

Soil nitrogen content differences are partially caused by pedoturbation within mound soils (Figures 5, 6, and 7). Mounds 1 and 2 had decreasing nitrogen contents with depth in the mound soil and an increase in the surface layer of the buried soil. Prairie dog fecal materials, carcasses, and plant materials deposited in galleries (Sheets et al., 1971) and at the mound surface are mixed with excavated soil in different degrees, causing differences in nitrogen contents in mound soil layers.

Adjacent non-mound soils (A and F, G and J) had decreasing nitrogen contents with depth and upper solum melanization (darkening) characteristic of soils derived from prairie vegetation (Smith et al., 1950). Nitrogen contents are more uniform with depth in mound profiles than in adjacent undisturbed soils. Addition and decomposition of prairie dog excreta and pedoturbation of mound materials homogenized nitrogen levels within mounds. Downhill

Figure 5. Nitrogen content of soil samples from mound 1 ($10 \times \%N = \text{g/Kg N}$). Letters A through F represent soil profiles across the mound surface.

Figure 6. Nitrogen content of soil samples from mound 2 ($10 \times \%N = \text{g/Kg N}$). Letters G through J represent soil profiles across the mound surface.

Figure 7. Nitrogen contents of soil profiles K through R from mound 3 (10 X %N = g/Kg N).

movement of excreta and the location of a burrow entrance near soil D might explain high nitrogen levels found in soil A.

Nitrogen levels decreased with depth in both mound and undisturbed soils at mound 3. The more abrupt decrease in nitrogen contents with depth in soils N, O, and P may be because Spearfish shale has been added to the mound soil above the boulder layer. Pedoturbation has mixed the original soil surface beneath mound 3 so that nitrogen contents there are similar to overlying and underlying soil layers.

C. Soil pH

Mound soil pH values increased irregularly with depth and were higher relative to upper layers of non-mound soils at all three studied mounds (Figures 8, 9, and 10). Prairie dog tunneling activity may extend downward to considerable depth (Koford, 1958; Sheets et al., 1971). Calcareous subsoil is moved upward and deposited on the original soil surface and on the mound. Leaching of calcium carbonate (CaCO_3) from this mound material caused recalcification of the buried soil. Low pH readings at mound 1 surface soils D and E are likely caused by downhill movement of excavated subsoil from this upslope portion of the mound. Soils D and E are probably older and have had deeper leaching of CaCO_3 .

Figure 8. Soil pH values of profiles A through F in mound 1 (saturated paste).

Figure 9. Soil pH values of profiles G through J in mound 2 (saturated paste).

Figure 10. Soil pH values of profiles K through R in mound 3 (saturated paste).

Secondary carbonates coated the surfaces of prismatic soil structural pedes in the buried original surface. Calcium carbonate-saturated water draining through calcareous mound soil moves between prism faces in the buried soil and flows horizontally into the prisms. The wetting of soil minerals removes water from the soil solution so that soil pedes become supersaturated with CaCO_3 . Dissolved CaCO_3 is then deposited on the surface of and within structural pedes as precipitated secondary carbonates. Secondary carbonates can accumulate rapidly on ped faces of buried archaeological soils less than 1,000 years old (Keller et al., 1984). At least 200 years would be required to coat ped faces in South Dakota (E. M. White, personal communication), dependent on percent CaCO_3 equivalent in mound soil, soil texture, precipitation, soil pH, and depth and frequency of soil wet-dry cycles.

D. Total Phosphorus

Soil total phosphorus contents are an indicator of mound age and pedoturbation. Differences in phosphorus distribution were apparent in mounds 1 and 2 (Figures 11 and 12). Phosphorus levels within and adjacent to mound 2 were generally higher than mound 1. Lower phosphorus contents in mound 1 may represent remnants of soil developed in a previously-forested environment (White et al., 1974b) at the site. Pedoturbation and enrichment of soil by excreta

Figure 11. Total phosphorus contents of soil profiles A through F from mound 1. (ppm P = mg Kg^{-1} P)

Figure 12. Total phosphorus contents of soil profiles G through J in mound 2.
(ppm P = mg Kg⁻¹ P)

may have masked characteristics of forested soil conditions at mound 2. Close proximity of mounds 1 and 2 (about 100 meters apart) supports the idea that more intensive pedoturbation has occurred at mound 2, opposed to non-forested conditions having always existed there.

Soils B, C, D, and E probably form a sequence from youngest to oldest across mound 1. Heterogeneity between layers and profiles is caused by subsoil pedoturbation with surface soil and subsidence of layers as supporting soil is tunneled, in addition to influence of past vegetation.

The larger volume of mixing in mound 2, compared to mound 1, suggests a longer period of forest absence and of prairie dog colonization at mound 2. Soil total phosphorus contents on the non-mound soil surface probably were increased by nutrient cycling and phosphorus deposition by prairie plants. High surface phosphorus contents are also due to excreta addition on and near the mound. Total phosphorus analyses conducted on prairie dog fecal pellets showed an average total phosphorus content of 1,924 mg Kg⁻¹, ranging from 1,724 to 2,384 mg Kg⁻¹ P. The high surface phosphorus content at soil G of mound 2 may be due to downslope movement of fecal pellets from the mound surface.

Combined effects of mixing of Spearfish subsoil with overlying Tilford silt loam and addition of fecal, carcass, bone, and plant residues resulted in higher total phosphorus levels in soil O and across the mound 3 surface

relative to non-mound soils there. Phosphorus cycling by vegetation may have influenced non-mound surface soils K and R (Figure 13). Total phosphorus was greater in the surface layer of the non-mound soil than in the subsurface soil immediately below the surface, but total phosphorus then increased with depth.

Soils M through O likely represent the mound center. Excavated subsoil deposited in tunnels and fecal material in galleries and at the mound surface could greatly affect phosphorus contents of these soil profiles. Increased prairie dog activity on and beneath the central mound area would increase phosphorus deposition there. In support of this idea, soil O has a much higher conductivity in subsurface soil layers than at any other location in mound 3. High conductivity there may be due to ions in prairie dog excreta.

Total phosphorus content of a soil varies little with soil maturity (Bohn, 1979). Differences in phosphorus contents within and between mound and non-mound soils reflect cumulative effects of pedoturbation, soil enrichment, nutrient cycling, and parent material variability above and below the boulder layer. Faunal pedoturbation resulted in phosphorus accumulation within disturbed soils in a number of studies (Baxter and Hole, 1967; Salem and Hole, 1968; Laycock and Richardson, 1975; Kalisz and

Figure 13. Total phosphorus contents of soil profiles K through R in mound 3.
(ppm P = mg Kg⁻¹ P)

Stone, 1983; Cowan et al., 1985, and others).

Mound phosphorus contents might initially increase or decrease at the site of a newly-constructed prairie dog mound, depending on the degree of plant removal, quantity of subsoil material used in mound construction relative to surface soil, and the subsoil phosphorus content. Phosphorus would likely accumulate over time as greater volumes of subsoil are disturbed and prairie dog fecal, bone, and plant residues are incorporated with mound soil. This description may represent what has taken place at the mound 3 site.

Table 1

Summary of Laboratory Analyses for Mound 1--ENE Trench (1984)

Profile	Depth (cm)	N ¹ (g/Kg)	pH	EC (dS/m)	OM ² (g/Kg)	Total P ³ (mg Kg ⁻¹)	Color	
							Dry	Moist
1-C	0-16	1.09	7.56	0.405	9.7	610	10YR 5/2	10YR 3/2
	16-26	0.68	7.71	0.520	6.7	630	10YR 5/2	10YR 3/2
	26-36	0.69	7.79	0.800	4.8	620	10YR 6/2	10YR 4/2
	36-45	1.07	7.83	0.425	10.8	390	10YR 4/1	10YR 3/2
	45-62	1.14	7.91	0.280	10.4	540	10YR 4/1	10YR 3/2
	62-70	0.92	7.94	0.270	7.1	480	10YR 5/2	10YR 3/2
	70-85	0.90	7.91	0.405	6.2	500	10YR 5/3	10YR 3/3
1-B	0-9	1.40	7.59	0.610	15.3	610	10YR 5/2	10YR 4/1
	9-21	0.84	7.56	0.600	7.6	700	10YR 6/2	10YR 4/2
	21-35	1.04	7.54	0.775	9.8	760	10YR 5/2	10YR 3/2
	35-45	1.22	7.89	0.435	12.7	420	10YR 5/2	10YR 4/2
	45-60	1.29	7.93	0.485	11.2	460	10YR 4/1	10YR 3/1
	60-70	1.12	7.83	0.415	9.3	460	10YR 5/2	10YR 3/2
	70-85	1.01	7.74	0.790	7.3	470	10YR 5/3	10YR 3/3
1-A (N-M) ⁴	0-16	3.02	5.96	0.375	16.2	610	10YR 4/1	10YR 2/2
	16-32	1.36	6.33	0.630	12.2	530	10YR 5/1	10YR 3/2
	32-50	0.74	6.59	0.410	5.5	420	10YR 5/2	10YR 3/3
	50-72	0.74	6.84	0.400	6.3	480	10YR 5/2	10YR 3/3
	72-90	0.72	6.77	0.650	5.1	530	10YR 5/3	10YR 3/3
	90-117	0.44	8.02	0.470	2.5	580	10YR 5/3	10YR 3/3

¹ g/Kg N = 10 X %N. ² g/Kg OM = 10 X %OM. ³ mg Kg⁻¹ = ppm.

⁴ (N-M) = Non-Mound soil.

Table 2

Summary of Laboratory Analyses for Mound 1--WSW Trench (1984)

<u>Profile</u>	<u>Depth (cm)</u>	<u>N¹ (g/Kg)</u>	<u>pH</u>	<u>EC (dS/m)</u>	<u>OM (g/Kg)</u>	<u>Total P (mg Kg⁻¹)</u>	<u>Color</u>	
							<u>Dry</u>	<u>Moist</u>
1-D	0-9	2.92	6.58	0.840	15.5	620	10YR 5/2	10YR 3/2
	9-22	1.01	6.98	0.795	9.1	680	10YR 5/2	10YR 4/2
	22-32	0.91	7.35	0.620	9.2	560	10YR 5/2	10YR 3/2
	32-50	0.83	7.84	0.560	7.8	530	10YR 4/1	10YR 3/2
	50-64	1.01	7.73	0.750	9.8	540	10YR 5/1	10YR 3/1
	64-73	1.05	7.89	0.325	9.4	520	10YR 5/1	10YR 3/1
	73-85	0.91	7.62	0.710	7.5	480	10YR 5/3	10YR 3/3
1-E	0-14	1.38	6.78	0.620	12.5	500	10YR 5/2	10YR 3/2
	14-26	0.86	7.63	0.850	7.0	630	10YR 4/1	10YR 3/2
	26-50	0.76	7.78	0.490	7.4	460	10YR 5/2	10YR 4/2
	50-65	0.87	7.97	0.380	7.4	580	10YR 5/2	10YR 3/2
	65-76	0.94	7.90	0.500	7.5	480	10YR 5/3	10YR 3/2
	76-85	0.94	7.94	0.730	7.0	500	10YR 5/3	10YR 3/3
1-F (N-M)	0-14	1.81	5.99	0.440	15.9	660	10YR 4/1	10YR 3/1
	14-41	1.29	6.38	0.375	10.8	670	10YR 5/1	10YR 3/2
	41-57	0.94	7.84	0.420	7.1	540	10YR 6/2	10YR 4/2
	57-75	0.83	7.81	0.680	5.1	550	10YR 6/3	10YR 4/2
	75-87	0.76	7.89	0.460	4.9	540	10YR 6/3	10YR 4/2
	87-117	0.56	7.90	0.680	2.1	720	10YR 6/3	10YR 5/3

Table 3

Summary of Laboratory Analyses for Mound 2--ENE Trench (1984)

Profile	Depth (cm)	N (g/Kg)	pH	EC (dS/m)	OM (g/Kg)	Total P (mg Kg ⁻¹)	Color	
							Dry	Moist
2-H	0-14	2.00	6.95	1.200	16.2	730	10YR 5/2	10YR 4/2
	14-22	1.50	7.14	1.250	11.5	710	10YR 5/2	10YR 3/2
	22-34	0.81	7.94	0.610	6.1	540	10YR 5/3	10YR 4/3
	34-56	1.36	7.82	0.460	10.0	620	10YR 5/2	10YR 4/2
	56-68	1.52	7.65	0.455	12.8	560	10YR 5/1	10YR 3/2
	68-84	1.02	7.20	0.595	7.4	610	10YR 5/3	10YR 4/2
	84-93	1.06	7.85	0.805	7.8	460	10YR 5/3	10YR 4/2
	93+	0.73	7.59	0.670	4.9	650	10YR 6/3	10YR 5/3
2-G (N-M)	0-5	6.62	5.34	0.600	22.6	900	10YR 5/2	10YR 3/2
	5-12	2.93	5.50	0.290	19.5	760	10YR 5/2	10YR 3/2
	12-29	1.17	6.26	1.560	9.1	750	10YR 5/3	10YR 3/2
	29-44	0.97	7.35	0.570	7.6	690	10YR 6/3	10YR 4/2
	44-75	0.66	7.64	0.560	5.3	620	10YR 6/3	10YR 4/3
	75+	0.66	7.96	0.580	5.2	700	10YR 7/2	10YR 5/3

Table 4

Summary of Laboratory Analyses for Mound 2--WSW Trench (1984)

<u>Profile</u>	<u>Depth (cm)</u>	<u>N (g/Kg)</u>	<u>pH</u>	<u>EC (dS/m)</u>	<u>OM (g/Kg)</u>	<u>Total P (mg Kg⁻¹)</u>	<u>Color</u>	
							<u>Dry</u>	<u>Moist</u>
2-I	0-14	1.69	6.98	0.960	16.8	740	10YR 5/2	10YR 3/2
	14-29	1.03	7.61	0.730	7.2	650	10YR 6/3	10YR 4/2
	29-40	1.08	7.78	0.530	8.4	680	10YR 4/2	10YR 3/2
	40-56	0.88	7.73	0.580	8.6	690	10YR 5/2	10YR 3/2
	56-70	1.02	7.71	0.530	7.6	690	10YR 4/3	10YR 3/3
	70-86	0.92	7.78	0.540	7.3	650	10YR 5/2	10YR 4/2
	86-97	0.63	7.90	0.620	4.5	560	10YR 6/3	10YR 5/3
	97+	0.91	7.87	0.580	6.5	690	10YR 6/3	10YR 5/3
2-J (N-M)	0-6	4.21	5.94	0.440	21.3	860	10YR 4/1	10YR 3/1
	6-14	2.03	5.95	0.290	18.8	760	10YR 4/1	10YR 3/1
	14-25	1.44	6.61	0.380	12.4	690	10YR 5/2	10YR 4/2
	25-42	0.95	7.03	0.510	6.7	270	10YR 5/3	10YR 4/2
	42-58	0.85	6.88	0.525	5.9	680	10YR 5/3	10YR 4/2
	58-69	0.81	6.98	0.500	5.9	690	10YR 5/3	10YR 4/2
	69-91	0.35	7.12	0.550	2.4	460	10YR 6/3	10YR 5/3

Table 5

Summary of Laboratory Analyses for Mound 3 (1984)

Profile	Depth (cm)	N (g/Kg)	pH	EC (dS/m)	OM (g/Kg)	Total P (mg Kg ⁻¹)	Color Dry	Moist
3-K (N-M)	0-8	2.72	6.54	0.540	25.3	580	7.5YR 4/2	5YR 3/2
	8-14	1.67	6.89	0.520	13.5	150	5YR 4/2	5YR 3/2
	14-30	1.41	7.23	0.800	10.0	650	5YR 4/3	5YR 4/2
	30-43	1.37	7.87	0.550	9.6	640	5YR 6/3	5YR 4/3
	43-50	1.35	7.78	0.500	9.0	710	5YR 5/3	5YR 4/3
<hr/>								
3-L	0-6	2.79	7.21	0.490	24.6	640	5YR 4/3	5YR 3/2
	6-23	1.76	7.61	0.450	13.1	600	5YR 4/3	5YR 3/3
	23-35	1.61	7.67	0.660	11.6	540	5YR 5/3	5YR 4/3
	35-49	1.22	7.90	0.480	9.3	540	5YR 6/2	5YR 5/3
	49-58	0.82	7.98	0.460	5.6	510	5YR 6/3	5YR 5/3
	58-66	0.90	7.90	0.530	5.6	500	7.5YR 6/4	7.5YR 5/4
	66-73	0.75	7.99	0.475	4.7	480	7.5YR 7/2	7.5YR 5/4
<hr/>								
3-M	0-6	2.18	7.50	0.470	19.4	710	5YR 5/6	2.5YR 3/4
	6-12	1.26	7.68	0.470	9.9	690	5YR 5/6	2.5YR 3/4
	12-22	1.26	7.56	0.480	8.8	690	5YR 5/4	2.5YR 3/4
	22-46	0.97	7.85	0.375	6.1	660	5YR 5/4	5YR 4/4
	46-57	0.96	7.92	0.440	6.2	620	5YR 5/3	5YR 4/3
	57-69	0.87	7.97	0.470	5.7	660	5YR 6/3	5YR 5/3
	69-82	0.67	8.03	0.480	4.8	580	5YR 6/3	7.5YR 5/4
	82-95	0.63	7.99	0.500	1.0	640	5YR 6/3	5YR 5/4

Table 6

Summary of Laboratory Analyses for Mound 3 (1984)

<u>Profile</u>	<u>Depth</u> <u>(cm)</u>	<u>N</u> <u>(g/Kg)</u>	<u>pH</u>	<u>EC</u> <u>(dS/m)</u>	<u>(g/Kg)</u>	<u>Total P</u> <u>(mg Kg⁻¹)</u>	<u>Color</u>	
							<u>Dry</u>	<u>Moist</u>
3-N	0-7	1.84	7.59	0.475	16.7	900	5YR 4/6	2.5YR 3/6
	7-16	1.13	7.77	0.430	8.9	800	5YR 4/6	2.5YR 3/6
	16-36	0.75	7.90	0.420	5.8	690	2.5YR 4/6	2.5YR 3/6
	36-56	0.74	8.01	0.440	5.5	560	2.5YR 4/6	2.5YR 3/6
	56-78	0.51	8.05	0.480	3.1	580	5YR 4/6	2.5YR 4/4
	78-94	0.55	8.20	0.570	3.1	540	5YR 5/4	2.5YR 4/6
	94-115	0.41	8.39	0.540	2.1	500	7.5YR 6/4	5YR 5/4
3-O	0-6	2.80	7.46	0.510	24.2	1530	5YR 5/4	2.5YR 3/6
	6-16	1.23	7.69	0.480	9.0	880	5YR 5/4	2.5YR 3/6
	16-38	1.36	7.73	0.470	9.7	710	5YR 5/4	2.5YR 3/6
	38-61	1.08	7.90	0.380	6.5	730	2.5YR 5/4	2.5YR 4/4
	61-82	0.63	7.93	0.510	3.9	710	2.5YR 5/4	2.5YR 3/6
	82-102	0.56	7.93	1.400	3.0	800	2.5YR 5/4	2.5YR 3/6
	102-113	0.65	7.88	1.600	2.8	880	2.5YR 5/4	2.5YR 3/6
	113-122	0.65	7.96	2.300	4.2	850	5YR 6/3	5YR 5/3
	122-155	0.61	7.65	1.820	3.1	760	2.5YR 5/4	2.5YR 4/4

Table 7

Summary of Laboratory Analyses for Mound 3 (1984)

<u>Profile</u>	<u>Depth</u> <u>(cm)</u>	<u>N</u> <u>(g/Kg)</u>	<u>pH</u>	<u>EC</u> <u>(dS/m)</u>	<u>OM</u> <u>(g/Kg)</u>	<u>Total P</u> <u>(mg Kg⁻¹)</u>	<u>Color</u> <u>Dry</u>	<u>Moist</u>
3-P	0-6	2.57	7.33	0.520	22.9	710	5YR 5/4	5YR 3/4
	6-23	1.84	7.70	0.430	12.4	580	5YR 5/4	5YR 3/4
	23-38	1.34	7.80	0.390	9.4	450	2.5YR 5/4	2.5YR 3/6
	38-55	0.61	7.88	0.390	4.3	500	2.5YR 5/4	2.5YR 3/6
	55-79	0.59	7.96	0.430	3.5	480	2.5YR 5/4	2.5YR 3/6
	79-95	0.39	8.03	0.480	2.1	580	5YR 6/4	5YR 4/4
	95-117	0.34	8.13	0.500	1.9	460	7.5YR 6/4	7.5YR 5/4
3-Q	0-8	2.36	7.40	0.440	19.8	600	5YR 4/4	5YR 3/4
	8-23	1.32	7.69	0.370	8.4	530	5YR 5/4	5YR 4/6
	23-32	0.82	7.78	0.440	5.6	500	5YR 5/4	5YR 4/6
	32-39	0.93	7.88	0.470	6.1	470	7.5YR 5/4	5YR 4/4
	39-47	0.82	7.75	0.500	5.9	460	7.5YR 6/4	7.5YR 4/4
3-R (N-M)	0-6	2.21	6.73	0.500	27.5	390	7.5YR 4/2	5YR 3/2
	6-15	1.77	7.02	0.430	14.5	370	5YR 4/2	5YR 3/2
	5-27	1.38	7.17	0.490	9.9	400	5YR 4/3	5YR 3/3
	27-35	1.36	7.29	0.630	11.4	470	7.5YR 6/4	5YR 4/4

Table 8

Ranges and Means of Soil Properties Measured for Mounds 1-3 (1984)

<u>Mound</u>		<u>g/Kg N</u>	<u>pH</u>	<u>dS/m EC</u>	<u>g/Kg OM</u>	<u>mg Kg⁻¹ P</u>
1	Range	0.44-3.02	5.96-8.02	0.270-0.850	2.1-16.2	390-760
	Mean	1.08	7.46	0.548	8.7	553
2	Range	0.35-6.62	5.34-7.96	0.290-1.560	2.4-22.6	270-900
	Mean	1.44	7.14	0.637	9.9	657
3	Range	0.34-2.80	6.54-8.39	0.370-2.300	1.0-27.5	150-1530
	Mean	1.22	7.70	0.584	9.3	618

SOIL CHANGES ACROSS MOUNDS (1985)

Taxonomic and legal descriptions of studied mounds were obtained from the Soil Conservation Service and are given in Tables 9 and 10. Mound areas were moderately to excessively well-drained soils formed in alluvium-colluvium or residuum and were not on steep slopes except in newer and marginal areas of colony expansion.

A. Soil Color

Distinct color changes were observed across most mound surfaces (Table 11). Hues remained constant or became darker (ie., became or intergraded to a 10YR hue) from the mound center outward to the non-mound soil. Soil values became darker and chromas more gray or blackish with distance from the mound center. Duplicates a and b taken from sampling rings usually had similar colors. Most color differences between duplicates were found in sampling rings near the mound center. Smaller numbers of subsamples collected for inner ring composite samples and excavation and deposition of different-colored soil layers around the central burrow would cause potential color variability between duplicates

Surface soil color distribution from mound center to adjacent non-mound soil may reflect mound age and extent of pedoturbation. Younger-aged mounds (mounds 1, 8, 10, 11, 15, and 16) and older mounds abandoned for a long

Table 9

Classification of Soils Sampled for Prairie Dog Mound Analysis¹

<u>Soil Series</u>	<u>Classification</u>	
Bullflat	Typic Argiborolls	Fine-loamy, mixed
Cordeston	Cumulic Haploborolls	Fine-loamy, mixed
Hilger	Typic Argiborolls	Loamy-skeletal, mixed
Tilford	Torriorthentic Haplustolls	Fine-silty, mixed, mesic
Winetti	Typic Ustifluvents	Loamy-skeletal, mixed, (calcareous), frigid

¹ From "Classification of South Dakota Soils 1986". South Dakota State University Agricultural Experiment Station Technical Bulletin Number 58. Gary Lemme, Plant Science Department, South Dakota State University, and USDA-SCS Soil Survey Staff.

Table 10

Studied Soils and Mound Locations

<u>Mound</u>	<u>Soil Mapping Unit</u>	<u>Legal Description¹</u>
1	Bullflat-Cordeston (silt) loams, 0-10% slopes.	500'N, 650'W of NE corner, Section 7, T5S, R6E.
2	Bullflat-Cordeston (silt) loams, 0-10% slopes.	300'N, 300'W of NE corner, Section 7, T5S, R6E.
3 ²	Tilford silt loam, 0-15% slopes	1775'S, 250'W of NE corner, Section 22, T5S, R6E.
4 ²	Tilford silt loam, 0-15% slopes	2250'S, 250'W of NE corner Section 22, T5S, R6E.
5 ²	Tilford silt loam, 0-15% slopes	2550'S, 250'W of NE corner, Section 22, T5S, R6E.
6	Tilford silt loam, 0-15% slopes	2550'S, 1450'E of NW corner, Section 12, T6S, R5E.
7	Tilford silt loam, 0-15% slopes	2500'N, 1230'E of SW corner, Section 12, T6S, R5E.
8 ³	Hilger cobbly loam, 0-40% slopes	1025'N, 630'W of SE corner, Section 11, T6S, R5E.
9	Winetti cobbly loam, 2-10% slopes	1600'N, 800'W of SE corner, Section 11, T6S, R5E.

Table 10 (continued)

<u>Mound</u>	<u>Soil Mapping Unit</u>	<u>Legal Description</u>
10 ³	Hilger cobbly loam, 0-40% slopes	750'S, 900'E of NW corner, Section 14, T6S, R5E.
11 ³	Hilger cobbly loam, 0-40% slopes	1000'S, 1650'W of NE corner, Section 14, T6S, R5E.
12 ³	Hilger cobbly loam, 0-40% slopes	920'S, 1600'W of NE corner, Section 14, T6S, R5E.
13	Bullflat silt loam, 0-3% slopes	500'E, 350'N of SW corner, Section 9, T5S, R6E.
14	Bullflat silt loam, 0-3% slopes	350'N, 250'W of SE corner, Section 8, T5S, R6E.
15	Hilger cobbly loam, 0-6% slopes	1600'N, 1550'E of SW corner, Section 11, T5S, R5E.
16	Hilger cobbly loam, 0-6% slopes	1700'N, 1400'E of SW corner, Section 11, T5S, R5E.

¹ Mound site descriptions are approximate (Custer County, SD, 5th principle
meridian.)

² Soils were identified as Vale silt loam in mapping, but because of limited
acreage were correlated as Tilford.

³ Soil could be an inclusion of another soil. Hilger soil most likely on
summit and shoulder landscape positions.

Table 11

Munsell Soil Colors from Mound and Adjacent Surface Soil (0-10 cm)

<u>Meters¹</u> <u>Distance</u>	<u>Mound</u>					
	<u>1</u>		<u>2</u>		<u>3</u>	
0.25a	A	4/3 3/3	A	5/3 3/3	C	4/6 3/4
0.25b	A	4/3 3/3	A	5/3 3/3	C	4/6 3/4
0.75a	A	4/2 3/2	A	4/3 3/2	C	4/4 3/3
0.75b	A	4/2 3/2	A	4/3 3/2	C	4/6 3/4
1.5a	A	4/2 3/2	A	4/3 3/2	C	4/6-4/4 4/3
1.5b	A	4/2 3/2	A	4/3 3/2	C	4/4-4/6 4/3
2.5a	A	4/2 2/2	A	4/3 3/2	C	5/3 4/4
2.5b	A	4/2 2/2	A	4/3 3/2	C	4/4 3/3
3.5a	-		A	4/3 3/2	C	4/3 3/3
3.5b	-		A	4/3 3/2	C	4/4 3/3
4.5a	-		-		C	4/4 3/4
4.5b	-		-		C	4/4 3/4
5.5a	-		-		C	4/3 3/3
5.5b	-		-		C	4/3 3/3
6.5a	-		-		C	4/2 3/2
6.5b	-		-		C	4/2 3/2
Cont a	A	4/2 2/2	A	4/2 2/2	A	4/3 3/3
Cont b	A	4/2 2/2	A	4/2 2/2	A	4/3 3/3

Table 11 (continued)

Meters Distance	<u>Mound</u>					
	<u>4</u>		<u>5</u>		<u>6</u>	
0.25a	B,C 5/4	C 4/3	B,C 5/4	C 4/3-D 4/4	C 4/6	4/4
0.25b	C,B 5/4	C 4/4	B,C 5/4	C 4/3-D 4/4	C 4/6	4/4
0.75a	-		B,C 5/4	C 4/3-D 4/4	C 4/6	4/4
0.75b	-		B,C 5/4	C 4/3-D 4/4	C 4/6	4/4
1.5a	C 4/4	3/4	A 4/3	3/3	C,B 4/4	C 3/4
1.5b	C 4/4	3/4	A 4/3	3/3	C,B 4/4	C 3/4
2.5a	-		A 4/2	3/2	B 4/4-C 4/6	C 4/2-D 4/6
2.5b	-		A 4/2	3/2	B 4/4	C 4/2
3.5a	C 5/3-5/4	4/3	-		A 4/3	3/3
3.5b	C 5/3-5/4	4/3	-		A 4/3	3/3
4.5a	A 4/3	C 3/3	-		-	
4.5b	A 4/3	C 3/3	-		-	
5.5a	A 4/3	B 3/2	-		-	
5.5b	A 4/3	B 3/2	-		-	
Cont a	A 4/3	B 3/2	A 3/3	3/2	A 4/2	3/2
Cont b	A 4/3	B 3/2	A 3/3	3/2	A 4/2	3/2

Table 11 (continued)

<u>Meters Distance</u>	<u>7</u>		<u>Mound 8</u>		<u>9</u>	
0.25a	B 5/4	C 4/4	C 4/6	4/3	B 5/2-C 5/4	A 4/2-C 4/6
0.25b	B 5/4	C 4/4	C 4/6	4/3	B 5/2-C 5/4	A 4/2-C 4/6
0.75a	B 4/4	C 3/4	C 4/6	4/3	A 5/3-C 5/4	A 4/2-C 4/6
0.75b	B 4/4	C 3/4	C 4/6	4/3	A 5/3-C 5/4	A 4/2-C 4/6
1.5a	B 4/4	C 3/4	C 4/3	3/3	A 4/3	4/2
1.5b	B 4/4	C 3/4	C 4/3	3/3	A 4/3	4/2
2.5a	B 4/2	3/2	-		A 4/3	4/2
2.5b	B 4/2	3/2	-		A 4/3	4/2
3.5a	A 4/3	3/3-3/2	-		A 4/2	3/2
3.5b	A 4/3	3/3-3/2	-		A 4/2	3/2
Cont a	A 5/1	3/1	C 4/3	3/3	A 4/2	3/2
Cont b	A 5/1	3/1	C 4/4	3/3	A 4/2	3/2

Table 11 (continued)

<u>Meters</u> <u>Distance</u>	<u>10</u>		<u>Mound</u> <u>11</u>		<u>12</u>	
0.25a	D 4/8	D 4/6	A 4/2	3/2	B 6/4-A 5/3	B 5/4-A 4/3
0.25b	D 4/8-C 4/3	D 4/6-C 3/3	A 4/3	3/2	B 6/4-A 5/3	B 5/4-A 4/3
0.75a	D 4/6-C 4/3	D 3/6-C 3/3	A 3/3	3/2	B 6/4-A 3/3	B 5/4-A 3/2
0.75b	D 4/6-C 4/3	D 3/6-C 3/3	A 3/3	3/2	B 6/4-A 3/3	B 5/4-A 3/2
1.5a	C 4/3	3/3	-		A 4/3-B 6/4	A 3/3
1.5b	C 4/3	3/3	-		A 4/3	3/3
Cont a	C 4/3	3/3	A 4/2	3/2	A 3/3	3/2
Cont b	C 4/3	3/3	A 4/2	3/2	A 3/3	3/2

Table 11 (continued)

Meters Distance	<u>13</u>		<u>14</u>		<u>15</u>		<u>16</u>	
0.25a	A 4/3	3/2	A 4/3	3/2	A 6/3-5/3	4/3	A 5/3	3/3
0.25b	A 4/3	3/2	A 4/3	3/2	A 6/3-5/3	4/3	A 5/3	3/3
0.75a	A 4/2	3/2	A 4/3	3/2	A 5/3	3/3	A 5/3	3/3
0.75b	A 4/2	3/2	A 4/3	3/2	A 5/3	3/3	A 5/3	3/3
1.5a	A 4/2	3/2	A 4/3	3/2	A 5/3	3/3	A 4/3	3/2
1.5b	A 4/3	3/2	A 4/3	3/2	A 5/3	3/3	A 4/3	3/2
2.5a	A 4/3	3/2	A 4/3	3/2	A 4/2	3/2	A 4/2	2/2
2.5b	A 4/3	3/2	A 4/3	3/2	A 4/2	3/2	A 4/2	2/2
3.5a	A 4/3	3/2	A 4/3	3/2	-		-	
3.5b	A 4/3	3/2	A 4/3	3/2	-		-	
4.5a	A 4/3	3/2	A 4/3	3/2	-		-	
4.5b	A 4/3	3/2	A 4/3	3/2	-		-	
5.5a	A 4/3	3/2	-		-		-	
5.5b	A 4/3	3/2	-		-		-	
Control a	A 4/3	3/2	A 4/3	3/2	A 4/2	2/2	A 4/2	2/2
Control b	A 4/3	3/2	A 4/3	3/2	A 4/2	2/2	A 4/2	2/2

¹ a and b represent separate composite samples taken at distances from mound center.

² A, B, C, and D are 10YR, 7.5YR, 5YR, and 2.5YR hues, and Munsell colors are for dry and moist soil samples, respectively.

period of time (mounds 2, 13, and 14) had soil colors similar to the original soil surface because prairie dogs collected surface soil for mound construction (Koford, 1958) and because vegetation growth on abandoned mounds added organic matter to darken soil surface color, respectively.

Younger mounds constructed in differing parent materials or older mounds actively or recently used had soil colors inherited from subsoil parent materials. Soil colors usually had higher values and chromas and frequently had more reddish hues than the adjacent undisturbed soil. Mounds 5 and 12 were thought to be young relative to other mounds studied. Parent material changes were near the soil surface so that a shallow burrow re-located different-colored strata to the surface. Mound 12 was constructed on the footslope of a ridge above the intersection of two drainage channels. Wind Cave National Park is at the edge of the Black Hills dome where sedimentary beds have been tilted. Soil parent materials change rapidly across slope if tilted beds are narrow. The color difference between mounds 11 and 12 probably is caused by this change. The two mounds were constructed about 25 meters apart.

Mounds 6, 7, and 9 were actively-used mounds in the Bison Flats dog town. Despite the large size of each mound, burrowing had not obliterated the original soil

surface beneath each mound. Burrowing probably was confined to central tunnels through which subsoil was moved to the surface. Subsoil in these mounds had reddish hues, higher (lighter) values and more yellowish chromas than the adjacent undisturbed soil.

Mound 3 and 4 soil colors are derived from parent materials of reddish (2.5YR and 5YR) hues. Soil colors would gradually become less reddish with time on the mound surface, but both mounds retained 5YR hues in the abandoned section of the Red Valley dog town. Mounds 3 and 4 are large, have steep side slopes which would accelerate runoff of precipitation, and are exposed to wind erosion and increased evapotranspiration. Plant regrowth is decreased on each mound because of droughtiness and soil melanization from organic matter progresses more slowly than on smaller mounds. Leaching of CaCO_3 is also reduced and surface soil pH values are higher because of reduced water infiltration. Furthermore, the air-dry clay surface on mounds will bind water at soil moisture tensions approaching or exceeding the range of plant water extractability. Low average annual precipitation (38 cm) in this section of the Red Valley also contributed to less rapid mound color changes compared to mounds constructed in towns on relatively moist sites.

B. Soil Texture

Interaction of soil-formation factors (parent material, climate, time, topography, and organisms) is often complex, yet noticeable patterns of soil texture distribution across mound surfaces were observed (Table 12). Surface soils generally became finer-textured moving from the mound center to adjacent undisturbed soil surfaces. Undisturbed surface soils were silt loam or loam texture beside all studied mounds.

Textural distribution between mound and non-mound soils varied from the same texture across mound areas to abrupt texture changes. Texture differences across mound sites are due in part to different soil horizons being excavated, because subsoil is exposed at the soil surface as a result of pedoturbation. Some differences can also be attributed to wind and water erosion of finer-textured sediments, especially from large mounds elevated much higher than the original soil surface. Koford (1958) referred to soil loss by wind erosion in drier months, but concluded that not enough is known about prairie dogs to measure their immediate or ultimate effects on erosion through mound-building activity or vegetation removal. Variability in slope, vegetation, climate, alluvium and residuum parent materials, and the depth and degree of pedoturbation are other factors related to soil texture distribution across mound sites.

Table 12

Soil Textures from Mound and Adjacent Surface Soil
by Ribbon Method (0-10 cm)

Meters from Mound Center	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
0.25a	Cl ¹	l	l	sil	sicl
0.25b	Cl	l	l	sil	sicl
0.75a	Cl	Cl	l	-	sicl
0.75b	Cl	Cl	l	-	sicl
1.5a	Cl	Cl	l	sil	sil
1.5b	Cl	Cl	l	sil	sil
2.5a	Csil ¹	Cl	l	-	sil
2.5b	Csil	Cl	sil	-	sil
3.5a	-	Cl	l	sil	-
3.5b	-	Cl	Cl	sil	-
4.5a	-	-	sil	sil	-
4.5b	-	-	sil	sil	-
5.5a	-	-	-	-	-
5.5b	-	-	-	-	-
6.5a	-	-	-	-	-
6.5b	-	-	-	-	-
Control a	sil	sil	sil	sil	sil
Control b	sil	sil	sil	sil	sil

Table 12 (continued)

<u>Meters from Mound Center</u>	<u>6</u>	<u>7</u>	<u>8</u>	<u>9</u>	<u>10</u>
0.25a	scl	sicl	sicl	sicl	sl
0.25b	scl	sicl	sicl	sicl	sl
0.75a	scl	sicl	sicl	sicl	sl
0.75b	scl	sicl	sicl	sicl	sl
1.5a	scl	sicl	sicl	sicl	sl
1.5b	cl	sicl	sicl	sicl	sl
2.5a	cl	sicl	-	sicl	-
2.5b	cl	sicl	-	sicl	-
3.5a	sil	sicl	-	sic	-
3.5b	sil	sicl	-	sic	-
4.5a	-	-	-	-	-
4.5b	-	-	-	-	-
Control a	sil	sil	sil	sil	1
Control b	sil	sil	sil	sil	1

Table 12 (continued)

Meters from Mound Center	<u>11</u>	<u>12</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>
0.25a	1	c	sl	sl	col ¹	sl
0.25b	1	c	sl	sl	col	sl
0.75a	1	cl	sl	sl	col	1
0.75b	1	cl	sl	sl	col	1
1.5a	-	1	sl	1	ccl ¹	1
1.5b	-	1	sl	1	ccl	1
2.5a	-	-	sl	1	1	1
2.5b	-	-	sl	1	1	1
3.5a	-	-	sl	1	-	-
3.5b	-	-	sl	1	-	-
4.5a	-	-	sl	1	-	-
4.5b	-	-	sl	1	-	-
5.5a	-	-	1	-	-	-
5.5b	-	-	1	-	-	-
Control a	sil	1	sil	sil	sil	1
Control b	sil	1	sil	sil	sil	1

¹ Cl=coarse loam. Csil=Coarse silt loam. sil=silt loam.
 1=loam. silcl=silty clay loam. c=clay. col=cobbly loam.
 ccl=cobbly clay loam. scl=sandy clay loam.
 cl=clay loam. sic=silty clay. sl=sandy loam.

Few studies have dealt with relationships between rodent burrowing and soil texture, concentrating instead on the volume of soil raised and number of mounds created by excavation activity (Grinnell, 1923; Miller, 1948; Miller and Bond, 1960; Reid et al., 1966, and others). Deep, well-drained soils without excessive gravel or sand content represent typical sites for prairie dog colonization (Hardy, 1945; Sheets et al., 1971; Dalstead et al., 1981), although Thorp (1949) studied a colony of prairie dogs established in soil with silt loam surface texture underlain by sand and gravel 2 to 3 meters below the soil surface where prairie dog pedoturbation altered the mound soil to a loam texture. Crotovinas are a Russian term used to describe abandoned tunnels and nests of rodents filled in with soil material (Joffe, 1949; Hole, 1981) which contribute to redistribution of different-colored and textured soils. However, the most comprehensive study of prairie dogs noted only that inhabited soils were "relatively heavy" (fine-textured) and that burrows occur in many kinds of soil (Koford, 1958). Mounds are apparently limited to soils with enough fine-textured soil material to preserve tunnels and galleries from caving in from water infiltration and increased sediment loads deposited on the mound surface by subsoil excavation.

Excavation of an underlying contiguous bed of

finer-textured parent material may explain the increasingly coarser-textured soil surface moving away from the mound 12 center. Soils were clay texture surrounding the mound center, clay loam across the mound surface, and loam texture on the non-mound soil surface. The mound is part of post-1982 colony expansion onto poorly-vegetated slopes at the southern edge of the Bison Flats dog town. Time has been insufficient for erosion to move the clay-sized soil fraction downslope. Several authors noted increases in fine-textured soil material as a component of termite (Nye, 1955; Lee and Wood, 1971) and ant mounds (Baxter and Hole, 1967; Salem and Hole, 1968; Cowan et al., 1985) and worm casts (Thorp, 1949; Hole, 1961). However, prairie dog subsoil excavation appears to be at best non-selective regarding particle size. Surface soil texture is more likely influenced by subsoil parent material and by climatic and topographic effects after soil excavation.

Mounds 5, 7, 8, and 9 had slight texture changes from silty clay loam to silt loam across mound to non-mound surface soils. Uniformity of subsoil parent material and insufficient time for slope and wind and water erosion to affect mound surface soil texture may all be factors in the nearly uniform texture distribution across these mounds. In support of the above ideas, mounds 7, 8, and 9 are in active use. Continual excavation of finer-textured subsoil would counteract climatic and topographic influences on

mound soil texture. In addition, the site of mounds 8 and 9, near the depressional area of an old streambed, has been subject to stream deposition of various uniform-textured sediment layers.

Coefficients of variation (CV's) in clay content of Mollisols formed in alluvium or residuum were found to be as high as 57 percent in Mausbach's (1980) study on soil variability, with CV's of Entisols (mound 9) up to 61 percent.

No large rocks were found in mound soils at study sites. Hansen and Morris (1968) recorded similar findings for northern pocket gophers (Thomomys talpoides). Little movement of rocks greater than 2.5 cm in diameter was noted, and relatively fewer rocks were moved under dry soil conditions and when soils contained few large rocks.

C. Soil pH

The pH of most mound soils decreased with increasing distance from the mound center (Table 13). Mean pH values of mound surface soils are significantly different from non-mound surface soils except for mound 11 (Table 14).

Soil pH varies little (CV = 15 percent) between soil profiles within landscape units of a few hectares (Wilding et al., 1983). Mausbach et al. (1980) examined groups of pedons, each representing different parent mate-

Table 13

Average 1:1 pH of the 0-10 cm layer of Prairie Dog Mounds with Increasing Distance from the Mound Center.

<u>Mound</u>	<u>0.25</u>	<u>0.75</u>	<u>1.5</u>	<u>2.5</u>	<u>3.5</u>	<u>4.5</u>	<u>5.5</u>	<u>6.5</u>	<u>Non-Mound¹</u>
1	7.43	7.39	7.12	6.56	-	-	-	-	6.32
2	7.00	7.12	7.09	6.76	6.34	-	-	-	6.08
3	7.71	7.64	7.64	7.60	7.59	7.34	7.17	6.84	6.49
4	7.88	-	7.78	-	7.52	7.29	6.77	-	6.60
5	7.82	7.76	7.50	7.25	-	-	-	-	6.74
6	7.43	7.42	7.16	6.81	6.13	-	-	-	6.05
7	7.61	7.69	7.68	7.43	6.99	-	-	-	6.00
8	7.51	7.55	7.43	-	-	-	-	-	7.38
9	7.63	7.60	7.56	7.50	7.12	-	-	-	6.83
10	6.44	6.88	6.52	-	-	-	-	-	6.47
11	6.76	6.82	-	-	-	-	-	-	6.75
12	7.39	7.58	7.28	-	-	-	-	-	6.89
13	5.31	5.45	5.69	5.71	5.74	5.75	5.85	-	6.00
14	5.80	5.77	5.73	5.86	5.76	5.90	-	-	6.01
15	7.67	7.60	7.47	6.78	-	-	-	-	6.00
16	7.31	7.20	7.29	6.15	-	-	-	-	6.01

¹ Average of two composite samples taken 3 meters beyond the mound edge.

Table 14

Analysis of Variance for Average 1:1 pH of Mound
Surfaces Compared to Adjacent Non-Mound Soil.

<u>Mound Number</u>	<u>Mound</u>	<u>Soil</u>	<u>df</u>	<u>F¹</u>
1	7.13	6.32	4	76.17**
2	6.86	6.08	5	18.90**
3	7.45	6.49	8	62.82**
4	7.45	6.60	5	52.96**
5	7.58	6.74	4	113.43**
6	6.99	6.05	5	24.38**
7	7.48	6.00	5	432.74**
8	7.50	7.38	3	20.86*
9	7.48	6.83	5	71.97**
10	6.61	6.47	3	32.88*
11	6.79	6.75	2	0.80(NS)
12	7.42	6.89	3	24.52*
13	5.64	6.00	7	6.44*
14	5.80	6.01	6	6.24*
15	7.38	6.00	4	132.51**
16	7.01	6.01	4	54.90**

¹ **=significant at 1% level of probability.

*=significant at 5% level of probability.

(NS)=not significant.

rials, up to 32 kilometers apart and found A horizon pH values with CV's of 10 to 16 percent for residuum and alluvium-derived soils. Comparisons within mounds and within colonies are more statistically meaningful with respect to mound ages and the degree of chemical and physical alteration of soil when soil properties with limited variability are measured.

Calcium is the dominant exchangeable cation in most plains soils. Lime (CaCO_3) leached from A and upper B horizons can accumulate in Bk and C horizons of warm, very dry plains soil (Westin and Malo, 1978). Excavated calcareous subsoil deposited at the soil surface increases mound surface pH values relative to leached, undisturbed non-mound soil. Differences of greater than 1.0 pH unit between mound and undisturbed soil surfaces were common, especially on the majority of older and larger mounds. Older mounds are composed of a greater volume of subsoil than recently-constructed mounds. Leaching of carbonates would be inhibited by frequent addition of calcareous subsoil at the mound center. Accordingly, pH values would remain high for a longer time there.

Mounds 8, 10, and 11 were constructed in Hilger cobbly loam since post-1976 expansion of the Bison Flats dog town. Younger mounds are composed of appreciable amounts of surface soil (Koford, 1958). Depth of burrowing may be insufficient to relocate calcareous subsoil at the

surface of mounds 10 and 11, and pH values are uniform for mound and non-mound surfaces as a result. The mound 8 site, on the footslope position of a toposequence above an old streambed, may also have had additions of calcareous Winetti cobbly loam surface soil eroded from upslope. These additions could increase non-mound surface soil pH values so that variability in pH between mound and undisturbed (non-mound) soil surfaces was not evident.

Long-term nutrient cycling through vegetation could explain pH increases moving across mound to non-mound soil surfaces at mounds 13 and 14. Both are old, abandoned mounds constructed on a level terrace. Western wheatgrass grew densely on non-mound soil but was sparse on mounds. Scattered clumps of little bluestem were found on mounds but were infrequent on adjacent non-mound soils. Decomposing plant litter high in calcium and other bases may have partially neutralized soil surface acidity on undisturbed soil (Jenny, 1980). Calcium leached from mound surface soils as CaCO_3 was not recycled upward as effectively or for as long a period of time by little bluestem and shallow-rooted annual plants which invaded the disturbed soil after mound abandonment.

Sandy loam soil textures found across much of mound 13 and the center of mound 14 would be more well-drained, arid locations than silt loam-textured undis-

turbed soils. White (1961) observed little bluestem growth confined to weakly-developed soils of gilgai microridges in some areas of western South Dakota. Microridges are similar to mound soils because pedoturbation (physical and biological, respectively) created weakly-developed soil profiles in both cases. White also noted that western wheatgrass tended to occupy moist microvalley positions between microridges. Finer-textured non-mound soils are more structurally well-developed, favoring establishment of western wheatgrass and propagation of that species by rhizomes. Long-term nutrient cycling without soil disturbance would counter leaching of bases more effectively than on mound soils, where leaching may not have been compensated for by nutrient cycling.

Statistical comparisons of pH values between mounds from the same study area generally were not significant (Table 15). Mound 2 is likely much older than mound 1 since past forested soil colors have been obscured by increased pedoturbation relative to mound 1. Lower surface pH values across mound 2 are the result of longer mound surface exposure to calcium leaching compared to mound 1. Mounds 10, 11, and 12 are contiguous and were constructed on cobbly, sloping sites with tilted beds of sedimentary rock. Notable color differences between mounds support the idea of different parent materials at these three

Table 15

Analysis of Variance for Average 1:1 pH Values Between Selected Mounds from the Same Study Area.

<u>Mound Comparison</u> ¹		<u>average pH</u>		<u>pH range</u>		<u>F</u> ²
<u>a</u>	<u>b</u>	<u>a</u>	<u>b</u>	<u>a</u>	<u>b</u>	
1 vs 2		7.04	6.73	6.30-7.46	6.07-7.18	26.30*
3 vs 4		7.28	7.42	6.47-7.71	6.54-7.99	8.15(NS)
6 vs 7		6.96	7.09	5.98-7.50	5.96-7.73	11.61(NS)
8 vs 9		7.48	7.35	7.37-7.58	6.73-7.68	8.16(NS)
10 vs 11		6.59	6.78	6.44-6.90	6.71-6.84	38.20*
11 vs 12		6.78	7.29	6.71-6.84	6.88-7.60	514.48**
13 vs 14		5.59	5.86	5.21-6.04	5.71-6.01	14.50(NS)
15 vs 16		7.09	6.84	5.89-7.74	5.82-7.32	10.22(NS)

¹ inner two sampling rings and control samples from each mound were used for mound comparisons.

² **=significant at 1% level of probability. *=significant at 5% level.
(NS)=not significant below 10% (0.1) level of probability.

mounds of similar ages. Younger-aged mounds would have a pH distribution influenced primarily by parent material; Large pH differences between parent materials would be statistically significant.

Over time, climatic effects on mound soil pH become increasingly important, homogenizing pH differences between mounds by leaching greater relative quantities of CaCO_3 from calcareous mound soil than from non-calcareous mounds. Significant pH differences between older mounds from the same study area are in this way less likely.

D. Total Phosphorus in Surface Layers

Total phosphorus contents of the soil surface layer (0-10 cm), analyzed in 1985, tended to decrease with distance from the mound center (Table 16). Older mounds generally had significantly more total phosphorus in surface layers than adjacent undisturbed soil. Younger mounds had fairly uniform total phosphorus in mound and undisturbed surface layers (Table 17).

Differences in total phosphorus were not always significant. Comparisons made between mounds from the same study area were not significant except for mounds 10 and 11. However, when orthogonal comparisons were used to maximize degrees of freedom and partition the sum of squares, more significant differences in phosphorus contents were found between mound and non-mound soil surfaces

Table 16

Mean Total Phosphorus Content (mg Kg⁻¹) of the 0-10cm layer of Prairie Dog Mounds with Increasing Distance from the Mound Center.

<u>Mound</u>	<u>0.25</u>	<u>0.75</u>	<u>1.5</u>	<u>2.5</u>	<u>3.5</u>	<u>4.5</u>	<u>5.5</u>	<u>6.5</u>	<u>Non-¹ Mound</u>
1	718	718	640	556	-	-	-	-	587
2	679	636	676	646	673	-	-	-	703
3	710	680	678	634	624	609	526	474	474
4	680	-	656	-	684	561	516	-	521
5	700	680	624	616	-	-	-	-	591
6	609	660	671	686	691	-	-	-	730
7	862	832	834	764	798	-	-	-	804
8	870	808	905	-	-	-	-	-	850
9	848	837	761	812	718	-	-	-	742
10	126	136	230	-	-	-	-	-	260
11	481	392	-	-	-	-	-	-	441
12	590	482	387	-	-	-	-	-	363
13	690	675	693	705	618	555	503	-	543
14	602	573	612	643	607	565	-	-	547
15	650	616	612	441	-	-	-	-	435
16	500	539	453	479	-	-	-	-	488

¹ Average of two composite samples taken 3 meters beyond the mound edge.

Table 17

Statistical Analysis of Mean Phosphorus Contents of Mound Surfaces Compared to Non-Mound Soil.

<u>Mound Number</u>	<u>Total P (mg Kg⁻¹)</u>		<u>df</u>	<u>F¹</u>
	<u>Mound</u>	<u>Soil</u>		
1	658	587	4	12.21*
2	662	703	5	2.48(NS)
3	617	474	8	9.38*
4	619	521	5	7.84*
5	655	591	4	33.80**
6	663	730	5	20.89**
7	818	804	5	1.24(NS)
8	844	850	3	1.84(NS)
9	795	742	5	1.51(NS)
10	164	260	3	11.67*
11	437	441	2	8.78(NS)
12	486	363	3	42.10**
13	634	543	7	12.15*
14	600	547	6	8.01*
15	580	435	4	12.70*
16	493	488	4	1.15(NS)

¹ **=significant at 1% probability. *=significant at 5% probability.

(NS)=not significant at $\leq$ 5% probability.

(Table 18). Within mounds, pedoturbation could homogenize total phosphorus contents if mound soils were thoroughly mixed during excavation. On the other hand, incomplete mixing of soil material, parent material variability, slope, climatic, and biotic factors make predictions of surface phosphorus contents difficult.

Smith et al. (1950) observed decreasing total phosphorus contents with depth in Tama silt loam in Iowa to the A3 and increases in the B horizon before reaching a maximum in the C horizon. Likewise, Lipps and Chesnin (1950) found maximum phosphorus contents in the parent material of Chernozem soils. Total phosphorus contents were variable in horizons above parent materials for different soils. They attributed the accumulation of phosphorus in surface horizons to upward cycling by rhizosphere plants.

Other studies have found total phosphorus contents decreased gradually or irregularly with depth (Alway and Rost, 1916; Nye and Bertheux, 1957; White and Gartner, 1974a). Total phosphorus contents of small, young mounds may be more directly influenced by parent material phosphorus levels than older and larger mounds. Small mounds with shallowly-excavated burrows contain smaller quantities of subsoil relative to surface soil, whereas older mounds are primarily subsoil (Tileston and Lechleitner, 1966). In addition, fecal wastes and phosphorus are less likely to accumulate on the small surface area of young mounds.

Table 18

Orthogonal Comparisons of Mean Total Phosphorus Content Between Selected
Mound and Undisturbed Surface Soils (0-10cm).

<u>Mound Numbers</u>	<u>Samples</u>		<u>Significance²</u>
	<u>Mound¹</u>	<u>Soil</u>	
	(mg Kg ⁻¹)		
1,2,3,4,5,13,14	671	568	F = 0.01
6,7,8,9,10,11,12,15,16	595	568	F = 0.05
6,7,8,9,10,11,12	599	598	NS
15,16	583	461	F = 0.01
13	682	543	F = 0.01
14	588	547	NS
1	718	587	F = 0.01
2	657	701	NS

¹ Mound soil is represented by the two inner sampling rings.

² F = 0.01, F = 0.05, and NS are significant at 1%, 5%, and not significant at $\leq$ 5% probability, respectively.

Significant differences between mound and non-mound phosphorus contents, particularly mounds 10 and 12, may be due to an abrupt parent material change near the soil surface. Subsoil may have been used in greater proportion than surface soil to construct each mound.

Mound 15 had significantly more total phosphorus on mound soils relative to non-mound soil because of large numbers of fecal pellets which had been excreted on the mound surface or excavated there from subsoil tunnels and galleries.

Surface layer phosphorus contents were markedly different between mounds 1 and 2 despite their location 100 meters apart. Mound 1 had a significantly higher surface layer phosphorus content in mound compared to non-mound soil, yet mound 2 is thought to be older than mound 1 because of the larger size, greater area of disturbed soil beneath mound 2, relative absence of previously-forested soil colors with depth, and lower surface pH values in the mound 2 soil. Phosphorus distribution in mound 2 fails to support the preceding evidence. Mound 2 may have been abandoned soon after subsoil was deposited on the mound surface. In this case, phosphorus contents on the mound surface would reflect parent material phosphorus levels instead of phosphorus accumulated from prairie dog activity. Mound 2 may also have been used intermittently

for a long period of time so that phosphorus accumulation had not proceeded at a rate comparable to mound 1. Greater downhill movement of eroded, phosphorus-rich colluvial sediments may explain the higher total phosphorus contents on non-mound surfaces at mound 2. In addition, Koford (1958) found that badgers (Taxidea taxus) were not only common in Wind Cave National Park but that they were capable of digging to the deepest recesses of burrows, bringing the lining of prairie dog nests to the surface. Pedoturbation of the mound 2 surface during badger digging could partially alter mound surface phosphorus distribution, although no large-scale digging in the mound 2 surface was apparent.

Mound Formation Rate

Area of disturbed original soil surface beneath mounds was proportional to total mound surface area (Table 19). Measurements of the relationship were plotted logarithmically, obtaining a high coefficient of determination ($r = 0.78$) between total and disturbed soil surface areas (Figure 14). Variability about the regression line is dependent on the amount of surface soil scraping by prairie dogs, years of mound activity, ease of subsoil excavation, and availability of adjacent sites for mound construction. Mounds appeared to be more numerous and smaller in the Bison Flats dog town relative to other studied areas. Available area for colony expansion is great here (Lovaas, 1972). In the Red Valley, the hogback ridge and shallow, rocky soil profile favored enlargement of existing mounds and limited colony expansion due to excessive slope.

As time of mound use by prairie dogs increases, subsoil additions to mounds could be expected to increase if subsoil is friable enough for the construction of longer tunnels and more galleries. Steep slopes and shallow soils were found to impede mound formation in most areas of the park; Smaller, younger mounds were located on many of these sites of marginal prairie dog habitat if colony expansion had made emigration necessary.

Soil samples collected from the buried soil surface

Table 19

Total Mound Surface Area and Area of Disturbed
Original Soil Surface.

<u>Mound</u> ¹	Mound Area (m ²)	
	<u>Total</u>	<u>Disturbed</u>
1	15	4
2	30	20
3	161	116
4	140	96
5	13	0.3
6	50	1
7	44	3
8	13	4
9	36	1
10	7	0.2
11	2	0.2
12	8	0.2
13	128	42
14	98	13
15	21	4
16	12	0.3

¹ Mounds are grouped according to the five major areas studied at Wind Cave National Park.

(1984) and surface of mound 1 (1985) were C14-dated as 1340+-80 and 190 years old. Ruhe (1983) indicated that C14 dates estimate the mean residence time of soil organic matter. He reported C14 dates of 410+-110, 840+-200, and 1545+-110 years before present at 15, 38, and 61 centimeter mean sampling depths of a southern Iowa soil. Actively-growing plant roots make up a smaller portion of soil organic matter with depth. Average age of organic matter therefore increases with depth where pedoturbation is insufficient to redistribute organic matter more uniformly. In comparison to Ruhe's data, the original soil surface of mound 1 was likely 400 to 800 years old when buried during mound construction and the C14 date has increased over time as easily oxidizable organic matter has been decomposed in the original soil surface beneath mound 1. Four hundred years is a realistic minimum date for the age of mound 1.

Mound 3 is located in the Red Valley in a warmer, drier area of the park 180 meters lower in elevation than mound 1. Profile characteristics and mound size indicate that forest encroachment into grassland was not a factor in this section of the Red Valley and that mound development likely progressed uninterrupted for thousands of years. The extreme drought period 7,000 years before present coincided with the time grasslands were most extensive along grass-forest ecotones and probably represents

the time mound 3 was first constructed. Food shortages due to drought or overgrazing are a cause of dog town expansion (Koford, 1958). When vegetation became sparse, parts of dog towns were abandoned and prairie dogs moved toward more vegetated areas on town margins (Bailey, 1926). Digging of new burrows and vegetation modification by prairie dogs made sites favorable for future and continued occupation. Mounds constructed near mound 3 were likely used continuously because of difficulty in tunneling through the shallow subsoil boulder layer.

Using age estimates of mounds 1 and 3 and disturbed soil area beneath mounds, a linear scale of mound ages was plotted (Figure 15). Ages are consistent with known or estimated ages for mounds except for mounds 8, 9, and 15. These mounds may have been recolonized after abandonment or poisoning. Bison Flats dog town, site of mounds 6 through 12, was cited by Lovaas (1972) as having very great potential for expansion. Prairie dog control and eradication programs in 1952 and 1953 used poisoned grain to eliminate prairie dogs in large areas of this town. Many mounds found at Bison Flats may therefore have been recently recolonized instead of constructed. Mounds 6 and 7 were constructed before 1976, while mounds 8 and 9 and mounds 10, 11, and 12 were thought to have been constructed after 1976 and 1982, respectively. Intermittent mound use

before or after observed continuous colonization and different rates of subsoil excavation both affect the linear relationship between variables used in Figure 15.

It is significant that the three oldest and largest mounds (mounds 3, 4, and 13) also have the greatest decreases in total phosphorus from inner to non-mound surfaces (except for mounds 12 and 15). The three mounds may be examples of long-term phosphorus accumulation by prairie dog activity. Mound 12 likely exposed dissimilar parent materials during construction, while mound 15 had numerous fecal pellets scattered across the mound surface which became part of collected soil samples at that site. If phosphorus-rich subsoil were the explanation for significant differences between mound and non-mound soil surface total phosphorus contents at mounds 3 and 4, then similar large decreases in surface total phosphorus contents between mound and undisturbed soils at mounds 6 and 7 would also be expected since Tilford silt loam is found at each site. No such trend is noticeable. In fact, outer sampling rings still within the mound area (6.5, 3.5, and 5.5 for mounds 3, 4, and 13, respectively) have total phosphorus contents quite similar to adjacent non-mound soil surfaces at each site. Since the outer mound surfaces are composed partially or wholly (mound 3) of subsoil and probably have had less phosphorus accumulation through prairie dog activity compared to nearer the mound center, it is

likely that large differences in total phosphorus contents with depth do not exist in these soil profiles. Significant differences in total phosphorus across mound surfaces could therefore be attributed to prairie dog influences on mound soils instead of large differences in phosphorus content with depth.

In addition, younger mounds often had small decreases or increases in total phosphorus across mound to non-mound surfaces. Phosphorus-rich subsoil may therefore not be a factor in distribution of total phosphorus contents across younger mounds. In most cases, recent mound construction precludes phosphorus accumulation on mound surfaces because fecal, carcass or bone, and plant materials have not been incorporated into mound soil or deposited on younger mound surfaces as frequently as on older mounds. Younger mounds also tend to be smaller in size relative to older mounds. Prairie dogs may therefore spend less time confined to the dimensions of a smaller mound, which would have accumulated phosphorus more rapidly than a larger mound if excretion occurred frequently on the smaller mound surface.

General Relationships

Prairie dogs both influence and are influenced by interaction of five soil-forming factors: parent material, climate, organisms (biotic factor), topography, and time. Each determines the expression of observable and measurable soil properties across a landscape.

Relationships between dog towns and vegetation are visible in the field, but there is a notable lack of research concerning influence of burrowing animals on vegetation (USDA, 1978). Prairie dogs disturb soil through excavation and cut down tall plants within dog towns, favoring establishment of edible forbs and short grasses and removing obstacles to vision and movement (Koford, 1958). Past authors often attributed range depletion to prairie dog colonization (Merriam, 1902; Bell, 1921; Taylor and Loftfield, 1924; Bailey, 1926) without considering that thriving populations of prairie dogs may have been caused by range overgrazing of cattle. The resulting flora would be composed of succulent young-growth short grasses and forbs which would promote expansion of dog towns.

Zonation of vegetation due to prairie dog colonization is more apparent in active or recently-abandoned towns than in long-abandoned towns. In the former cases, prairie dogs have maintained a vegetative subclimax within

mound and town areas through recent foraging activity and cutting of tall plants. Osborn and Allen (1949) described vegetation zonation around a studied prairie dog town as representing seven stages of plant succession found locally where succession proceeded without environmental disturbance. The central town area represented the earliest successional stage (bare ground and mat forbs), with concentric rings outward from the town center representative of more advanced stages of plant succession. The outer vegetation zone of Osborn and Allen was composed of climax-stage tall grasses (big and little bluestem and indian-grass).

Mounds 6 through 12 and mounds 15 and 16 exhibited vegetational zonation, although zonation was not observed to be entirely concentric or composed of many plant successional stages on or near these active mounds. Plant growth was limited around each mound relative to adjacent areas outside dog towns. Silvery wormwood and red threeawn were only somewhat abundant on or near mounds 6 through 12 at Bison Flats, while short, isolated stands of western wheatgrass grew only at a distance from each mound. Mounds 15 and 16 exhibited abrupt changes in vegetation. Silvery wormwood, white sage, and pussytoes (Antennaria spp.) grew on and near mound surfaces. The southern edge of the town was bordered by a tall, dense stand of little bluestem which grew as close as 10 meters from mound 16. Close

proximity of little bluestem to active mounds was indicative of recent colonization of this area of Rankin Ridge by prairie dogs.

In the latter case, plant succession proceeds toward climax stage after mounds and towns are long-abandoned or have too small a population of prairie dogs to maintain a vegetative subclimax. Mounds 13 and 14 were apparently long-abandoned. Western wheatgrass grew profusely around but sparsely on mound surfaces, while bluegrama and buffalo-grass were identified as understory grasses on and around each mound. Scattered clumps of little bluestem were found on both mounds. No forbs grew in fair abundance on or near mounds 13 and 14 except for yellow sweet clover (Melilotus officinalis). In the absence of prairie dog foraging and cutting, more advanced stages of plant succession and taller plants have replaced or are replacing bare ground and forbs at sites formerly occupied by prairie dogs. Similarly, Osborn and Allen (1949) observed plant succession and abandonment of a prairie dog town for 10 years after private grazing was terminated. Plant succession proceeded towards climax stage tall grasses, and prairie dogs were unable to maintain enough of a shortgrass subclimax within the colony to meet nutritional needs. Prairie dogs were thought to occupy the study area only with the assistance of heavy cattle grazing to favor the

shortgrass successional stage. Osborn and Allen's study lends credibility to the idea that prairie dog colonization in itself is an effect of vegetative conditions rather than a cause.

Fewer and larger mounds were present in areas 1 (mounds 1 and 2), 2 (mounds 3, 4, and 5), and 4 (mounds 13 and 14). Small prairie dog populations which appeared to have existed at these sites may have been unable to maintain shortgrass subclimax at the expense of taller plant encroachment from the borders of these smaller towns. Furthermore, survival of colonies is higher where more individuals can detect danger or where a catastrophe such as flooding or predators is less likely to eliminate an entire dog town (King, 1955).

Recent work investigating ecological relationships between herbivores and vegetation has been undertaken at Wind Cave (Coppock et al., 1983a, b; Detling and Painter, 1983; Archer and Detling, 1984; Ingham and Detling, 1984). Significant findings included that prairie areas uncolonized by prairie dogs had the greatest live plant biomass per unit area and that 3 to 8 years of prairie dog colonization, although producing greatest total diversity of plant species on dog towns, produced the least live plant biomass (Coppock et al., 1983a). Forbs and dwarf shrubs composed >95 percent of biomass on old prairie dog towns while graminoids made up >85 percent of plant biomass on

uncolonized prairie. Dead plant biomass and litter declined on town sites and significantly less leaf biomass was produced from tillers of big bluestem and threadleaf sedge (Carex filifolia) as prairie dog colonization time increased, thought to be due to increased competition for and multiple defoliation of tillers (Coppock et al., 1983a; Ingham and Detling, 1984). Western wheatgrass growing within studied prairie dog towns adapted to multiple defoliations by producing greater plant leaf blade/leaf sheath ratios, more horizontal leaves, fewer leaves/tiller, and narrower leaf blades relative to western wheatgrass within grazing exclosures (Detling and Painter, 1983). Studies of plant growth form and defoliation were not attempted here, but observations made in the field support plant biomass and composition findings cited by the previous authors.

Removal of plant cover on and near mounds affects soil properties there in various ways. Total plant biomass is reduced by immediate removal of plant cover and by long-term reduction of plant growth after multiple defoliation and subsequent slower growth rates of edible plants. Poor plant cover on and near mounds leads to decreased levels of soil organic matter and nitrogen. Long-term plant removal on mounds 1 through 3 may be a cause of lower soil surface organic matter levels relative to non-mound surface soils.

However, results of nitrogen and phosphorus analyses indicate that addition of nutrient-rich fecal material partially or wholly compensates for forage removal before mound abandonment, especially on mound 3. Loss of organic matter from plant sources is highest on and near mounds and should decrease with distance from the mound center if foraging activity reduces plant cover to a greater extent at and near the mound site.

Vegetation removal at mound sites also affects surface soil microclimate within prairie dog towns. Decreased soil organic matter and more direct incidence of solar radiation and wind on exposed soil decreases soil water-holding capacity and dries soil surfaces into a hard crust after mounds are abandoned or subsoil is no longer excavated. Infiltration of precipitation is thereby reduced along with increased soil surface temperatures. Vegetation establishment on mounds may therefore proceed slowly even after mound abandonment, although regrowth of vegetation on mounds and dog towns may also be rapid (Koford, 1958, p. 60). Presumably Koford's observation was the trend on mounds (younger than those observed in this study) where soil drying was not intense and enrichment of mound soil with excreta enhanced plant growth. Expansion of dog towns would proceed most rapidly when plant growth is limited by overgrazing or drought. The 1976 expansion of Bison Flats dog town, also the approximate construction

date of mounds 8 and 9, may have been a response to poor forage growth at existing town sites caused by very dry soil conditions that year.

In addition, mound soils become increasingly susceptible to wind and water erosion in the absence of plant cover. In locations where prairie dog colonies are dense there are large areas of bare to poorly-vegetated soil. These areas tend to become larger in times of drought or when grass cover is repeatedly clipped close to the soil surface by prairie dogs and other wildlife. Wind erosion is a common sight on prairie dog colonies in the Conata Basin of western South Dakota (USDA, 1978).

Subsoil deposited in mounds is less chemically weathered and would initially support less vegetation than A horizon soil present before pedoturbation. Furthermore, wind erosion of mounds removes greater quantities of phosphorus-bearing soil than are removed from lower inter-mound areas. Water erosion may also redeposit phosphorus-rich soil from higher central mound surfaces so that phosphorus accumulates toward or around the mound rim, depending on mound texture, chemical nature of subsoil material, length of time phosphorus has accumulated on the mound, and mound topography.

Prairie dogs compensate for phosphorus erosion from mound surfaces by addition and incorporation of fecal

pellets on mounds. Young, recently-excavated mounds 8, 10, 11, 15, and 16 were covered with fecal pellets in various stages of decomposition at the time of sampling. Pellets may also have been redeposited on the mound surface after excavation from tunnels and galleries within each mound. Increased quantities of excavated subsoil are mixed with fecal material as mounds are enlarged. In this way a larger volume of soil across and within each mound becomes enriched with phosphorus from fecal pellets.

Bone and carcass remains may also be excavated and deposited on mounds. Although no prairie dog carcasses were observed in this study, prairie dog bones were often found on younger-aged mounds. Animal wastes and bone are known to be high in phosphorus content (Waggamann, 1969), although phosphorus in bone would be slowly available for combination with soil exchange sites relative to fecal phosphorus. It was thought that the absence of bones on old mounds was due to the fact that each mound was no longer a completely bare, exposed area of soil and that bones had been removed by other organisms in the time passed since mound abandonment, notably mammals smaller than prairie dogs. Plant residue wastes dropped on mounds as part of plant cutting or foraging or excavated from nests within mounds could also contribute small quantities of phosphorus to mound surface soils.

The net effect of phosphorus redistribution by the

sources mentioned is likely to be an accumulation of phosphorus in mound material as mounds are enlarged and inhabited by prairie dogs over time. Soil phosphorus levels were not measured adjacent to dog towns on uncolonized areas. However, phosphorus concentrations there would probably be intermediate to mound (high phosphorus) and defoliated non-mound (lower phosphorus) soils.

SUMMARY AND CONCLUSIONS

Pedoturbation by prairie dogs alters soil development through:

1. Burrowing, tunneling and excavation of soil in construction of crater or dome-shaped mounds.

With time, prairie dog activity creates:

2. Mound soils lighter in color, coarser-textured in the mound center, higher in pH and total phosphorus, lower in organic matter, and poorly-vegetated compared to adjacent non-mixed soils.

These soil changes are more permanent than prairie dog modification of vegetation at mound sites, but may also be less visible in examination of dog towns. Soil properties can be used to estimate prairie dog mound ages or at least the length of time the mound has been used on a intermittent or continuous basis.

Sampling procedures and analyses conducted in the 1984 study investigated vertical distribution of mound and adjacent non-mound soil properties across three large prairie dog mounds. Soil samples from the 1985 study were collected from sampling ring grids designed to systematically determine collection points across 16 mounds to a depth of 10 cm.

Addition of greater volumes of subsoil to mound surfaces over time results in lighter-colored mound soil relative to adjacent undisturbed surface soil. Mound soil surfaces may be coarser-textured than non-mound surfaces due to removal of finer-textured sediments from mounds by wind and water erosion. Redeposition of carbonates at mound surfaces of excavated subsoil raised soil pH values across and with depth in mounds, although influences of long-term nutrient cycling, parent material differences, and variable ages of studied mounds may create exceptions to this finding.

Accumulation of phosphorus in mound soil was apparent at several mound sites where mounds were believed to have been inhabited by prairie dogs for long periods of time. Phosphorus accumulation was also found more frequently on large mounds. Mounds are enlarged with time and phosphorus accumulates on mounds used over many years. Prairie dog excretion habits were not observed in this study or documented by other authors, but it is believed that excreta are more likely deposited on large mound surfaces since a proportionately greater degree of prairie dog activity would be confined to the dimensions of a large mound compared to a small, recently-constructed mound.

Removal of vegetation from mound and non-mound soil surfaces during colonization will decrease quantities of phosphorus present in plant material there, but increases

in fecal and plant phosphorus within mounded soil and fecal phosphorus on mound surfaces due to prairie dog activity are likely. The role of wind and water erosion in the removal of phosphorus-bearing soil sediments from mound surfaces is difficult to quantify, but is thought to be a factor contributing to phosphorus removal from large, elevated mound surfaces. The net result of long-term prairie dog activity is accumulation of phosphorus in mound soil areas over time.

Other localized effects of mound construction include decreased soil organic matter on mound surfaces, less granular soil structure and more soil surface crusting on abandoned mounds, reduced total plant biomass in active dog towns, drier and warmer mound soil surfaces, and increased likelihood of wind and water erosion from exposed or poorly-vegetated mound and undisturbed soil surfaces.

Ideally, prairie dog influences on soil development could be isolated from related biotic, chemical, and physical factors to examine colonization and mound construction effects on soil properties. Although this idealism is not totally possible, results obtained from this study may be used as guidelines in further research involving pedoturbation by burrowing rodents.

Bibliography

- Alway, F. J., and C. O. Rost. 1916. The vertical distribution of phosphorus in the horizons of some Nebraska soils in relation to profile development. *Soil Science* 2:493-497.
- Archer, S., and J. K. Detling. 1984. The effects of defoliation and competition on regrowth of tillers of two North American mixed-grass prairie graminoids. *Oikos* 43:351-357.
- Bailey, V. 1926. A biological survey of North Dakota. United States Department of Agriculture, North American Fauna, 49.
- Baxter, F. P., and F. D. Hole. 1967. Ant pedoturbation in a prairie soil. *Soil Science Society of America Proceedings* 31:425-428.
- Biswell, H. H. 1973. Fire ecology in ponderosa pine-grassland. Tall Timbers Fire Ecology Conference Proceedings 12:69-96.
- Bohn, H. L., B. L. McNeal, and G. A. O'Connor. 1979. Soil Chemistry. Wiley-Interscience Publishers, New York. 329 p.
- Buntley, G. J., and R. I. Papendick. 1960. Worm-worked soils of eastern South Dakota, their morphology and classification. *Soil Science Society of America Proceedings* 24:128-132.
- Clark, T. W. 1979. The hard life of the prairie dog. *National Geographic* 156:270-281.
- Coppock, D. L., J. K. Detling, J. E. Ellis, and M. I. Dyer. 1983a. Plant-Herbivore interactions in a North American mixed grass prairie. I. Effects of Black-tailed prairie dogs on intraseasonal aboveground plant biomass and nutrient dynamics and plant species diversity. *Oecologia* 56:1-9.
- Coppock, D. L., J. E. Ellis, J. K. Detling, and M. I. Dyer. 1983b. Plant-Herbivore interactions in a North American mixed-grass prairie. II. Responses of Bison to modification of vegetation by prairie dogs. *Oecologia* 56:10-15.

- Costello, D. F. 1970. The World of the Prairie Dog. J. B. Lipencott Publishers, New York. 160 p.
- Cowan, J. A., G. S. Humphreys, P. B. Mitchell, and C. L. Murphy. 1985. An assessment of pedoturbation by two species of mound-building ants, Camponotus intrepides and Iridomyrmex purpureus. Australian Journal of Soil Research 22:95-107.
- Crocker-Bedford, D. C., and J. J. Spillett. 1977. Homeranges of Utah prairie dogs. Journal of Mammalogy 58:672-673.
- Dalstead, K. J., S. S. Blair, B. K. Worchester, and R. Klukas. 1981. Application of remote sensing to prairie dog management. Journal of Range Management 34:218-223.
- Darton, N. H., and S. Paige. 1925. Geologic Atlas of the United States, Central Black Hills Folio Number 19. United States Geological Survey, Washington, D. C.
- Detling, J. K., and E. L. Painter. Defoliation responses of western wheatgrass populations with diverse histories of prairie dog grazing. Oecologia 57:65-71.
- Dodge, R. I. 1876. The Black Hills. Ross and Haines Publishers, Minneapolis. Reprinted 1965. 155 p.
- Eidt, R. C. 1977. Detection and examination of anthrosols by phosphate analysis. Science 197:1327-1333.
- Eidt, R. C. 1984. Advances in Abandoned Settlement Analysis: Application to Prehistoric Anthrosols in Columbia, South America. University of Wisconsin-Milwaukee Press. 159 p.
- Feldman, R. M., and R. A. Heimlich. 1980. The Black Hills. Kendall-Hunt Publishers, Dubuque, Iowa. 190 p.
- Garrett, M. G. 1982. Dispersal of Prairie Dogs in Wind Cave National Park, South Dakota. M. S. Thesis, Iowa State University, Ames. 76 p.
- Gartner, F. R., and W. W. Thompson. 1973. Fire in the Black Hills forest-grass ecotone. Tall Timbers Fire Ecology Conference Proceedings 12:37-68.
- Gottermann, H. L. 1973. Natural phosphate sources in relation to phosphate budgets; A contribution to the understanding of eutrophication. Water Research 7:3-17.

- Grinnell, J. 1923. The burrowing rodents of California as agents in soil formation. *Journal of Mammalogy* 4:137-149.
- Gruell, G. E. 1983. Fire and Vegetative Trends in the Northern Rockies: Interpretations from 1871-1982 Photographs. United States Forest Service Technical Report INT-158. 117 p.
- Hansen, R. M., and M. J. Morris. 1968. Movement of rocks by northern pocket gophers. *Journal of Mammalogy* 49: 391-399.
- Hardy, R. 1945. The influence of types of soil upon the local distribution of some mammals in southwestern Utah. *Ecological Monographs* 15:71-108.
- Hole, F. D. 1961. A classification of pedoturbations and some other processes and factors of soil formation in relation to isotropism and anisotropism. *Soil Science* 91:375-377.
- Hole, F. D. 1981. Effects of animals on soil. *Geoderma* 25:75-112.
- Humphrey, R. R. 1958. The desert grassland, a history of vegetational change and an analysis of causes. *Botanical Review* 24:193-252.
- Ingham, R. E., and J. K. Detling. 1984. Plant-Herbivore interactions in a North American mixed-grass prairie. *Oecologia* 63:307-313.
- Jackson, M. L. 1958. *Soil Chemical Analysis*. Prentice-Hall Publishers, Englewood Cliffs, New Jersey. 498 p.
- Jenny, H. 1980. *The Soil Resource: Origin and Behavior*. Ecological Studies 37. Springer-Verlag Publishers, New York. 377 p.
- Joffe, J. S. 1949. *Pedology*. Pedology Publications, New Brunswick, New Jersey. 662 p.
- Johnson, H. N. 1949. A climatological review of the Black Hills. *South Dakota School of Mines and Technology, Black Hills Engineer* 29:3-35.
- Kalisz, P. J., and E. L. Stone. 1983. Soil Mixing by scarab beetles and pocket gophers in north-central Florida. *Soil Science Society of America Journal* 48: 169-172.

- Keller, M., P. Froelich, and P. Winham. 1984. Test excavations at sites 39JK63 and 39JK68, Pass Creek. White River Badlands Regional Research Report, Volume 4. 48 p.
- Kelso, L. H. 1939. Food Habits of Prairie Dogs. United States Department of Agriculture Circular Number 529. 15 p.
- King, J. A. 1955. Social Behavior, Social Organization, and Population Dynamics in a Black-tailed Prairie Dog Town in the Black Hills of South Dakota. Laboratory of Vertebrate Zoology, University of Michigan, Ann Arbor.
- Koford, C. B. 1958. Prairie dogs, whitefaces, and bluegrama. Wildlife Monographs 3:1-78.
- Laycock, W. A., and B. Z. Richardson. 1975. Long term effects of pocket gopher control on vegetation and soils of an alpine grassland. Journal of Range Management 28:458-462.
- Lee, K. E., and T. G. Wood. 1971. Termites and Soils. Academic Press, London. 250 p.
- Lipps, R. C., and L. Chesnin. 1950. The phosphorus status of some azonal, prairie, and Chernozem soils in eastern Nebraska. Soil Science Society of America Proceedings 15:329-333.
- Longhurst, W. 1944. Observations on the ecology of the Gunnison prairie dog in Colorado. Journal of Mammalogy 25:24-36.
- Lovaas, A. L. 1972. Report on the Prairie Dogs of Wind Cave National Park. Wind Cave National Park Annual Report. 18 p.
- Ludlow, W. 1875. Report of a Reconnaissance of the Black Hills of Dakota Made in 1874. United States Engineer Department, Government Printing Office, Washington, D. C. 121 p.
- Mausbach, M. J., B. R. Brasher, R. D. Yeck, and W. D. Nettleton. 1980. Variability of measured properties in morphologically matched pedons. Soil Science Society of America Journal 44:358-363.

- McIntosh, A. C. 1927. Some sources of the Black Hills flora. South Dakota School of Mines and Technology, Black Hills Engineer 15:38-44.
- McIntosh, A. C. 1949. A botanical survey of the Black Hills of Dakota. South Dakota School of Mines and Technology, Black Hills Engineer 28:3-74.
- McKelvey, V. E. 1973. Abundance and Distribution of Phosphorus in the Lithosphere. In Griffith, E. J., A. Beaton, J. M. Spencer, and D. T. Mitchell, editors. Environmental Phosphorus Handbook, New York.
- McLean, E. O. 1982. Soil pH and Lime Requirement. In Methods of Soil Analysis, Part 2, Chemical and Microbiological Properties, Agronomy Monographs 9 (2nd edition). American Society of Agronomy-Soil Science Society of America, Madison, Wisconsin. p. 199-209.
- Miller, M. A. 1948. Seasonal trends in burrowing of pocket gophers (Thomomys). Journal of Mammalogy 29: 38-44.
- Miller, R. S., and H. E. Bond. 1960. The summer burrowing activity of pocket gophers. Journal of Mammalogy 41: 469-475.
- Newton, H., and W. P. Jenny. 1880. Report of the Geology and Resources of the Black Hills of Dakota. United States Geological Survey, Government Printing Office, Washington, D. C. 566 p.
- Nye, P. H. 1955. Some soil-forming processes in the humid tropics. IV. The action of soil fauna. Journal of Soil Science 6:73-83.
- Nye, P. H., and M. H. Bertheux. 1957. The distribution of phosphorus in forest and savannah soils of the Gold Coast and its agricultural significance. Journal of Soil Science 49:141-159.
- Osborn, B. 1942. Prairie dogs in shinnery (oak scrub) savannah. Ecology 23:110-115.
- Osborn, B., and P. E. Allen. 1949. Vegetation of an abandoned prairie dog town in tall grass prairie. Ecology 30:322-332.

- Progulske, D. R. 1972. Passport to 1874. South Dakota State University Agricultural Experiment Station, Brookings. South Dakota Farm and Home Research 23: 17-35.
- Puecker, T. K. 1968. Johann George Kohl, a theoretical geographer of the 19th century. The Professional Geographer 20:247-250.
- Reid, V. H., R. M. Hansen, and A. L. Ward. 1966. Counting mounds and earth plugs to census mountain pocket gophers. Journal of Wildlife Management 30:327-334.
- Ruhe, R. V. 1983. Aspects of Holocene Pedology in the United States. In: Late Quaternary Environments of the United States, Volume 2, The Holocene. H. E. Wright, Jr., editor. University of Minnesota Press, Minneapolis. 277 p.
- Salem, M. Z., and F. D. Hole. 1968. Ant pedoturbation in a forest soil. Soil Science Society of America Proceedings 32:363-367.
- Schripsema, J. R. 1977. Ecological changes on pine-grassland burned in spring, late spring, and winter. M. S. Thesis, South Dakota State University, Brookings. 99 p.
- Sheets, R. G., R. L. Linder, and R. B. Dahlgren. 1971. Burrow systems of prairie dogs in South Dakota. Journal of Mammalogy 52:451-453.
- Smith, G. D., W. H. Allaway, and F. F. Riecken. 1950. Prairie Soils of the Upper Mississippi. In: Advances in Agronomy, Volume 2. Academic Press, Incorporated, New York. p. 157-205.
- Soil Conservation Service. 1972. Soil Survey Laboratory Methods and Procedures for Collecting Soil Samples. Soil Survey Investigations Report 1, United States Department of Agriculture. 63 p.
- Soil Survey Staff. 1975. Soil Taxonomy. Agricultural Handbook 436. Soil Conservation Service, United States Department of Agriculture. Government Printing Office, Washington, D. C. 754 p.
- Steel, R. G. D., and J. H. Torrie. 1960. Principles and Procedures of Statistics. McGraw-Hill Book Company, New York. 481 p.

- Steel, R. G. D., and J. H. Torrie. 1980. Principles and Procedures of Statistics: A Biometrical Approach. 2nd edition. McGraw-Hill Book Company, New York. 633 p.
- Summers, C. A., and R. L. Linder. 1978. Food habits of the black-tailed prairie dog in western South Dakota. Journal of Range Management 31:134-136.
- Taylor, W. P., and J. V. G. Loftfield. 1924. Damage to Range Grasses by the Zuni Prairie Dog. United States Department of Agriculture Bulletin 1227:1-15.
- Thilenius, J. F. 1971. Vascular Plants of the Black Hills of South Dakota and Adjacent Wyoming. United States Department of Agriculture Forest Research Paper RM-71. 43 p.
- Thilenius, J. F. 1972. Classification of Deer Habitat in the Ponderosa Pine Forest of the Black Hills, South Dakota. United States Department of Agriculture Forest Research Paper RM-91. 28 p.
- Thompson, W. W., and F. R. Gartner. 1971. Native forage response to clearing low quality ponderosa pine. Journal of Range Management 24:272-277.
- Thorp, J. 1949. Effects of certain animals that live in soils. Science Monthly 68:180-191.
- Tileston, J. V., and R. R. Lechleitner. 1966. Some comparisons of the black-tailed and white-tailed prairie dogs in north-central Colorado. American Midland Naturalist 75:292-316.
- United States Department of Agriculture: Forest Service. 1978. Prairie Dog Management: Final Environmental Statement. 188 p.
- United States Geological Survey. 1957. Wind Cave Quadrangle, Custer County, South Dakota. 7.5 minute series topographic map, revised 1979.
- Uresk, D. W. 1984. Black-tailed prairie dog food habits and forage relationships in western South Dakota. Journal of Range Management 37:325-329.
- Van Bruggen, T. 1976. The Vascular Plants of South Dakota. Iowa State University Press, Ames. 538 p.

- Waggamann, W. H. 1969. Phosphoric Acid, Phosphates and Phosphate Fertilizers, 2nd edition. New York. 453 p.
- Watanabe, F. J., and S. R. Olsen. 1975. Test of an ascorbic acid method for determining phosphorus in water and NaHCO_3 extracts of soil. Soil Science Society of America Proceedings 29:677-678.
- Westin, F. C., and D. D. Malo. 1978. Soils of South Dakota. South Dakota State University Agricultural Experiment Station-United States Department of Agriculture, Soil Conservation Service Bulletin 656. 118 p.
- White, E. M. 1960. A possible relationship of little bluestem distribution to soils. Journal of Range Management 14:243-247.
- White, E. M., and D. C. Carlson. 1984. Estimating soil mixing by rodents. Proceedings of the South Dakota Academy of Sciences 63:34-37.
- White, E. M., and F. G. Gartner. 1974. Phosphorus and iron contents of some Black Hills soils. Proceedings of the South Dakota Academy of Sciences 53:29-35.
- White, E. M., F. G. Gartner, and W. W. Thompson. 1974. Content and adsorption of phosphorus in Black Hills prairie and forest soils. Soil Science Society of America Proceedings 38:965-967.
- White, E. M., J. R. Johnson, and J. T. Nichols. 1969. Prairie-forest transition soils of the South Dakota Black Hills. Soil Science Society of America Proceedings 33:932-936.
- Wilding, L. P., N. E. Smeck, and G. F. Hall, editors. 1983. Pedogenesis and Soil Taxonomy. I. Concepts and Interactions. Elsevier Publishers, New York. 303 p.